



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			NDIDATE JMBER		

BIOLOGY 0610/61

Paper 6 Alternative to Practical

October/November 2011

1 hour

Candidates answer on the Question Paper

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use				
1				
2				
3				
Total				

This document consists of 13 printed pages and 3 blank pages.



BLANK PAGE

1 Catalase is an enzyme found in plant and animal tissues. It catalyses the breakdown of hydrogen peroxide into water and oxygen. For Examiner's Use

$$2H_2O_2 \rightarrow 2H_2O + O_2$$

hydrogen peroxide water oxygen

The activity of this enzyme can be measured by collecting the volume of oxygen gas given off as shown in Fig. 1.1.




Fig. 1.1

Some students compared the catalase activity in two plant tissues, **sweet potato**, *Ipomoea batanus*, and **Irish potato**, *Solanum tuberosum*.

- 2.0 g of **sweet potato** was cut into small pieces.
- The small pieces were placed in a flask together with 25 cm³ of hydrogen peroxide.
- The bung and delivery tube were fitted to the flask, as shown in Fig. 1.1.
- The volume of oxygen gas released was measured after 4 minutes (experiment 1).
- This was repeated three times (experiments 2, 3 and 4).
- The same procedure was carried out with 2.0 g of Irish potato cut into small pieces.
- The results are shown in Table 1.1.

Table 1.1

For Examiner's Use

ovnoriment	volume of oxygen gas / cm ³						
experiment	sweet potato	Irish potato					
1	32.0	12.5					
2	20.0	9.0					
3	35.5	8.5					
4	28.0	10.0					
total	115.5						
mean	28.9						

(a) (i)	The total volume of oxygen gas and the mean volume of oxygen gas have bee calculated for the sweet potato .	en
	Calculate these values for the Irish potato .	

Show your working below.

	Write your answers in Table 1.1.	[2]
(ii)	Suggest why the tissues were cut into small pieces before being added to flask.	the
		[1]

	Draw	a hori	zonta	l line	acros	s yo	ur baı	r cha	art to	show	the m	ean va	lue.		
									7-7-1-1						
		+++++		1-1-1-1		 - - -									
					+	!-! -!-!-									
						- - - - - - - -			‡- 						
				1-1-1-1	+++++				<u> </u>						
		-1-1-1-1-				I- I -I-I-									
				1111											
						!-! -!-									
					<u> </u>										
						! ! ! ! !									
				1.1.1.1	; 	!-! -!-!-									
						<u> </u>									
						1- 1 -1-1-									
		-1-1-1-1-													
						!-! -!-									
						<u> </u>									
		-1-1-1-				!-! -!-!-									
	ì							-1-1-1-	111111						
(ii)	Sugge	est tw	n res	sons	s for t	he v	/ariati	on i	n th	ne resi	ılts of	the fo	ur sw	eet i	nota
("')	experi			13011	3 101 1		anan	011 1		10 1030	1113 01	tile ic	ui 34	וייייייי	pota

For Examiner's Use

[2]

(c)	Suggest and explain two ways in which a similar investigation could be planned to collect more reliable data.	Exa
	[4]	
	[Total: 14]	

For Examiner's Use

BLANK PAGE

2 Birds have feathers to cover their bodies.

Fig. 2.1 shows two types of feather, **A** and **B**.



(a) (i) Make a labelled outline drawing of feather A.

For Examiner's Use

PMT

[4]

(ii)	Suggest a function of feather A.	For Examiner's Use
	[1]	
(iii)	Suggest a function of feather B and explain how it is adapted for this function.	
	[2]	

Question 2 continues on the next page.

Fig. 2.2 shows feather ${\bf B}$ placed upon a grid. You will use this to help you to estimate the surface area of feather ${\bf B}$.

For Examiner's Use




Fig. 2.2

(b) (i) Measure the size of one of the grid squares. Use this to calculate the surface area of feather **B**.

Show your working.

area of feather **B** ____cm² [3]

For Examin Use	escribe a more accurate method that you could use to find the actual surface ea of feather B .	(ii)
	[2]	
	[Total: 12]	

ner's

3 Fig. 3.1 shows three cress seedlings grown under different conditions. The seeds came from the same plant and the seedlings have been grown for the same length of time.

For Examiner's Use

The seedlings are drawn to scale.




Fig. 3.1

(a) The cress seedlings, **D**, **E** and **F** differ in appearance. For each seedling state how it is different and suggest an explanation.

(i)	seedling D
	[3]
(ii)	seedling E
	[3]

For Examiner's Use

(iii)	seedling F	
	[3]	

Question 3 continues on the next page.

(b) Seeds develop within the fruits of flowering plants.Fig. 3.2 shows two types of fruit, G and H, from species similar to cress.

For Examiner's Use




Fig. 3.2

t H.

1	
2	
	[2]

(ii) Complete Table 3.1 by recording two differences between fruit **G** and fruit **H**.

Table 3.1

fruit G	fruit H
1	
2	

[2]

Suggest how the seeds may be dispersed from these two fruits.		For Examiner's
	ı	Use
[1]	
	Suggest how the seeds may be dispersed from these two fruits.	Suggest how the seeds may be dispersed from these two fruits. [1]

[Total: 14]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.