


1.



The figure above shows a sketch of part of the curve  $C$  with equation

$$y = \sin(\ln x), \quad x \geq 1.$$

The point  $Q$ , on  $C$ , is a maximum.

- (a) Show that the point  $P(1, 0)$  lies on  $C$ . (1)
- (b) Find the coordinates of the point  $Q$ . (5)
- (c) Find the area of the shaded region between  $C$  and the line  $PQ$ . (9)

**(Total 15 marks)**

1. (a)  $x = 1; y = \sin(\ln 1) = \sin 0 = 0$   
 $\therefore P = (1, 0)$  and P lies on C B1 c.s.o. 1

(b)  $y' = \frac{1}{x} \cos(\ln x)$  M1, A1


$y' = 0$  at Q  $\therefore \cos(\ln x) = 0 \therefore \ln x = \frac{\pi}{2}$  M1

$$x = e^{\frac{\pi}{2}}$$

$\therefore Q = \left( e^{\frac{\pi}{2}}, \sin(\ln e^{\frac{\pi}{2}}) \right)$  A1

$= (e^{\frac{\pi}{2}}, 1)$  A1 5

(c)



Area =  $\int_1^{e^{\frac{\pi}{2}}} \sin(\ln x) dx$  - Area  $\Delta PQR$  (correct approach) M1

Area  $\Delta PQR = \frac{1}{2} \times 1 \times (e^{\frac{\pi}{2}} - 1)$  B1

for integral; let  $\ln x = u \therefore x = e^u$  (substitution) M1

$$\frac{1}{x} dx = du \therefore dx = e^u du$$

$\underline{E} = \int_0^{\frac{\pi}{2}} \sin u \cdot (e^u du)$  A1

$= \left[ e^u \sin u \right]_0^{\frac{\pi}{2}} - \int e^u \cos u du$  M1

$= e^{\frac{\pi}{2}} - \left[ e^u \cos u \right]_0^{\frac{\pi}{2}} - \int e^u \sin u du$  M1

$$\therefore 2I = e^{\frac{\pi}{2}} + 1$$

$$I = \frac{1}{2}(1 + e^{\frac{\pi}{2}}) = 1$$

(I) A1

$$\therefore \text{Area} = \frac{1}{2}(1 + e^{\frac{\pi}{2}}) - \frac{1}{2}(-1 + e^{\frac{\pi}{2}}) = 1$$

A1

**[9]**

1. This was the question in which many candidates earned their highest marks. It was also the one for which most 5 marks were gained. Virtually all candidates scored the first mark. Differentiation was generally good in part (b) and many candidates scored all 5 of these marks. A common error was to state that  $\ln x=1$ . There were also many good attempts at part (c). Nearly all recognized the need to take the difference of two areas. Those who sought to find the area of the triangle by forming the equation of the line and then integrating usually came unstuck in a mass of algebra and they rarely obtained the correct value. Fortunately most simply used half the base  $\times$  height! Integration of  $y$  was usually well done. Similar numbers of candidates used direct integration by parts ( $x\sin(\ln x)$  etc.) as used the substitution  $u=\ln x$ , resulting in  $e^u \sin u \, du$ . Many were able to complete the two cycles of parts and obtain the correct answer.