

Centre No.						Paper Reference	Surname	Initial(s)
Candidate No.						6 6 6 6 / 0 1	Signature	

Paper Reference(s)
6666/01

Edexcel GCE

Core Mathematics C4

Advanced

Tuesday 22 January 2008 – Afternoon
Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

Materials required for examination
Mathematical Formulae (Green)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
Total	

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature. Check that you have the correct question paper. You must write your answer for each question in the space following the question. When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided. Full marks may be obtained for answers to ALL questions. The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). There are 8 questions in this question paper. The total mark for this paper is 75. There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.
©2008 Edexcel Limited.

Printer's Log. No.
N26282A



Turn over

1.




Figure 1

The curve shown in Figure 1 has equation $y = e^x \sqrt{\sin x}$, $0 \leq x \leq \pi$. The finite region R bounded by the curve and the x -axis is shown shaded in Figure 1.

(a) Complete the table below with the values of y corresponding to $x = \frac{\pi}{4}$ and $\frac{\pi}{2}$, giving your answers to 5 decimal places.

x	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π
y	0			8.87207	0

(2)

(b) Use the trapezium rule, with all the values in the completed table, to obtain an estimate for the area of the region R . Give your answer to 4 decimal places.

(4)



6. The points A and B have position vectors $2\mathbf{i} + 6\mathbf{j} - \mathbf{k}$ and $3\mathbf{i} + 4\mathbf{j} + \mathbf{k}$ respectively.

The line l_1 passes through the points A and B .

- (a) Find the vector \overrightarrow{AB} . (2)

- (b) Find a vector equation for the line l_1 . (2)

A second line l_2 passes through the origin and is parallel to the vector $\mathbf{i} + \mathbf{k}$. The line l_1 meets the line l_2 at the point C .

- (c) Find the acute angle between l_1 and l_2 . (3)

- (d) Find the position vector of the point C . (4)



7.




Figure 3

The curve C has parametric equations

$$x = \ln(t + 2), \quad y = \frac{1}{(t + 1)}, \quad t > -1.$$

The finite region R between the curve C and the x -axis, bounded by the lines with equations $x = \ln 2$ and $x = \ln 4$, is shown shaded in Figure 3.

(a) Show that the area of R is given by the integral

$$\int_0^2 \frac{1}{(t + 1)(t + 2)} dt. \tag{4}$$

(b) Hence find an exact value for this area. (6)

(c) Find a cartesian equation of the curve C , in the form $y = f(x)$. (4)

(d) State the domain of values for x for this curve. (1)



8. Liquid is pouring into a large vertical circular cylinder at a constant rate of $1600 \text{ cm}^3 \text{ s}^{-1}$ and is leaking out of a hole in the base, at a rate proportional to the square root of the height of the liquid already in the cylinder. The area of the circular cross section of the cylinder is 4000 cm^2 .

(a) Show that at time t seconds, the height h cm of liquid in the cylinder satisfies the differential equation

$$\frac{dh}{dt} = 0.4 - k\sqrt{h}, \text{ where } k \text{ is a positive constant.} \quad (3)$$

When $h = 25$, water is leaking out of the hole at $400 \text{ cm}^3 \text{ s}^{-1}$.

(b) Show that $k = 0.02$ (1)

(c) Separate the variables of the differential equation

$$\frac{dh}{dt} = 0.4 - 0.02\sqrt{h},$$

to show that the time taken to fill the cylinder from empty to a height of 100 cm is given by

$$\int_0^{100} \frac{50}{20 - \sqrt{h}} dh. \quad (2)$$

Using the substitution $h = (20 - x)^2$, or otherwise,

(d) find the exact value of $\int_0^{100} \frac{50}{20 - \sqrt{h}} dh$. (6)

(e) Hence find the time taken to fill the cylinder from empty to a height of 100 cm, giving your answer in minutes and seconds to the nearest second. (1)



Leave
blank

Question 8 continued

Lined area for writing answers to Question 8.

Q8

(Total 13 marks)

TOTAL FOR PAPER: 75 MARKS

END

