




1.

Figure 1



A particle of weight 24 N is held in equilibrium by two light inextensible strings. One string is horizontal. The other string is inclined at an angle of  $30^\circ$  to the horizontal, as shown in Figure 1. The tension in the horizontal string is  $Q$  newtons and the tension in the other string is  $P$  newtons. Find

(a) the value of  $P$ , (3)

(b) the value of  $Q$ . (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---



2.

**Figure 2**



A uniform plank  $AB$  has weight 120 N and length 3 m. The plank rests horizontally in equilibrium on two smooth supports  $C$  and  $D$ , where  $AC = 1$  m and  $CD = x$  m, as shown in Figure 2. The reaction of the support on the plank at  $D$  has magnitude 80 N. Modelling the plank as a rod,

- (a) show that  $x = 0.75$  **(3)**

A rock is now placed at  $B$  and the plank is on the point of tilting about  $D$ . Modelling the rock as a particle, find

- (b) the weight of the rock, **(4)**

- (c) the magnitude of the reaction of the support on the plank at  $D$ . **(2)**

- (d) State how you have used the model of the rock as a particle. **(1)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave  
blank

**Question 2 continued**

Lined area for writing the answer to Question 2, containing approximately 30 horizontal lines.

**Q2**

**(Total 10 marks)**

--	--



N 2 3 5 6 0 A 0 5 1 6



Leave blank

4. A particle  $P$  of mass  $0.3 \text{ kg}$  is moving with speed  $u \text{ m s}^{-1}$  in a straight line on a smooth horizontal table. The particle  $P$  collides directly with a particle  $Q$  of mass  $0.6 \text{ kg}$ , which is at rest on the table. Immediately after the particles collide,  $P$  has speed  $2 \text{ m s}^{-1}$  and  $Q$  has speed  $5 \text{ m s}^{-1}$ . The direction of motion of  $P$  is reversed by the collision. Find

(a) the value of  $u$ , (4)

(b) the magnitude of the impulse exerted by  $P$  on  $Q$ . (2)

Immediately after the collision, a constant force of magnitude  $R \text{ newtons}$  is applied to  $Q$  in the direction directly opposite to the direction of motion of  $Q$ . As a result  $Q$  is brought to rest in  $1.5 \text{ s}$ .

(c) Find the value of  $R$ . (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---












6.

Figure 3



A box of mass  $30\text{ kg}$  is being pulled along rough horizontal ground at a constant speed using a rope. The rope makes an angle of  $20^\circ$  with the ground, as shown in Figure 3. The coefficient of friction between the box and the ground is  $0.4$ . The box is modelled as a particle and the rope as a light, inextensible string. The tension in the rope is  $P$  newtons.

(a) Find the value of  $P$ . **(8)**

The tension in the rope is now increased to  $150\text{ N}$ .

(b) Find the acceleration of the box. **(6)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave  
blank

**Question 6 continued**

Lined writing area for the answer to Question 6. The page contains approximately 30 horizontal lines for writing.

**(Total 14 marks)**

**Q6**

--	--



7.

Figure 4




Figure 4 shows two particles  $P$  and  $Q$ , of mass 3 kg and 2 kg respectively, connected by a light inextensible string. Initially  $P$  is held at rest on a fixed smooth plane inclined at  $30^\circ$  to the horizontal. The string passes over a small smooth light pulley  $A$  fixed at the top of the plane. The part of the string from  $P$  to  $A$  is parallel to a line of greatest slope of the plane. The particle  $Q$  hangs freely below  $A$ . The system is released from rest with the string taut.

(a) Write down an equation of motion for  $P$  and an equation of motion for  $Q$ . (4)

(b) Hence show that the acceleration of  $Q$  is  $0.98 \text{ m s}^{-2}$ . (2)

(c) Find the tension in the string. (2)

(d) State where in your calculations you have used the information that the string is inextensible. (1)

On release,  $Q$  is at a height of 0.8 m above the ground. When  $Q$  reaches the ground, it is brought to rest immediately by the impact with the ground and does not rebound. The initial distance of  $P$  from  $A$  is such that in the subsequent motion  $P$  does not reach  $A$ . Find

(e) the speed of  $Q$  as it reaches the ground, (2)

(f) the time between the instant when  $Q$  reaches the ground and the instant when the string becomes taut again. (5)

---

---

---

---

---

---

---

---



Leave  
blank

**Question 7 continued**

Horizontal lines for writing answers.

**Q7**

Grading box

**(Total 16 marks)**

**TOTAL FOR PAPER: 75 MARKS**

**END**

