

3. Three forces \mathbf{F}_1 , \mathbf{F}_2 and \mathbf{F}_3 acting on a particle P are given by

$$\mathbf{F}_1 = (7\mathbf{i} - 9\mathbf{j}) \text{ N}$$

$$\mathbf{F}_2 = (5\mathbf{i} + 6\mathbf{j}) \text{ N}$$

$$\mathbf{F}_3 = (p\mathbf{i} + q\mathbf{j}) \text{ N}$$

where p and q are constants.

Given that P is in equilibrium,

(a) find the value of p and the value of q .

(3)

The force \mathbf{F}_3 is now removed. The resultant of \mathbf{F}_1 and \mathbf{F}_2 is \mathbf{R} . Find

(b) the magnitude of \mathbf{R} ,

(2)

(c) the angle, to the nearest degree, that the direction of \mathbf{R} makes with \mathbf{j} .

(3)



8.




Figure 2

A particle P of mass 4 kg is moving up a fixed rough plane at a constant speed of 16 m s^{-1} under the action of a force of magnitude 36 N . The plane is inclined at 30° to the horizontal. The force acts in the vertical plane containing the line of greatest slope of the plane through P , and acts at 30° to the inclined plane, as shown in Figure 2. The coefficient of friction between P and the plane is μ . Find

- (a) the magnitude of the normal reaction between P and the plane, **(4)**
- (b) the value of μ . **(5)**

The force of magnitude 36 N is removed.

- (c) Find the distance that P travels between the instant when the force is removed and the instant when it comes to rest. **(5)**



