




Question Number	Scheme	Marks
1. (a)	$s = ut + \frac{1}{2}at^2 : 50 = 5 \times 4 + \frac{1}{2} \times a \times 4^2$ $\Rightarrow 30 = 8a \Rightarrow a = 3.75 \text{ m s}^{-1}$	M1 A1 A1 (3)
(b)	$30^2 = 5^2 + 2 \times 3.75 \times s$ $\Rightarrow s = 116\frac{2}{3} \text{ m}$	M1 A1 ft A1 (3) (6 marks)
2.	<p>Considering momentum of A: $3.6 = 0.5(5 + v)$ $\Rightarrow v = 2.2 \text{ m s}^{-1}$</p> <p>Considering momentum of B: $3.6 = m(3 + 1)$ or $m(3 - 1)$ $m = 0.9$ or $m = 1.8$</p>	M1 A1 A1 (3) M1 A1 (one) M1 A1 (both) (4) (7 marks)
3. (a)	<p>$M(C): 16 \times 30 = w \times 20 + 5 \times 70$ (3 terms) $\Rightarrow w = 6.5 \text{ N}$</p>	M1 A1 A1 (3)
(b)	<p>$M(D): 3.5d + 6.5(d - 50) = 5(100 - d)$ $\Rightarrow d = 55 \text{ cm}$</p>	M1 A2ft (-1 eeo) A1 (4)
(c)	Tension equal along string, i.e. tensions = weights throughout or no contributions from strings in moments equation	B1 (1) (8 marks)

(ft = follow through mark; -1eeoo = minus one mark for each error or omission)

Question Number	Scheme	Marks
4. (a)	 $F = \frac{2}{5}R$ $R(\uparrow): R \cos 30^\circ - F \cos 60^\circ = 6g$ $R \frac{\sqrt{3}}{2} - \frac{2}{5}R - \frac{1}{2} = 6g$ $\Rightarrow R = 88.3 \text{ N (or } 88 \text{ N)}$	B1 M1 A1 A1 (4)
(b)	$R(\leftarrow): P = R \cos 60^\circ + F \cos 30^\circ$ $= 74.7 \text{ N (or } 75 \text{ N)}$	M1 A1 A1 (3)
(c)	 <p>Component of weight (\checkmark) $= 6g \cos 60^\circ$ $= 29.4 \text{ N}$</p> <p>$R' = 6g \cos 30^\circ = 50.9 \text{ N}$</p> <p>$F_{\max} = 0.4 R' = 20.36 \text{ N}$</p> <p>Since $29.4 > 20.36$, the box moves</p>	B1 M1 A1 M1 A1 cso (5) (12 marks)
5. (a)	 $\tan \theta = \frac{1}{2} \Rightarrow \theta = 26.6^\circ$ <p>angle required $= 153.4^\circ$</p>	M1 A1 A1 (3)
(b)	$\mathbf{a} = \frac{1}{3} [(\mathbf{i} - 2\mathbf{j}) - (-5\mathbf{i} + 7\mathbf{j})]$ $= (2\mathbf{i} - 3\mathbf{j}) \text{ m s}^{-2}$	M1 A1 (2)
(c)	$\mathbf{F} = m\mathbf{a} = 4\mathbf{i} - 6\mathbf{j}$ $ \mathbf{F} = \sqrt{(16 + 36)} = 7.21 \text{ N}$	M1 M1 A1 (3)
(d)	$\mathbf{v} = (-5 + 2t)\mathbf{i} + (7 - 3t)\mathbf{j}$	M1 A1ft (2)
(e)	$\mathbf{v} \text{ parallel to } \mathbf{i} + \mathbf{j} \Rightarrow \frac{-5 + 2t}{7 - 3t} = 1$ $\Rightarrow t = 2.4 \text{ s}$	M1 M1 A1 (3) (13 marks)

(cso = correct solution only)

Question Number	Scheme	Marks
6. (a)	<p>shape (3, 2.5)</p>	B1 B1 (2)
(b)	$\text{Area} = 27 = \frac{1}{2} \times 1.5 \times 3 + 3T + \frac{1}{2} \times 2.5 \times 3$ $\Rightarrow T = 7 \text{ s}$	M1 A1 A1 (3)
(c)	<p>shape $0 \leq t \leq 8.5$ shape $t > 8.5$ (2, 7 (ft), 2.5)</p>	B1 B1 B1 (3)
(d)	<p>(System)</p> $T - 200g = 200 \times 2$ $\Rightarrow T = 2360 \text{ N}$	M1 A1 A1 (3)
(e)	<p>(Man)</p> $R - 80g = -80 \times 1.2$ $\Rightarrow R = 688 \text{ N}$	M1 A1 A1 (3)
		(14 marks)

Question Number	Scheme	Marks
7. (a)		
	$R = 2mg \Rightarrow F = 2\mu mg$	B1
	A: $T - 2\mu mg = 2ma$	M1 A1
	B: $mg \times \frac{1}{2} - T = ma$	M1 A1
	Eliminating T : $3ma = \frac{1}{2}mg - 2\mu mg$	M1
	$a = \frac{1}{6}(1 - 4\mu)g$ (*)	A1 (7)
(b)	$\mu = 0.2 \Rightarrow a = \frac{1}{30}g$	B1
	when string breaks: $v^2 = 2 \times \frac{1}{30}g \times h = \frac{1}{15}gh$	M1 A1
	A decelerating with deceleration $f \Rightarrow 2mf = 2\mu mg$	
	$f = \mu g = \frac{1}{5}g$	B1
	Hence distance travelled during deceleration is given by $\frac{1}{15}gh = 2 \times \frac{1}{5}gd$	M1
	$\Rightarrow d = \frac{1}{6}h$	
	\therefore Total distance = $\frac{7}{6}h$	A1 cso (6)
(c)	Any two from: weight of pulley; friction at pulley; friction on slope; weight of string; string extensible; 'spin' of particle	B1 B1 (2)
		(15 marks)

((*)) indicates final line is given on the paper; cso = correct solution only)