

3.




Figure 1

A particle of mass 2 kg is suspended from a horizontal ceiling by two light inextensible strings, PR and QR . The particle hangs at R in equilibrium, with the strings in a vertical plane. The string PR is inclined at 55° to the horizontal and the string QR is inclined at 35° to the horizontal, as shown in Figure 1.

Find

- (i) the tension in the string PR ,
- (ii) the tension in the string QR .

(7)



4.




Figure 2

A lift of mass 200 kg is being lowered into a mineshaft by a vertical cable attached to the top of the lift. A crate of mass 55 kg is on the floor inside the lift, as shown in Figure 2. The lift descends vertically with constant acceleration. There is a constant upwards resistance of magnitude 150 N on the lift. The crate experiences a constant normal reaction of magnitude 473 N from the floor of the lift.

(a) Find the acceleration of the lift. **(3)**

(b) Find the magnitude of the force exerted on the lift by the cable. **(4)**



5.




Figure 3

A beam AB has length 5 m and mass 25 kg. The beam is suspended in equilibrium in a horizontal position by two vertical ropes. One rope is attached to the beam at A and the other rope is attached to the point C on the beam where $CB = 0.5$ m, as shown in Figure 3. A particle P of mass 60 kg is attached to the beam at B and the beam remains in equilibrium in a horizontal position. The beam is modelled as a uniform rod and the ropes are modelled as light strings.

(a) Find

- (i) the tension in the rope attached to the beam at A ,
- (ii) the tension in the rope attached to the beam at C .

(6)

Particle P is removed and replaced by a particle Q of mass M kg at B . Given that the beam remains in equilibrium in a horizontal position,

(b) find

- (i) the greatest possible value of M ,
- (ii) the greatest possible tension in the rope attached to the beam at C .

(6)



Leave
blank

Question 5 continued

Lined area for writing the answer to Question 5 continued.



6. A particle P is moving with constant velocity. The position vector of P at time t seconds ($t \geq 0$) is \mathbf{r} metres, relative to a fixed origin O , and is given by

$$\mathbf{r} = (2t - 3)\mathbf{i} + (4 - 5t)\mathbf{j}$$

(a) Find the initial position vector of P . (1)

The particle P passes through the point with position vector $(3.4\mathbf{i} - 12\mathbf{j})$ m at time T seconds.

(b) Find the value of T . (3)

(c) Find the speed of P . (4)

Horizontal lines for writing answers.



7. A train travels along a straight horizontal track between two stations, A and B . The train starts from rest at A and moves with constant acceleration 0.5 m s^{-2} until it reaches a speed of $V \text{ m s}^{-1}$, ($V < 50$). The train then travels at this constant speed before it moves with constant deceleration 0.25 m s^{-2} until it comes to rest at B .

(a) Sketch in the space below a speed-time graph for the motion of the train between the two stations A and B .

(2)

The total time for the journey from A to B is 5 minutes.

(b) Find, in terms of V , the length of time, in seconds, for which the train is

(i) accelerating,

(ii) decelerating,

(iii) moving with constant speed.

(5)

Given that the distance between the two stations A and B is 6.3 km,

(c) find the value of V .

(6)



8.




Figure 4

Two particles P and Q have mass 4 kg and 0.5 kg respectively. The particles are attached to the ends of a light inextensible string. Particle P is held at rest on a fixed rough plane, which is inclined to the horizontal at an angle α where $\tan \alpha = \frac{4}{3}$. The coefficient of friction between P and the plane is 0.5. The string lies along the plane and passes over a small smooth light pulley which is fixed at the top of the plane. Particle Q hangs freely at rest vertically below the pulley. The string lies in the vertical plane which contains the pulley and a line of greatest slope of the inclined plane, as shown in Figure 4. Particle P is released from rest with the string taut and slides down the plane.

Given that Q has not hit the pulley, find

- (a) the tension in the string during the motion, (11)
- (b) the magnitude of the resultant force exerted by the string on the pulley. (4)



