

## MECHANICS (C) UNIT 2 TEST PAPER 6

Take  $g = 9.8 \text{ ms}^{-2}$  and give all answers correct to 3 significant figures where necessary.

1. A ship, of mass 5000 tonnes, is moving through the sea at a constant speed of  $15 \text{ km h}^{-1}$ .
- (i) Calculate the momentum of the ship, in the form  $a \times 10^n$ , where  $0 < a < 10$  and  $n$  is an integer. State the units of your answer. [2]

Given that there is a constant force of magnitude 4000 N acting against the ship's motion due to air and water resistances,

- (ii) find the rate, in kW, at which the ship's engines are working. [3]

2. Two small smooth spheres  $P$  and  $Q$  are moving along a straight line in opposite directions with the same speed  $u$ , and collide directly. Immediately after the impact, the direction of  $P$ 's motion has been reversed and its speed has been halved.

The coefficient of restitution between  $P$  and  $Q$  is  $e$ .

- (i) Express the speed of  $Q$  after the impact in the form  $au(be + c)$ , where  $a$ ,  $b$  and  $c$  are constants to be found. [4]


- (ii) Deduce the range of values of  $e$  for which the direction of motion of  $Q$  remains unaltered. [3]

3. A small block of wood, of mass 0.5 kg, slides down a line of greatest slope of a smooth plane inclined at an angle  $\alpha$  to the horizontal, where  $\sin \alpha = \frac{2}{5}$ . The block is given an initial impulse of magnitude 2 Ns, and reaches the bottom of the plane with kinetic energy 19 J.

- (i) Find, in J, the change in the potential energy of the block as it moves down the plane. [2]

- (ii) Hence find the distance travelled by the block down the plane. [3]


- (iii) State two modelling assumptions that you have made. [2]

4.  A uniform rod  $XY$ , of length  $2a$  and mass  $m$ , is connected to a vertical wall by a smooth hinge at the end  $X$ . A horizontal light inelastic string connects the mid-point of  $XY$  to the wall and the rod is in equilibrium in this position.

- (i) Draw a diagram to show all the forces acting on the rod. [3]

Given that the tension in the horizontal string is of magnitude  $2mg$ ,

- (ii) find the angle which  $XY$  makes with the vertical. [5]


5.  The diagram shows a uniform lamina  $ABCDEF$ .

- (i) Calculate the distance of the centre of mass of the lamina from (a)  $AF$ , (b)  $AB$ . [6]

The lamina is hung over a smooth peg at  $D$  and rests in equilibrium in a vertical plane.

- (ii) Find the angle between  $CD$  and the vertical. [3]

6. The diagram shows a particle  $P$  of mass  $m$  kg moving on the inner surface of a smooth fixed hemispherical bowl of radius  $r$  m which is fixed with its axis vertical.  $P$  moves at a constant speed in a horizontal circle, at a depth  $h$  m below the top of the bowl.



- (i) Show that the force exerted on  $P$  by the bowl has magnitude  $\frac{mgr}{h}$  N. [4]

- (ii) Find, in terms of  $g$ ,  $h$  and  $r$ , the constant speed of  $P$ . [5]

7. A projectile is fired with speed  $8 \text{ ms}^{-1}$  from a point  $O$  on a horizontal plane, at an angle of elevation  $\alpha$ . It moves under gravity such that, at time  $t$  seconds after projection, it is at the point with coordinates  $(x, 4t - 4.9t^2)$  relative to  $O$ , where the  $x$  and  $y$  directions are respectively horizontal and vertically upwards.
- (i) Show that  $\sin \alpha = 0.5$  [3]
- (ii) Hence find  $x$  in terms of  $t$ . [2]
- (iii) Show that  $y = \frac{x}{\sqrt{3}} - \frac{gx^2}{96}$  [3]
- (iv) Deduce the value of  $x$  when the projectile returns to the horizontal plane. [2]
- On another occasion, the equation of the projectile's path is  $y = \frac{3x}{4} - \frac{gx^2}{32}$ .
- (v) Find the angle of projection and the initial speed of the projectile. [5]

### MECHANICS 2 (C) TEST PAPER 6 : ANSWERS AND MARK SCHEME

1. (i)  $5\,000\,000 \times (15\,000 \div 3600) = 2.08 \times 10^7 \text{ Ns or kg m s}^{-1}$  M1 A1  
 (ii)  $P = 4000 \times (150 \div 36) = 16.7 \text{ kW}$  M1 A1 A1 5
2. (i)  $[v - (-u/2)] / (-u - u) = -e$  M1 A1  
 $v + \frac{1}{2}u = 2ue$  Speed =  $\pm \frac{1}{2}u(4e - 1)$  M1 A1  
 (ii) If  $v < 0$  then  $4e - 1 < 0$ , so  $0 < e < \frac{1}{4}$  M1 A1 A1 7
3. (i) P.E. lost = K.E. gained =  $19 - \frac{1}{4} \times 0.5 \times 4^2 = 15 \text{ J}$  M1 A1  
 (ii)  $0.5gh = 15$   $h = \frac{30}{g}$   $d = h \div \sin \alpha = \frac{75}{g} = 7.65 \text{ m}$  M1 A1 A1  
 (iii) Modelled block as particle; ignored air resistance B1 B1 7
4. (i) Diagram showing weight, tension, two components of reaction or single reaction force at  $X$  B3  
 (ii)  $M(X) : 2mg a \cos \theta = mg a \sin \theta$   $\tan \theta = 2$   $\theta = 63.4^\circ$  M1 A1 A1 M1 A1 8
5. (i) (a)  $132(11) + 84(3) = 216\bar{x}$   $\bar{x} = 7.89$  M1 A1 A1  
 (b)  $132(3) + 84(13) = 216\bar{y}$   $\bar{y} = 6.89$  M1 A1 A1  
 (ii)  $\tan \alpha = 0.89 \div 1.89 = 0.471$   $\alpha = 25.2^\circ$  M1 A1 A1 9
6. (i) Reaction  $R$  acts on  $P$  towards centre of sphere, at  $\theta$  to vertical M1  
 where  $\cos \theta = h/r$   $R \cos \theta = mg$ , so  $R = \frac{mgr}{h}$  B1 M1 A1  
 (ii) Resolve towards centre :  $R \sin \theta = mv^2 / (r \sin \theta)$  B1  
 $v^2 = (mgr^2 / h)(\sin^2 \theta / m) = (gr^2 / h)(r^2 - h^2) / r^2$  M1 A1 A1  
 $v = \sqrt{[g(r^2 - h^2) / h]}$  A1 9
7. (i)  $y = (8 \sin \alpha)t - 4.9t^2 = 4t - 4.9t^2$  (given), so  $\sin \alpha = \frac{1}{2}$  M1 A1 A1  
 (ii)  $x = 8t \cos \alpha = (4\sqrt{3})t$  M1 A1

$$(iii) y = \frac{4x}{4\sqrt{3}} - \frac{gx^2}{2(4\sqrt{3})^2} = \frac{x}{\sqrt{3}} - \frac{gx^2}{96} \quad \text{M1 A1 A1}$$

$$(iv) \text{ When } y = 0, \frac{x}{\sqrt{3}} = \frac{gx^2}{96} \quad gx = \frac{96}{\sqrt{3}} \quad x = 5.66 \quad \text{M1 A1}$$

$$(v) \tan \alpha = 0.75 \quad \alpha = 36.9^\circ \quad \text{M1 A1}$$

$$2u^2 \cos^2 \alpha = 32 \quad u^2 \left( \frac{16}{25} \right) = 16 \quad u = 5 \text{ ms}^{-1} \quad \text{M1 A1 A1} \quad 15$$