

AQA Maths M2

Topic Questions from Papers

Differential Equations

Answers

<b>1</b>	$1600 \frac{dv}{dt} = -40v$ $\int \frac{1}{v} dv = \int -\frac{1}{40} dt$ $\ln v = -\frac{t}{40} + c$ $v = Ae^{-\frac{t}{40}}$ $t = 0, v = 20 \Rightarrow c = 20$ $v = 20e^{-\frac{t}{40}}$	M1		Applying Newton's second law with $40v$ and $\frac{dv}{dt}$ .
		A1		Correct equation
		dM1		Separating variables
		dM1		integrating to get $\ln v$ term.
		A1		Correct integral with or without $c$
dM1		Finding constant		
A1		7	Correct final result	
<b>Total</b>			<b>7</b>	

(Q5, Jan 2006)

<b>2 (a)</b>	$20 \frac{dv}{dt} = -10\sqrt{v}$ $\frac{dv}{dt} = -\frac{\sqrt{v}}{2}$ $\int \frac{1}{\sqrt{v}} dv = \int -\frac{1}{2} dt \quad \text{AG}$ $2\sqrt{v} = -\frac{t}{2} + c$ $t = 0, v = 25 \Rightarrow c = 10$ $v = \left(\frac{20-t}{4}\right)^2$	M1		applying Newton's second law with $\frac{dv}{dt}$
		A1		correct differential equation
		dM1		separating variables
		dM1		integrating
		A1		correct integrals
dM1		finding the constant of integration		
A1		7	correct final result from correct working	
<b>(b)</b>	$t = 20$	B1	1	correct time
<b>Total</b>			<b>8</b>	

(Q6, June 2006)

<b>3 (a)</b>	Max speed $\equiv$ zero acceleration used	M1		Implied
	$\frac{72000}{60}$ $\frac{72000}{60} = k \times 60$ $k = 20$	M1 A1	3	
<b>(b)(i)</b>	$20v = -500 \frac{dv}{dt}$ $\frac{dv}{dt} = -\frac{v}{25}$	M1 A1	2	see $\frac{dv}{dt}$ , $\pm$
	<b>(ii)</b> $25 \int \frac{dv}{v} = - \int dt$ $[25 \ln v]_{20}^{10} = -[t]_0^t$ $25 \ln 10 - 25 \ln 20 = -t$ $t = 25 \ln 2 \text{ or } 17.3 \text{ or } -25 \ln \frac{1}{2}$	M1 A1 A1 m1 A1 A1	6	M1 separating variables  <b>Alternative</b> $25 \ln v = -t (+ c)$ A1  $t = 0, v = 20, c = 25 \ln 20$ m1 $t = t, v = 10,$ $25 \ln 10 = -t + 25 \ln 20$ A1 $t = 25 \ln 2 \text{ or } 17.3$ A1
<b>Total</b>			<b>11</b>	

(Q7, Jan 2007)

<b>4 (a)</b>	Using $F = ma$ :	M1	2	Condone no ‘-’ AG Note: no use of $m \Rightarrow$ no marks in (a)
	$-\lambda mv = ma = m \frac{dv}{dt}$ $\therefore \frac{dv}{dt} = -\lambda v$	A1		
<b>(b)</b>	$\int \frac{dv}{v} = -\lambda \int dt$ $\ln v = -\lambda t + c$ $v = C e^{-\lambda t}$	M1 A1	4	Needs ‘+ c’  Needs correct working AG
	When $t = 0, v = U \Rightarrow C = U$ $v = U e^{-\lambda t}$	M1 A1		
<b>Total</b>			<b>6</b>	

(Q7, June 2007)

<b>5 (a)</b>	Power of engine is 8kW ∴ Force exerted by engine = $\frac{8000}{v}$	M1A1		M1 for Power = $Fv$
	Using $F = ma$ : $\frac{8000}{v} - kv^2 = 600 \frac{dv}{dt}$ $600 \frac{dv}{dt} - \frac{8000}{v} + kv^2 = 0$	m1 A1	4	AG
<b>(b)(i)</b>	When engine is turned off, power is zero: $-kv^2 = 600 \frac{dv}{dt}$	B1	1	AG
<b>(ii)</b>	$\int 600 \frac{dv}{v^2} = -\int k dt$ $-\frac{600}{v} = -kt + c$ When $t = 0, v = 20$ : ∴ $c = -\frac{600}{20} = -30$ ∴ $\frac{600}{v} = kt + 30$ When $v = 10, kt = 30$ : ∴ $t = \frac{30}{k}$	M1 A1 A1 M1 A1	5	Need '+ c'    $-\frac{30}{k}$ SC3
	<b>Total</b>		<b>10</b>	

(Q8, Jan 2008)

<b>6 (a)</b>	Using $F = ma$ $-0.05mv = m \frac{dv}{dt}$ ∴ $\frac{dv}{dt} = -0.05v$	B1	1	Need to see $m$ terms
	<b>(b)</b> $\int \frac{dv}{v} = -\int 0.05 dt$ $\ln v = -0.05t + c$ $v = Ce^{-0.05t}$ When $t = 0, v = 20$ , ∴ $C = 20$ $v = 20e^{-0.05t}$	B1 M1 M1 A1	4	Need first 2 terms } fully correct solutions
<b>(c)</b>	When $v = 10, 10 = 20e^{-0.05t}$ $e^{0.05t} = 2$ ∴ $t = \frac{1}{0.05} \ln 2$ $= 13.9$	M1 A1 A1	3	Accept $20 \ln 2$
<b>Total</b>			<b>8</b>	

(Q6, June 2008)

7 (a)	Using $F = ma$ : $-0.08v^2 = 0.05a$ $\therefore \frac{dv}{dt} = -1.6v^2$	B1 B1	2	AG; condone sign error in first B1
	(b) $\int \frac{dv}{v^2} = -1.6 \int dt$ $-\frac{1}{v} = -1.6t (+c)$ When $t = 0, v = 3 \Rightarrow c = -\frac{1}{3}$ $\frac{1}{v} = \frac{1}{3} + 1.6t$ * $\frac{1}{v} = \frac{1}{3} + \frac{8}{5}t$ $\frac{1}{v} = \frac{5 + 24t}{15}$ $v = \frac{15}{5 + 24t}$	M1 A1 M1 A1  A1		
<b>Total</b>			<b>7</b>	

(Q8, Jan 2009)

8 (a)	Using $F = ma$ : $-\lambda mv^{\frac{3}{2}} = m \frac{dv}{dt}$ $\therefore \frac{dv}{dt} = -\lambda v^{\frac{3}{2}}$	M1 A1	2	AG
	(b) $\int \frac{dv}{v^{\frac{3}{2}}} = -\lambda \int dt$ $-\frac{2}{\frac{1}{v^{\frac{1}{2}}}} = -\lambda t + c$ When $t = 0, v = 9 \Rightarrow c = -\frac{2}{3}$ $\frac{2}{\sqrt{v}} = \lambda t + \frac{2}{3}$ $\frac{\sqrt{v}}{2} = \frac{1}{\lambda t + \frac{2}{3}}$ $v = \left( \frac{6}{2 + 3\lambda t} \right)^2$ $v = \frac{36}{(2 + 3\lambda t)^2}$	M1 A1 M1 A1 A1 m1 A1		
(c)	When $v = 4$ , $\frac{2}{\sqrt{v}} = \lambda t + \frac{2}{3} \Rightarrow 1 = \lambda t + \frac{2}{3}$ $t = \frac{1}{3\lambda}$	M1A1 A1	7 3	AG or $\frac{36}{(2 + 3\lambda t)^2} = 4$ M1 $(2 + 3\lambda t)^2 = 9$ A1 $t = \frac{1}{3\lambda}$ A1 needs statement why $2 + 3\lambda t \neq -3$
<b>Total</b>			<b>12</b>	

(Q8, June 2009)

9 (a)	Using $F = ma$ , $-0.2mv^{\frac{1}{2}} = m \frac{dv}{dt}$ $\therefore \frac{dv}{dt} = -0.2v^{\frac{1}{2}}$	B1	1	AG Must see equ'n containing $m$
(b)	$\int \frac{dv}{v^{\frac{1}{2}}} = -\int 0.2 dt$ $2v^{\frac{1}{2}} = -0.2t + c$ When $t=0, v=16 \therefore C=8$	M1		m1 for + c
	$2v^{\frac{1}{2}} = -0.2t + 8$ $v = (4 - 0.1t)^2$	A1m1		
	When $t=0, v=16 \therefore C=8$	A1		
	$2v^{\frac{1}{2}} = -0.2t + 8$ $v = (4 - 0.1t)^2$	A1	5	AG
(c)	When $v=1, 1 = (4 - 0.1t)^2$ $4 - 0.1t = \pm 1$ $t=30$ or $50$ $t=30$	M1		[if use $2v^{\frac{1}{2}} = 8 - 0.2t$ no need to see 50]
	$t=30$ or $50$	A1		
	$t=30$	A1	3	$t \neq 50$ as ball stops when $t = 40$
(d)	Integrating $v = (4 - 0.1t)^2$ : $v = 16 - 0.8t + 0.01t^2$ $x = 16t - 0.4t^2 + \frac{0.01}{3}t^3 + d$ When $t=0, x=0 \Rightarrow d=0$ $x = 16t - 0.4t^2 + \frac{0.01}{3}t^3$ When speed is $1 \text{ ms}^{-1}, t = 30$ $x = 480 - 360 + 90$ $= 210$	M1		M1 for first 3 terms or $-\frac{10}{3}(4 - 0.1t)^3$
	When $t=0, x=0 \Rightarrow d=0$	A1		
	$x = 16t - 0.4t^2 + \frac{0.01}{3}t^3$			
	When speed is $1 \text{ ms}^{-1}, t = 30$ $x = 480 - 360 + 90$ $= 210$	m1		dep on M1 above
		A1	4	[No 'd', 3 marks only]
<b>Total</b>			<b>13</b>	

(Q5, Jan 2010)

<b>10</b>	$\frac{dv}{dt} = -\frac{\lambda}{v^4}$	M1		
	$\int v^4 dv = -\int \lambda dt$	m1		Condone one of $v^{-4}$ , $+\int \lambda dt$ , $\frac{1}{\lambda}$
	$\frac{4}{5}v^{\frac{5}{4}} = -\lambda t + c$	A1A1 m1		m1 for + c
	$t = 0, v = u \therefore c = \frac{4}{5}u^{\frac{5}{4}}$	A1		
	$\therefore v^{\frac{5}{4}} = u^{\frac{5}{4}} - \frac{5}{4}\lambda t$			
	$v = \left(u^{\frac{5}{4}} - \frac{5}{4}\lambda t\right)^{\frac{4}{5}}$	A1	7	
<b>Total</b>			<b>7</b>	

(Q5, June 2010)

<b>11 (a)(i)</b>	$F = 65g - 260v$ $= 65(9.8 - 4v)$	B1	1	Accept $260v - 65g$ AG must see $65g$ or $260$
	<b>(ii)</b> Using $F = ma$			
	$65 \frac{dv}{dt} = 65(9.8 - 4v)$	M1		Need to see terms in $m$ (condone - sign)
	$\frac{dv}{dt} = -4(v - 2.45)$	A1	2	AG
<b>(b)</b>	$\frac{1}{v - 2.45} \frac{dv}{dt} = -4$	B1		
	$\int \frac{1}{v - 2.45} dv = -\int 4 dt$			
	$\ln(v - 2.45) = -4t + c$	M1 A1		M1 log side correct $-4t + c$
	$v - 2.45 = Ce^{-4t}$ $t = 0, v = 19.6$ $\therefore C = 17.15$ or $e^{2.84}$	A1		Or $c = \ln 17.15$ or $2.84$
	$\therefore v = 2.45 + 17.15e^{-4t}$ $2.45 + 17.2e^{-4t}$	A1	5	
<b>Total</b>			<b>8</b>	

(Q8, Jan 2011)

<p><b>12 (a)</b></p> <p>Using <math>F = ma</math></p> $-2mv^{\frac{5}{4}} = m \frac{dv}{dt}$ $\therefore \frac{dv}{dt} = -2v^{\frac{5}{4}} \quad \mathbf{AG}$ <p><b>(b)</b></p> $\int \frac{dv}{v^{\frac{5}{4}}} = -2 \int dt$ $-\frac{4}{\frac{1}{v^4}} = -2t + c$ <p>When <math>t = 0, v = 16 \Rightarrow c = -2</math></p> $-\frac{4}{\frac{1}{v^4}} = -2t - 2$ $v^{\frac{1}{4}} = \frac{2}{1+t}$ $v = \left( \frac{2}{1+t} \right)^4 \quad \mathbf{AG}$	<p>B1</p> <p>M1</p> <p>A1</p> <p>dM1</p> <p>A1</p> <p>A1</p>	<p>1</p> <p>5</p> <p>6</p>	<p>B1: Must see <math>-2mv^{\frac{5}{4}} = m \frac{dv}{dt}</math> or <math>-2mv^{\frac{5}{4}} = ma</math> and correct final answer.</p> <p>M1: Two integrals with one in the form <math>\int f(v)dv</math> where <math>f(v) = v^{\pm \frac{5}{4}}</math> or <math>v^{\pm \frac{4}{5}}</math>. The other integral must not contain <math>v</math> terms.</p> <p>A1: Correct expression. Condone lack of <math>+c</math> for this A1, but no subsequent marks if no <math>c</math>.</p> <p>dM1: Using <math>t = 0</math> and <math>v = 16</math> to find <math>c</math>. A1: Obtaining <math>c = -2</math>.</p> <p>A1: Correct final answer. Must see <math>v^{\frac{1}{4}} = \frac{2}{1+t}</math> or <math>v^{-\frac{1}{4}} = \frac{1+t}{2}</math> or <math>\frac{1}{v^{\frac{1}{4}}} = \frac{1+t}{2}</math></p> <p>Or</p> <p>if they obtain <math>v = \left( \frac{2}{t+c} \right)^4</math></p> <p><math>v = 16, t = 0 \Rightarrow 16^{\frac{1}{4}} = \frac{2}{c}</math>, condone <math>c = 1</math> (no other root considered)</p>
<b>Total</b>		<b>6</b>	

(Q6, June 2011)



13	(a) using $F = ma$ $0.4 \frac{dv}{dt} = 2 - 4v$ $\frac{dv}{dt} = -10(v - 0.5)$	M1 A1	2	Needs line above
	(b) hence $\int \frac{1}{v-0.5} dv = -\int 10 dt$ $\ln(v - 0.5) = -10t + c$ $v - 0.5 = Ce^{-10t}$ $t = 0, v = 1$ $\therefore C = 0.5$ $\therefore v = 0.5 + 0.5e^{-10t}$	M1A1 m1  A1 A1	5	M1 for any side integrated correctly m1 for + c (and M1 gained)  condone $v = 0.5 + e^{-10t-0.693}$
	(c) when $v = 0.55, 0.55 = 0.5 + 0.5e^{-10t}$ $10 = e^{10t}$ $t = \ln 10 \div 10$ $= 0.230$	M1 A1 A1	3	substitute 0.55 into C's (b), after finding c, possible numerical error
<b>Total</b>			<b>10</b>	

(Q6, Jan 2012)

14	(a) Using $F = ma$ : $m \frac{dv}{dt} = 49 - 9.8v$ or $5g - 9.8v$ $\therefore \frac{dv}{dt} = -1.96(v - 5)$	M1 A1	2	Need to see $m \frac{dv}{dt}$ or $5 \frac{dv}{dt}$ or $a = \frac{49 - 9.81}{5}$  Must see $m$ terms (not $a = \dots$ )
	(b) $\int \frac{dv}{v-5} = -1.96 \int dt$ $\ln(v - 5) = -1.96t + c$ When $t = 0, v = 7 \Rightarrow c = \ln 2$ $\ln \frac{v-5}{2} = -1.96t$ $\frac{v-5}{2} = e^{-1.96t}$ $v = 5 + 2e^{-1.96t}$	M1 A1A1 A1  A1	5	And one side integrated Need + c, A1 each side OE  CAO
	<b>Total</b>			<b>7</b>

(Q7, June 2012)

<b>15 (a)</b>	Using $F = ma$ : $-4v^{\frac{1}{3}} = 12 \frac{dv}{dt}$ $\therefore \frac{dv}{dt} = -\frac{1}{3}v^{\frac{1}{3}}$ $-3 \int \frac{dv}{v^{\frac{1}{3}}} = \int dt$ $-3 \times \frac{v^{\frac{2}{3}}}{\frac{2}{3}} = t + c$ $-\frac{9}{2}v^{\frac{2}{3}} = t + c$ When $t = 0, v = 8 \Rightarrow c = -18$ $-\frac{9}{2}v^{\frac{2}{3}} = t - 18$ $v^{\frac{2}{3}} = 4 - \frac{2}{9}t$ $v = \left(4 - \frac{2}{9}t\right)^{\frac{3}{2}}$	B1		
		M1		condone -, 3 incorrect side
		A1		condone lack of + c
		M1A1		
		A1	6	
<b>(b)</b>	Particle is at rest when $4 - \frac{2}{9}t = 0$ The value of $t$ is 18	B1	1	
<b>Total</b>			<b>7</b>	

(Q5, Jan 2013)

<b>16 (a)</b>	Using $F = ma$ $1600 \frac{dv}{dt} = 4000 - 40v$ $\frac{dv}{dt} = \frac{4000 - 40v}{1600}$ $\frac{dv}{dt} = \frac{100 - v}{40}$	M1		
		A1	2	
<b>(b)</b>	$40 \frac{dv}{100 - v} = dt$ $40 \int \frac{dv}{100 - v} = \int dt$ $-40 \ln(100 - v) = t + c$ When $t = 0, v = 0 \Rightarrow c = -40 \ln 100$ $-40 \ln(100 - v) = t - 40 \ln 100$ $t = 40 \ln \frac{100}{100 - v}$ $e^{\frac{t}{40}} = \frac{100}{100 - v}$ $v = 100 - 100e^{-\frac{t}{40}}$ or $100(1 - e^{-\frac{t}{40}})$	B1		
		M1		
		A1		Condone lack of '+ c'
		M1A1		
		A1	6	
<b>Total</b>			<b>8</b>	

(Q6, June 2013)