

Section A: Pure Mathematics

1 Let $x = 10^{100}$, $y = 10^x$, $z = 10^y$, and let

$$a_1 = x!, \quad a_2 = x^y, \quad a_3 = y^x, \quad a_4 = z^x, \quad a_5 = e^{xyz}, \quad a_6 = z^{1/y}, \quad a_7 = y^{z/x}.$$

- (i) Use Stirling's approximation $n! \approx \sqrt{2\pi} n^{n+\frac{1}{2}} e^{-n}$, which is valid for large n , to show that $\log_{10}(\log_{10} a_1) \approx 102$.
- (ii) Arrange the seven numbers a_1, \dots, a_7 in ascending order of magnitude, justifying your result.

2 Consider the quadratic equation

$$nx^2 + 2x\sqrt{(pn^2 + q)} + rn + s = 0, \quad (*)$$

where $p > 0$, $p \neq r$ and $n = 1, 2, 3, \dots$.

- (i) For the case where $p = 3$, $q = 50$, $r = 2$, $s = 15$, find the set of values of n for which equation (*) has no real roots.
- (ii) Prove that if $p < r$ and $4q(p - r) > s^2$, then (*) has no real roots for any value of n .
- (iii) If $n = 1$, $p - r = 1$ and $q = s^2/8$, show that (*) has real roots if, and only if, $s \leq 4 - 2\sqrt{2}$ or $s \geq 4 + 2\sqrt{2}$.

3 Let

$$S_n(x) = e^{x^3} \frac{d^n}{dx^n} (e^{-x^3}).$$

Show that $S_2(x) = 9x^4 - 6x$ and find $S_3(x)$.

Prove by induction on n that $S_n(x)$ is a polynomial. By means of your induction argument, determine the order of this polynomial and the coefficient of the highest power of x .

Show also that if $\frac{dS_n}{dx} = 0$ for some value a of x , then $S_n(a)S_{n+1}(a) \leq 0$.

- 4 By considering the expansions in powers of x of both sides of the identity

$$(1+x)^n(1+x)^n \equiv (1+x)^{2n},$$

show that

$$\sum_{s=0}^n \binom{n}{s}^2 = \binom{2n}{n},$$

where $\binom{n}{s} = \frac{n!}{s!(n-s)!}$.

By considering similar identities, or otherwise, show also that:

- (i) if n is an even integer, then

$$\sum_{s=0}^n (-1)^s \binom{n}{s}^2 = (-1)^{n/2} \binom{n}{n/2};$$

(ii)
$$\sum_{t=1}^n 2t \binom{n}{t}^2 = n \binom{2n}{n}.$$

- 5 Show that if α is a solution of the equation

$$5\cos x + 12\sin x = 7,$$

then either

$$\cos \alpha = \frac{35 - 12\sqrt{120}}{169}$$

or $\cos \alpha$ has one other value which you should find.

Prove carefully that if $\pi/2 < \alpha < \pi$, then $\alpha < 3\pi/4$.

- 6 Find $\frac{dy}{dx}$ if

$$y = \frac{ax + b}{cx + d}. \quad (*)$$

By using changes of variable of the form $(*)$, or otherwise, show that

$$\int_0^1 \frac{1}{(x+3)^2} \ln \left(\frac{x+1}{x+3} \right) dx = \frac{1}{6} \ln 3 - \frac{1}{4} \ln 2 - \frac{1}{12},$$

and evaluate the integrals

$$\int_0^1 \frac{1}{(x+3)^2} \ln \left(\frac{x^2 + 3x + 2}{(x+3)^2} \right) dx \quad \text{and} \quad \int_0^1 \frac{1}{(x+3)^2} \ln \left(\frac{x+1}{x+2} \right) dx.$$

- 7 The curve C has equation

$$y = \frac{x}{\sqrt{(x^2 - 2x + a)}},$$

where the square root is positive. Show that, if $a > 1$, then C has exactly one stationary point.

Sketch C when (i) $a = 2$ and (ii) $a = 1$.

- 8 Prove that

$$\sum_{k=0}^n \sin k\theta = \frac{\cos \frac{1}{2}\theta - \cos(n + \frac{1}{2})\theta}{2 \sin \frac{1}{2}\theta}. \quad (*)$$

- (i) Deduce that, when n is large,

$$\sum_{k=0}^n \sin \left(\frac{k\pi}{n} \right) \approx \frac{2n}{\pi}.$$

- (ii) By differentiating (*) with respect to θ , or otherwise, show that, when n is large,

$$\sum_{k=0}^n k \sin^2 \left(\frac{k\pi}{2n} \right) \approx \left(\frac{1}{4} + \frac{1}{\pi^2} \right) n^2.$$

[The approximations, valid for small θ , $\sin \theta \approx \theta$ and $\cos \theta \approx 1 - \frac{1}{2}\theta^2$ may be assumed.]

Section B: Mechanics

9 In the Z -universe, a star of mass M suddenly blows up, and the fragments, with various initial speeds, start to move away from the centre of mass G which may be regarded as a fixed point. In the subsequent motion the acceleration of each fragment is directed towards G . Moreover, in accordance with the laws of physics of the Z -universe, there are positive constants k_1 , k_2 and R such that when a fragment is at a distance x from G , the magnitude of its acceleration is k_1x^3 if $x < R$ and is k_2x^{-4} if $x \geq R$. The initial speed of a fragment is denoted by u .

- (i) For $x < R$, write down a differential equation for the speed v , and hence determine v in terms of u , k_1 and x for $x < R$.
- (ii) Show that if $u < a$, where $2a^2 = k_1R^4$, then the fragment does not reach a distance R from G .
- (iii) Show that if $u \geq b$, where $6b^2 = 3k_1R^4 + 4k_2/R^3$, then from the moment of the explosion the fragment is always moving away from G .
- (iv) If $a < u < b$, determine in terms of k_2 , b and u the maximum distance from G attained by the fragment.

10 N particles $P_1, P_2, P_3, \dots, P_N$ with masses $m, qm, q^2m, \dots, q^{N-1}m$, respectively, are at rest at distinct points along a straight line in gravity-free space. The particle P_1 is set in motion towards P_2 with velocity V and in every subsequent impact the coefficient of restitution is e , where $0 < e < 1$. Show that after the first impact the velocities of P_1 and P_2 are

$$\left(\frac{1 - eq}{1 + q}\right)V \quad \text{and} \quad \left(\frac{1 + e}{1 + q}\right)V,$$

respectively.

Show that if $q \leq e$, then there are exactly $N - 1$ impacts and that if $q = e$, then the total loss of kinetic energy after all impacts have occurred is equal to

$$\frac{1}{2}me(1 - e^{N-1})V^2.$$

- 11 An automated mobile dummy target for gunnery practice is moving anti-clockwise around the circumference of a large circle of radius R in a horizontal plane at a constant angular speed ω . A shell is fired from O , the centre of this circle, with initial speed V and angle of elevation α . Show that if $V^2 < gR$, then no matter what the value of α , or what vertical plane the shell is fired in, the shell cannot hit the target.

Assume now that $V^2 > gR$ and that the shell hits the target, and let β be the angle through which the target rotates between the time at which the shell is fired and the time of impact. Show that β satisfies the equation

$$g^2\beta^4 - 4\omega^2V^2\beta^2 + 4R^2\omega^4 = 0.$$

Deduce that there are exactly two possible values of β .

Let β_1 and β_2 be the possible values of β and let P_1 and P_2 be the corresponding points of impact. By considering the quantities $(\beta_1^2 + \beta_2^2)$ and $\beta_1^2\beta_2^2$, or otherwise, show that the linear distance between P_1 and P_2 is

$$2R \sin\left(\frac{\omega}{g}\sqrt{V^2 - Rg}\right).$$

Section C: Probability and Statistics

- 12** It is known that there are three manufacturers A, B, C, who can produce micro chip MB666. The probability that a randomly selected MB666 is produced by A is $2p$, and the corresponding probabilities for B and C are p and $1 - 3p$, respectively, where $0 \leq p \leq \frac{1}{3}$. It is also known that 70% of MB666 micro chips from A are sound and that the corresponding percentages for B and C are 80% and 90%, respectively.

Find in terms of p , the conditional probability, $P(A|S)$, that if a randomly selected MB666 chip is found to be sound then it came from A, and also the conditional probability, $P(C|S)$, that if it is sound then it came from C.

A quality inspector took a random sample of one MB666 micro chip and found it to be sound. She then traced its place of manufacture to be A, and so estimated p by calculating the value of p that corresponds to the greatest value of $P(A|S)$. A second quality inspector also took a random sample of one MB666 chip and found it to be sound. Later he traced its place of manufacture to be C and so estimated p by applying the procedure of his colleague to $P(C|S)$.

Determine the values of the two estimates and comment briefly on the results obtained.

- 13** A stick is broken at a point, chosen at random, along its length. Find the probability that the ratio, R , of the length of the shorter piece to the length of the longer piece is less than r .

Find the probability density function for R , and calculate the mean and variance of R .

- 14** You play the following game. You throw a six-sided fair die repeatedly. You may choose to stop after any throw, except that you must stop if you throw a 1. Your score is the number obtained on your last throw. Determine the strategy that you should adopt in order to maximize your expected score, explaining your reasoning carefully.