

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

	CANDIDATE NAME					
	CENTRE NUMBER				CANDIDATE NUMBER	
* 4 1 8	PHYSICS					0625/05
°	Paper 5 Practical Test				May/June 2008	
•						1 hour 15 minutes
5 F	Candidates answer on the Question Paper.					
(J)						

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
4		
Total		

This document consists of **11** printed pages and **1** blank page.



For

Examiner's Use

1 In this experiment, you are to determine the weight of a mass using a balancing method. Carry out the following instructions referring to Fig. 1.1.



Fig. 1.1

- (a) Place the mass on the metre rule so that its centre is directly above the 10.0 cm mark.
- (b) Describe, with the aid of a diagram, how you judged that the centre of the mass was directly above the 10.0 cm mark.

[1]

- (c) Place the rule on the pivot so that the rule is as near as possible to being balanced.
- (d) Measure and record the distance *a* from the 50.0 cm mark (centre) of the rule to the pivot and the distance *b* from the centre of the mass to the pivot.

a =

b =[3]

(e)	Calculate the weight W of the mass, using the equation	For
	$W = \frac{ka}{b}.$	Examiner's Use
	The value of k is given to you on a card close to the apparatus. Record the value of k.	
	<i>k</i> =	
	<i>W</i> =[1]	
(f)	Repeat the steps (c) to (e), with the centre of the mass directly above the 20.0 cm mark	
	<i>a</i> =	
	<i>b</i> =	
	<i>W</i> =[3]	
(g)	Calculate and record the average of the two values for W. Show your working.	
	Average value for $W = \dots$ [2]	
	[Total: 10]	

For

Examiner's Use

- 2 In this experiment you are to compare the combined resistance of lamps in different circuit arrangements.
 - (a) The circuit shown in Fig. 2.1 has been set up for you.





- (i) Switch on. Measure and record in row 1 of Table 2.1 (on page 5) the current *I* in the circuit and the p.d. *V* across points **A** and **B**. Switch off.
- (ii) Calculate the combined resistance *R* of the three lamps using the equation

$$R = \frac{V}{I}.$$

Record this value of R in the table.

(iii) Complete the column headings for each of the *V*, *I* and *R* columns of the table.

For

Use

(iv) Disconnect the lamps and then reconnect them between points A and B as shown in Fig. 2.2. Examiner's



- (v) Switch on. Measure and record in row 2 of Table 2.1 the current I in the circuit and the p.d. V across points A and B. Switch off.
- (vi) Calculate the combined resistance R of the three lamps using the equation

$$R = \frac{V}{I}.$$

Record this value of *R* in Table 2.1.

Table 2.1

	VI	11	RI
Fig. 2.1			
Fig. 2.2			
Fig. 2.3			
Fig. 2.4			

(vii) Repeat steps (iv) to (vi) using the arrangements of the lamps shown in Fig. 2.3 and Fig. 2.4. [7]

- (b) Theory suggests that, if all three lamps have the same resistance under all conditions, the combined resistance R recorded in row 1 will be one ninth of the combined resistance R recorded in row 4.
 - (i) State whether, within the limits of experimental accuracy, your results support this theory. Justify your answer by reference to the results.

statement





You are provided with two containers labelled **A** (a beaker) and **B** (a measuring cylinder). You also have a supply of hot water.

- (a) (i) Pour 100 cm^3 of hot water into container A.
 - (ii) Measure the temperature θ of the hot water. Record this temperature in Table 3.1 (on page 8) for time t = 0 s.

© UCLES 2008

3

For

Examiner's Use

(iii) Start the stopwatch and then record the temperature of the water at 30s intervals for a total of four minutes.

Table 3.1				
	container A (beaker)	container B (measuring cylinder)		
t / s	θ Ι °C	<i>θI</i> °C		
0				
30				
60				
90				
120				
150				
180				
210				
240				

[3]

- (iv) Pour 100 cm^3 of hot water into container **B**.
- (v) Repeat steps (ii) and (iii).

For

Examiner's Use

- (b) (i) Use the data in Table 3.1 to plot a graph of the temperature of the water in the beaker (*y*-axis) against time (*x*-axis). Draw the best fit curve.
 - (ii) Using the same graph axes, plot another graph, of the temperature of the water in the measuring cylinder against time.



(c) The experiment you have just done was designed to investigate the effect of the surface area exposed to the air on the rate of cooling of hot water. State the effect of a larger surface area on the rate of cooling. Justify your answer by reference to your graph.

For

Use

In this experiment you are to determine a quantity called the refractive index of the material 4 of a transparent block. Examiner's

Carry out the following instructions referring to Fig. 4.1.



Fig. 4.1

- (a) Place the transparent block, largest face down, on the ray-trace sheet supplied. The block should be approximately in the middle of the paper. Draw the outline of the block ABCD.
- (b) Remove the block and draw the normal NN' to side AB so that the normal is 2.0 cm from A. Label the point G where NN' crosses DC. Label the point F where NN' crosses AB.
- (c) Draw the line EF at an angle of 30° to the normal as shown in Fig. 4.1.

(d)	Place the paper on the pin board.	For		
(e)	Place two pins P_1 and P_2 on line EF as shown in Fig. 4.1.			
(f)	Replace the block and observe the images of P_1 and P_2 through side CD of the block so that the images of P_1 and P_2 appear one behind the other. Place two pins P_3 and P_4 between your eye and the block so that P_3 and P_4 and the images of P_1 and P_2 , seen through the block, appear one behind the other. Mark the positions of P_1 , P_2 , P_3 and P_4 . Remove the block.			
(g)	Draw a line joining the positions of P_3 and P_4 . Continue the line until it meets CD . Label this point H .			
(h)	Measure and record the length <i>a</i> of the line GH .			
	<i>a</i> =[1]			
(i)	Draw the line HF .			
(j)	Measure and record the length <i>b</i> of the line HF.			
	<i>b</i> =[1]			
(k)	Extend the straight line EF through the outline of the block to a point J . The point J must be at least 5 cm from the block. The line EJ crosses the line CD . Label this point K .			
(I)	(i) Measure and record the length <i>c</i> of the line GK .			
	<i>C</i> =			
	(ii) Measure and record the length <i>d</i> of the line FK .			
	<i>d</i> =[1]			
(m)	Calculate the refractive index <i>n</i> of the material of the block using the equation			
	$n = \frac{cb}{ad}$			
	<i>n</i> =[2]			
	Tie your ray-trace sheet opposite this page.[5]			
	[Total: 10]			

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.