
1. The diagram shows a section through part of a cell as it would appear when seen with an electron microscope.



| This cell produces and secretes a protein. Describe the part played by organelles A, B at C in producing and secreting this protein. | nd |
|--|----|
| | |
| | |
| | |
| | |
| | |
| | |

(a)

(3)

(b) The table shows information about the different parts of this cell.

| Part of cell | Percentage of total cell volume | Number in the cell |
|--|---------------------------------|--------------------|
| Cytoplasm surrounding cell organelles | 54 | 1 |
| Mitochondria | 22 | about 1700 |
| Nucleus | 6 | 1 |
| Lysosomes | 1 | about 300 |
| Cisternae of rough endoplasmic reticulum | 9 | 1 |

| (i) | Which organelle is larger, a mitochondrion or a lysosome? Use calculations based on figures from the table to support your answer. | |
|------|--|---------|
| | Larger organelle; | |
| | | |
| | | |
| | | |
| | | |
| | | (2) |
| (ii) | In the drawing there appear to be a number of separate cisternae in the rough endoplasmic reticulum. The table gives the approximate number of cisternae as one. Suggest an explanation for the apparent difference. | |
| | | |
| | | |

| (3) |
|-----|
| |

2. Read the following passage.

A red blood cell is packed full of haemoglobin. When mature, it contains none of the organelles usually found in an animal cell. The nucleus, endoplasmic reticulum. mitochondria and ribosomes are all absent.

- More is known about the plasma membrane of a human red blood cell than about any other eukaryotic cell membrane. One reason for this is that the plasma membrane surrounding a red blood cell can be isolated without being contaminated by internal cell membranes. Red blood cell plasma membranes or "ghosts" can be prepared by putting the cells in a dilute salt solution. This causes the cells to swell and burst, leaving only the plasma membrane.
- 10 Red blood cell ghosts have been investigated and found to contain several different proteins. One of these proteins is spectrin. It is made up of long polypeptide chains which form a network on the inside of the membrane. Spectrin strengthens the membrane and is involved in maintaining the three-dimensional shape of the red blood cell.

(2)

(1)

Use the information from the passage and your own knowledge to answer the following questions.

(a) Complete the table by giving **two** ways in which the structure of a red blood cell differs from the structure of a bacterial cell.


| Red blood cell | Bacterial cell |
|----------------------|------------------------------|
| Contains haemoglobin | Does not contain haemoglobin |
| Contains spectrin | Does not contain spectrin |
| | |
| | |
| | |
| | |

| (b) | | noglobin is a protein. Explain why a mature red blood cell cannot make oglobin. | |
|-----|-----|---|-----|
| | | | |
| | | | |
| | | | |
| | | | (2) |
| (c) | (i) | Plasma membranes that have been isolated from red blood cells are not contaminated by internal cell membranes (lines 6 - 7). Explain why. | |
| | | | |

| | (ii) | When red blood cells are put in a dilute salt solution they swell (line 8). Use your knowledge of water potential to explain why. | |
|-------------|------|---|-----|
| | | | |
| | | | |
| | | | (2) |
| <i>(</i> 1) | a | | |
| (d) | | e people have red blood cells that do not contain spectrin. These red blood cells are rical in shape. They also burst more quickly when put in distilled water. | |
| | (i) | Explain why more oxygen is taken up by normal red blood cells than by these spherical cells. | |
| | | | |
| | | | (1) |
| | (ii) | Explain why red blood cells that do not contain spectrin burst more quickly when put into distilled water. | |
| | | | |
| | | | (1) |

| (e) | Polypeptides such as spectrin are formed from amino acids. Describe the structure of an amino acid molecule and explain how amino acids link together. |
|-----|--|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | (6) (Total 15 marks) |
| | (====================================== |

3. The drawing shows an electron micrograph of a section through part of an alveolus from a lung.



| Cell A is a eukaryotic cell. Give two featu which are not found in cell A . | res that may be found in a prokaryotic cell |
|---|---|
| | |
| | |
| | |
| | |
| | |
| | n one advantage of a biconcave disc over a |
| pherical cell of the same volume in transp | oorting oxygen. |
| pherical cell of the same volume in transp | porting oxygen. |
| pherical cell of the same volume in transp | oorting oxygen. |
| pherical cell of the same volume in transp | oorting oxygen. |
| pherical cell of the same volume in transp | oorting oxygen. |
| The diameter of a human red blood cell is | 7 μm. |
| The diameter of a human red blood cell is | 7 μm. |
| The diameter of a human red blood cell is | 7 μm. |

| | (ii) | In calculating the magnification, what assumption did you have to make about how the section was cut? | |
|-------|---------------------------|---|--------------|
| | | | |
| | | (Total 8 ma | (1) arks) |
| Read | the fo | ollowing passage. | |
| 5 | Hum prop Thes | an milk contains all the nutrients a young baby needs in exactly the right ortions. It is formed in the mammary glands by small groups of milk-producing cells. e cells absorb substances from the blood and use them to synthesise the lipids, bhydrates and proteins found in milk. Milk-producing cells are roughly cube-shaped have a height to breadth ratio of approximately 1.2:1. | |
| | cond | main carbohydrate in milk is lactose. Lactose is a disaccharide formed by the ensation of two monosaccharides, glucose and galactose. (A molecule of galactose he same formula as a molecule of glucose – the atoms are just arranged in a different) | |
| 10 | cyto _l diam | ose is synthesised in the Golgi apparatus and transported in vesicles through the blasm. Because lactose is unable to escape from these vesicles, they increase in eter as they move towards the plasma membrane. The vesicle membranes fuse with lasma membrane and the vesicles empty their contents out of the cell. | |
| Use t | | ormation from the passage and your own knowledge to answer the following | |
| (a) | (i) | The breadth of a milk-producing cell is 26 μm . Calculate the height of this cell. | |
| | | Height = μm | (1) |
| | | | |

4.

| | (ii) | Describe and explain how you would expect the height to breadth ratio of an epithelial cell from a lung alveolus to differ from the height to breadth ratio of a milk-producing cell. |
|-----|------|---|
| | | |
| | | |
| | | |
| | | |
| | | |
| (b) | How | many oxygen atoms are there in a molecule of |
| | (i) | galactose; |
| | | |
| | (ii) | lactose? |
| | | |
| | | |
| (c) | mem | lactose-containing vesicles increase in diameter as they move towards the plasma abrane of the milk-producing cell (lines 11-12). Use your knowledge of water ntial to explain why. |
| | | |
| | | |
| | | |
| | | |

| Suggest one advantage of milk-producing cells containing large numbers of mitochondria. | |
|--|----------|
| | |
| | |
| | |
| | |
| Some substances pass through the plasma membrane of a milk-producing cell by diffusion. Describe the structure of a plasma membrane and explain how different substances are able to pass through the membrane by diffusion. | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| (Tots | al 15 ma |

| a) | Explain l a short ti | • | ell allows it to take up a large amount of o | xygen i |
|----|-------------------------|----------------------------------|---|---------|
| | | | | |
| | | | | •••• |
| | | | | |
| | | | | |
| | | | | |
| | mixture | | al volumes of different liquids. A drop of d with an optical microscope. The table sl | |
| | Slide | Liquid added | Appearance of slide | |
| | A | Distilled water | No cells seen. Slide appears a uniform pale red colour | |
| | В | Sucrose solution | Cells are smaller in diameter than in an untreated sample of blood | |
| | C | Detergent (dissolves lipids) | No cells seen. Slide appears a uniform pale red colour | |
| o) | | rrounding a red blood cell? | e B tell you about the plasma membrane | |
| | | | | |
| | (ii) Ex | xplain the appearance of slide C | 2. | |

| (c) | | blood from which these samples were taken also contained monocytes and blocytes. How could you use the appearance of a nucleus to | |
|-----|------|---|-----------------|
| | (i) | distinguish between a monocyte and a red blood cell; | |
| | | | |
| | | | (1) |
| | (ii) | identify a granulocyte? | |
| | | | |
| | | | (1) |
| | | | (Total 7 marks) |

6. Read the following passage.

In a human, there are over 200 different types of cell clearly distinguishable from each other. What is more, many of these types include a number of different varieties. White blood cells, for example, include lymphocytes and granulocytes.

- Although different animal cells have many features in common, each type has adaptations.


 associated with its function in the organism. As an example, most cells contain the same organelles, but the number may differ from one type of cell to another. Muscle cells contain many mitochondria, while enzyme-secreting cells from salivary glands have particularly large amounts of rough endoplasmic reticulum.
- The number of a particular kind of organelle may change during the life of the cell. An example of this change is provided by cells in the tail of a tadpole. As a tadpole matures into a frog, its tail is gradually absorbed until it disappears completely. Absorption is associated with an increase in the number of lysosomes in the cells of the tail.

| Use | inform | ation from the passage and your own knowledge to answer the following questions. | | | |
|-----|--------|--|-----|--|--|
| (a) | Expl | Explain the link between. | | | |
| | (i) | mitochondria and muscle cells (lines 6 - 7); | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | (3) | | |
| | (ii) | rough endoplasmic reticulum and enzyme-secreting cells from salivary glands (lines 7 - 8). | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | (2) | | |
| (b) | | information in the passage to explain how a tadpole's tail is absorbed as a tadpole ges into a frog. | | | |
| | | | | | |
| | ••••• | | | | |

| (c) | Starting with some lettuce leaves, describe how you would obtain a sample of undamaged chloroplasts. Use your knowledge of cell fractionation and ultracentrifugation to answer this question. |
|-----|--|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | (6) (Total 13 marks) |


7. Moulds belong to a group of organisms called fungi. When mould is examined with a microscope it is seen to consist of long, colourless threads called hyphae.

A student investigated the growth of fungal hyphae. The diagram shows part of a hypha seen under a microscope at the beginning of a lesson and again at the end of the lesson.



| | one piece of evidence from the diagram that fungi are eukaryotic. | |
|--------------|---|--|
| (i) | By how much had the hypha grown during the lesson? Show your working. | |
| | Answer: μm | |
| (ii) | Explain how you could use your answer to calculate the rate of growth of this hypha. | |
| | | |
| Unde show | er the microscope, small granules were seen in the hypha. Describe how you could that these granules consisted of starch. | |
| | | |
| | | |
| | | |

8. Liver was ground to produce a homogenate. The diagram shows how fractions containing different cell organelles were produced from the filtered homogenate.



| (a) | Explain why the homogenate was filtered before spinning at low speed in the centrifuge. |
|-----|---|
| | |
| | |
| | |
| | |
| | |

| chondria. Suggest the main | (b) The org | |
|-------------------------------------|-------------|----|
| | (i) | |
| | (ii) | |
| parated in this way? | (c) Wh | |
| | | |
| en with a transmission electron | (d) Exp | |
| | | |
| (Total 7 mar | | |
| Describe how the length of the cell | (a) A p | 9. |
| | | |
| | | |
| | | |

| (b) | The water potential of a plant cell is –400 kPa. The cell is put in a solution with a water potential of –650 kPa. Describe and explain what will happen to the cell. | | | | |
|-----|---|-----|--|--|--|
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | (3) | | | |
| | | | | | |
| (c) | A group of students investigated the effect of sucrose concentration on the change in | | | | |

(c) A group of students investigated the effect of sucrose concentration on the change in length of cylinders of tissue cut from a young carrot. They measured the initial lengths of the carrot cylinders, then placed one in each of a number of sucrose solutions. After 18 hours, they removed the carrot cylinders and measured their final lengths. Some of the results are shown in the table.

| Concentration of sucrose / mol dm ⁻³ | Percentage decrease in length of carrot cylinder |
|---|--|
| 0.4 | 4.2 |
| 0.5 | 8.7 |
| 0.6 | 13.0 |
| 0.7 | 16.8 |
| 0.8 | 18.1 |
| 0.9 | 18.1 |
| 1.0 | 18.1 |

| | 1.0 | 18.1 | | | |
|-----|---|------|-----|--|--|
| (i) | The carrot cylinders were left for 18 hours in the sucrose solutions. Explain why they were left for a long time. | | | | |
| | | | (1) | | |

| (ii) | Explain how you would use a graph to predict the concentration of sucrose that would result in no change in length of the carrot cylinders. | |
|-------------|---|----|
| | | |
| | | |
| | | |
| | | 2) |
| | | |
| (iii) | Young carrots store sugars in their tissues but, in older carrots, some of this is converted to starch. How would using cylinders of tissue from older carrots affect | |
| | the results obtained for a sucrose solution of 0.6 mol dm ⁻³ ? Give a reason for your answer. | |
| | | |
| | | |
| | | |
| | (Total 10 marks | |
| The diagram | n shows a mitochondrion. | |
| | | |
| | X X Y | |

10.

(a)

(i)


(ii)

Name the parts labelled \mathbf{X} and \mathbf{Y} .

X

Y

Scientists isolated mitochondria from liver cells. They broke the cells open in an ice-cold, isotonic solution. They then used a centrifuge to separate the cell organelles. The diagram shows some of the steps in the process of centrifugation.



Suggest which pellet, A, B or C contained the mitochondria.

(b)

| | | | (1) |
|-----|------|-------------------------------|-----|
| (c) | Expl | ain why the solution used was | |
| | (i) | ice-cold | |
| | | | |
| | | | (1) |
| | (ii) | isotonic. | |

| (d) | People with mitochondrial disease have mitochondria that do not function properly. |
|-----|--|
| | Some people with mitochondrial disease can only exercise for a short time. Explain why a |
| | person with mitochondrial disease can only exercise for a short time. |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | (2) |
| | (Total 8 marks) |
| | (Tutai o marks) |