1	Catalase is an enzyme that breaks down hydrogen peroxide inside cells. Red blood cells contain
	catalase.

Some dogs have an inherited condition in which catalase is not produced. This condition is known as acatalasia and it is caused by a mutation in the gene for catalase.

(a) Define the terms g	gene and gene	mutation.
--	---------------	-----------

gene	 	
gene mutation		

[2]

(b) A geneticist was asked to investigate the inheritance of acatalasia in dogs.

The normal allele is represented by **B** and the mutant allele is represented by **b**.

The geneticist made the diagram in Fig. 3.1 to show the inheritance of acatalasia in a family of dogs. The shaded symbols indicate the dogs with acatalasia.




Fig. 3.1

	_									
(i) 9	Stata tha	genotypes	of the	annh	identified	as 1	2 and	l 3 in	Fig '	ว 1

l	
2	
3	[3]

(ii)		log 4 with dog 5 . Approxim	nately half of the offspring had alasia.	
	Complete the genetic dia	agram to show how this is	possible.	
		dog 4	dog 5	
	parental phenotypes	normal	has acatalasia	
	parental genotypes			
	gametes	, (+	
	Punnett square			
	offspring phenotypes			[3]
(iii)		the type of cross that you		[4]
				[1]
			[Tota	ıl: 9]

2			in is a red pigment found in carnation flowers. Some carnation plants have a gene thocyanin.	for
	(a)	(i)	A flower grower bred red carnations.	
			Describe how growers selectively breed plants.	
				[3]
				ĮO,
		(ii)	Explain the disadvantages of using sexual reproduction to breed red carnations.	
				[2]
				L - .
	(b)	Meios	is is necessary for sexual reproduction of carnation plants.	
		Define	e the term <i>meiosis</i> .	
				lo.
				[2]

Carna	ition plants show co-dor	ninance for the	e anthocyan	in gene. Th	ere are two alle	eles:
• F	^A – allele for anthocyani	n pigment (red	d flowers)			
• F	^N – allele for no anthocy	anin pigment	(white flowe	rs)		
(i)	State the genotype of	a carnation pla	ant that is he	eterozygous	for this gene.	
						[1]
(ii)	Describe the phenotyp	e of a heteroz	ygous carna	ation plant f	or this gene.	
						[1]
(iii)	The breeder crossed a genetic diagram, the					
					•	
	parental genotypes	F ^A	F"	×	F ^A F ^A	
	gametes			+)
	Punnett square					
	offspring genotypes .					
	offspring phenotypes					
	proportion of pure bree	eding carnatio	n plants			
						[4]

[Total: 13]

3 (a) Sex in cats is determined in the same way as in humans.

Complete the diagram below to show how sex is determined in cats.

male cat × female cat

female cat (XX)

	gametes	X	X
male cat			
()			

offspring ratio

(b) A scientist investigated the inheritance of fur colour in cats.

The gene for coat colour is located on the X chromosome. The gene has two alleles:


- B black
- **b** orange.

The X chromosome with the allele for black is represented by $\mathbf{X}^{\mathbf{B}}$.

The X chromosome with the allele for orange is $\dot{\mathbf{X}}^{\mathbf{b}}$.

A female cat can be a mixture of these colours, described as calico.

Fig. 3.1 shows the inheritance of this condition in a family of cats.



(i)	State the genotypes of cats 1, 4, and 5 in Fig. 3.1.	
	cat 1	
	cat 4	
	cat 5	[3]
(ii)	Coat colour in cats is an example of discontinuous variation.	
	Explain why coat colour is an example of discontinuous variation.	
		[3]

[Total: 9]

4 Fig. 5.1 shows the nematode, *Caenorhabditis elegans*.



Fig. 5.1

(a)	(i)	State the genus of this nematode.
		[1]
	(ii)	State two structural features of nematodes.
		1
		2[2]
(b)		natodes feed on dead and decaying material. Explain why this gives nematodes an ortant role in ecosystems.
		[3]

(c) Fig. 5.2 shows the life cycle of *C. elegans*. The diploid number of this species is 12.




Fig. 5.2

(i)	Suggest why there is very little genetic variation in the offspring of the adult nematode shown in Fig. 5.2.
	[2]
(ii)	State the haploid number of <i>C. elegans</i> .
	[1]