

Version 1.2



**General Certificate of Education
June 2010**

Chemistry

CHEM5

Energetics, Redox and Inorganic Chemistry

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2010 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance

Q	Part	Sub Part	Marking Guidance	Mark	Comments
1	(a)		$\text{CaF}_2(\text{s}) \rightarrow \text{Ca}^{2+}(\text{g}) + 2\text{F}^{-}(\text{g})$	1	
1	(b)	(i)	Enthalpy change for formation of 1 mol of substance From its elements Reactants and products/all substances in their standard states	1 1 1	Allow <u>heat energy change</u> , NOT energy Or normal states at 298 K, 1 bar (100 kPa)
1	(b)	(ii)	$\text{Ca}(\text{s}) + \text{F}_2(\text{g}) \rightarrow \text{CaF}_2(\text{s})$	1	
1	(b)	(iii)	$\Delta H_f(\text{CaF}_2) = \Delta H_a(\text{Ca}) + 1\text{st IE}(\text{Ca}) + 2^{\text{nd}} \text{IE}(\text{Ca}) + \text{BE}(\text{F}_2) + 2\text{xEA}(\text{F}) - \Delta H_L(\text{CaF}_2)$ $= 193 + 590 + 1150 + 158 + (2 \times -348) - 2602$ $= -1207 \text{ kJ mol}^{-1}$	1 1 1	Or labelled diagram Correct answer scores 3 -842 scores 2 (transfer error) -859 scores 1 only (using one E.A.) Units not required, wrong units lose 1 mark
1	(c)		Electrostatic attraction stronger/ionic bonding stronger/attraction between ions stronger/more energy to separate ions Because fluoride (ion) smaller than chloride	1 1	Molecular attraction /atoms/intermolecular forces CE=0 Do not allow F or fluorine
1	(d)	(i)	$\Delta H = \Delta H_L + \Sigma \Delta H_{\text{hyd}} = 2237 - 1650 + (2 \times -364)$ $= -141 \text{ kJ mol}^{-1}$	1 1	Can be on cycle/diagram Correct answer scores 2 Units not required, wrong units lose 1 mark

1	(d)	(ii)	Decreases Reaction exothermic/ ΔH -ve (Equilibrium)shifts to left/backwards (as temperature rises)/ equilibrium opposes the change	1 1 1	If ans to (d)(i) positive allow increases If (d)(i) +ve allow endothermic/ ΔH +ve If (d) (i) +ve allow shifts to right/forwards / equilibrium opposes the change If no answer to (d) (i) assume $-ve \Delta H$ used If effect deduced incorrectly from any ΔH CE=0 for these 3 marks
1	(e)		u.v. absorbed: electrons/they move to higher energy (levels)/ electrons excited visible light given out: electrons/they fall back down/move to lower energy (levels)	1 1	Must refer to absorbing u.v. NOT visible light or this must be implied.

Q	Part	Sub Part	Marking Guidance	Mark	Comments
2	(a)		<p><u>Macromolecular</u></p> <p>Covalent bonding (between atoms)</p> <p>Many/strong bonds to be broken (or lots of energy required)</p>	<p>1</p> <p>1</p> <p>1</p>	<p>Or <u>giant</u> molecule</p> <p>Or <u>giant</u> covalent (also gains M2)</p> <p>Do not allow giant atomic</p> <p>Ionic/metallic CE=0 for all 3 marks</p> <p>Do NOT allow if between molecules</p> <p>Lose both bonding marks if contradiction e.g. mention of intermolecular forces</p> <p>Note: 'covalent bonds between molecules' loses M2 but not M3</p>
2	(b)		Al ₂ O ₃ <u>ionic</u>	1	Allow <u>ionic</u> + covalent/ <u>ionic</u> with covalent character
2	(c)		2Al + 3/2O ₂ → Al ₂ O ₃	1	Allow multiples Ignore state symbols
2	(d)		Insoluble/impermeable/non-porous	1	Or does not react/inert Do not allow thick layer Must imply property of Al ₂ O ₃ not Al
2	(e)		Na ₂ O + H ₂ O → 2NaOH	1	Or Na ₂ O + H ₂ O → 2Na ⁺ + 2OH ⁻
2	(f)	(i)	Al ₂ O ₃ + 6HCl → 2AlCl ₃ + 3H ₂ O	1	Ionic equations with Al ₂ O ₃ possible e.g. Al ₂ O ₃ + 6H ⁺ → 2Al ³⁺ + 3H ₂ O Do not allow formation of Al ₂ Cl ₆

2	(f)	(ii)	$\text{Al}_2\text{O}_3 + 2\text{NaOH} + 3\text{H}_2\text{O} \rightarrow 2\text{NaAl}(\text{OH})_4$	1	<p>Other equations with Al_2O_3 are possible e.g.</p> $\text{Al}_2\text{O}_3 + 2\text{OH}^- + 3\text{H}_2\text{O} \rightarrow 2[\text{Al}(\text{OH})_4]^-$ $\text{Al}_2\text{O}_3 + 2\text{OH}^- + 7\text{H}_2\text{O} \rightarrow 2[\text{Al}(\text{H}_2\text{O})_2(\text{OH})_4]^-$
2	(g)		<p>SiO_2 acidic/Lewis acid/electron pair acceptor</p> $\text{SiO}_2 + 2\text{NaOH} \rightarrow \text{Na}_2\text{SiO}_3 + \text{H}_2\text{O}$	1 1	<p>Allow SiO_2 not amphoteric Do NOT allow BL acid</p> <p>Other equations with SiO_2 are possible e.g.</p> $\text{SiO}_2 + 2\text{OH}^- \rightarrow \text{SiO}_3^{2-} + \text{H}_2\text{O}$ $\text{SiO}_2 + 2\text{OH}^- + 2\text{H}_2\text{O} \rightarrow \text{Si}(\text{OH})_6^{2-}$

Q	Part	Sub Part	Marking Guidance	Mark	Comments
3	(a)		Same phase/state	1	
3	(b)		Because only exist in one oxidation state	1	Allow do not have variable oxidation states
3	(c)		$2\text{I}^- + \text{S}_2\text{O}_8^{2-} \rightarrow \text{I}_2 + 2\text{SO}_4^{2-}$	1	Ignore state symbols Allow multiples
3	(d)		Both (ions) have a negative charge	1	Or both have the same charge Or (ions) repel each other Do not allow both molecules have the same charge (contradiction)
3	(e)		$2\text{Fe}^{2+} + \text{S}_2\text{O}_8^{2-} \rightarrow 2\text{Fe}^{3+} + 2\text{SO}_4^{2-}$ $2\text{Fe}^{3+} + 2\text{I}^- \rightarrow 2\text{Fe}^{2+} + \text{I}_2$ Positive and negative (ions)/oppositely charged (ions)	1 1 1	Equations can be in any order Mark independently
3	(f)		Equations 1 and 2 can occur in any order	1	Allow idea of Fe^{3+} converted to Fe^{2+} then Fe^{2+} converted back to Fe^{3+}

Q	Part	Sub Part	Marking Guidance	Mark	Comments
4	(a)		Partially filled/incomplete d sub-shell/orbital/shell	1	Ignore reference to f orbitals Do not allow d block Do not allow half-filled d orbitals
4	(b)		Has ligand(s) linked by co-ordinate bonds	1 1	Allow molecules/ions with lone pairs Allow dative/donation of lone pair
4	(c)		(Blue) light is absorbed (from incident white light) Due to electrons moving to higher levels / electrons excited Red light (that) remains (is transmitted) / light that remains (transmitted light) is the colour observed	1 1 1	 Allow d → d transitions Allow red light reflected
4	(d)	(i)	Circle round any O ⁻ Circle round either N	1 1	List principle
4	(d)	(ii)	$\text{EDTA}^{4-} + [\text{Co}(\text{H}_2\text{O})_6]^{2+} \rightarrow [\text{CoEDTA}]^{2-} + 6\text{H}_2\text{O}$	1	Allow missing square brackets Ignore state symbols
4	(d)	(iii)	Increase in entropy/ ΔS positive Because 2 mol (of particles/molecules/species/entities) form 7 mol	1 1	Or increase in disorder Allow 'increase in number' as stated in words or as shown by any numbers deduced correctly from an incorrect equation Do not allow increase in ions/atoms

4	(e)	(i)	Co-ordinate/dative/dative covalent bond	1	Allow pair of electrons donated by nitrogen/ligand Do not allow pair of electrons donated from Iron/Fe
			Covalent bond	1	Shared electron pair
4	(e)	(ii)	Transport of oxygen/O ₂	1	Allow any statement that implies oxygen carried (around the body) Do not allow transport of carbon dioxide (CO ₂). This also contradicts the mark (list principle)
4	(e)	(iii)	Because it bonds to the iron/haemoglobin	1	Allow blocks site /CO has greater affinity for haemoglobin /carboxyhaemoglobin more stable than oxyhaemoglobin
			Displaces <u>oxygen</u>	1	Or prevents transport of <u>oxygen</u> QoL

Q	Part	Sub Part	Marking Guidance	Mark	Comments
5	(a)		W is CuCl_4^{2-}	1	
			Yellow-green/yellow/green	1	Not necessary to indicate solution Do not allow precipitate/solid
			$[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 4\text{Cl}^- \rightarrow \text{CuCl}_4^{2-} + 6\text{H}_2\text{O}$	1	Allow $+ 4\text{HCl} \rightarrow 4\text{H}^+$
5	(b)		X is $\text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2$	1	Allow $\text{Cu}(\text{OH})_2$ /copper hydroxide
			Blue precipitate/solid	1	Ignore shades
			$[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{NH}_3 \rightarrow \text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2 + 2\text{NH}_4^+$	1	Allow any balanced equation/equations leading to this hydroxide or $\text{Cu}(\text{OH})_2$ But must use ammonia
5	(c)		Y is $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$	1	
			Deep/dark/royal <u>blue</u> solution	1	QoL
			$\text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2 + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 2\text{H}_2\text{O} + 2\text{OH}^-$	1	Accept equation for formation from $\text{Cu}(\text{OH})_2$
5	(d)		Z is CuCO_3	1	Allow copper carbonate
			Green solid/precipitate	1	Allow blue-green precipitate
			$[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + \text{CO}_3^{2-} \rightarrow \text{CuCO}_3 + 6\text{H}_2\text{O}$	1	
5	(e)	(i)	$\text{Cu}^{2+}(\text{aq}) + \text{Fe}(\text{s}) \rightarrow \text{Cu}(\text{s}) + \text{Fe}^{2+}(\text{aq})$	1	Allow hydrated ions State symbols not essential but penalise if wrong
		Blue	1	Do not allow description of solids	
		Green	1	Allow yellow/(red-)brown/orange	

5	(e)	(ii)	<p>Any two correct points about copper extraction from two of these three categories:</p> <p>Any relevant mention of lower energy consumption</p> <p>Any relevant mention of benefits of less mining (of copper ore)</p> <p>Less release of CO₂ (or CO) into the atmosphere</p>	Max 2	<p>Do not allow reference to electricity alone or to temperature alone.</p> <p>Allow avoids depletion of (copper ore) resources</p> <p>Not just greenhouse gases. Must mention CO₂ or CO</p>
---	-----	------	--	-------	---

Q	Part	Sub Part	Marking Guidance	Mark	Comments
6	(a)		$\Delta H = \sum \Delta H_f(\text{products}) - \sum \Delta H_f(\text{reactants})$ $= -201 - 242 - (-394)$ $= -49 \text{ kJ mol}^{-1}$	1 1 1	+49 kJ mol ⁻¹ = 1 mark units not required, wrong units lose 1 mark
6	(b)		$\Delta S = \sum S(\text{products}) - \sum S(\text{reactants})$ $= 238 + 189 - (214 + 3 \times 131)$ $= -180 \text{ J K}^{-1} \text{ mol}^{-1}$	1 1 1	+180 = 1 mark units not required, wrong units lose 1 mark
6	(c)		$\Delta G = \Delta H - T\Delta S$ <p>(ΔS is negative so) at high temp $-T\Delta S$ (is positive and) greater than ΔH / large</p> <p>So $\Delta G > 0$</p> <p>(Limiting condition $\Delta G = 0$ so) $T = \Delta H / \Delta S$</p> $= 272 \text{ K}$ <p>Reaction is too slow at this temperature/to speed up the reaction</p>	1 1 1 1 1 1	<p>If use G not ΔG penalise M1 but not M2 and M3</p> <p>Do not award M2 or M3 if positive ΔS value used</p> <p>Independent mark unless positive ΔS value used</p> <p>Allow 297-298 if used given values. Do not award M5 if T -ve or if M4 should give T -ve</p>

6	(d)	<p>$\text{CH}_3\text{OH} + 3/2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$</p> <p>2.5 mol give 3 mol (gases)</p> <p>Therefore ΔS is positive/entropy increases</p> <p>(combustion exothermic so ΔH -ve so $\Delta H - T\Delta S$) and hence ΔG always negative (less than zero)</p>	1	<p>Allow multiples. Ignore state symbols. Do not allow equation for wrong compound but mark on provided number of moles increases or stays the same. If no equation or equation that gives a decrease in the number of moles, CE = 0</p>
			1	<p>Allow statement 'increase in number of moles/molecules' If numerical values given, they must match the equation in M1 Ignore the effect of incorrect state symbols on the number of moles of particles unless used correctly</p>
			1	<p>If correct deduction from wrong equation is $\Delta S = 0$ or ΔS very small must say ΔH -ve</p>
			1	<p>Allow G instead of ΔG Can score 3 out of 4 marks if equation wrong but leads to increase or no change in number of moles M4 dependent on M3</p> <p>Note, if equation wrong AND there is an incorrect deduction about the change in number of moles, CE = 0</p>
6	(e)	<p>$\text{CO}_2/\text{CO}/\text{CH}_4$ may be produced during H_2 manufacture/building the plant/transport/operating the plant</p>	1	

Q	Part	Sub Part	Marking Guidance	Mark	Comments
7	(a)		Hydrogen /H ₂ <u>gas/bubbles</u>	1	Allow 1 bar instead of 100 kPa Do not allow 1 atm
			1.0 mol dm ⁻³ HCl / H ⁺	1	
			At 298K and 100kPa	1	
			Pt (electrode)	1	
7	(b)		Li ⁺ + MnO ₂ + e ⁻ → LiMnO ₂	1	Ignore state symbols
			-0.13(V)	1	
7	(c)		Fe ³⁺ ions reduced to Fe ²⁺	1	Can score from equation/scheme
			Because $E(\text{Fe}^{3+}/\text{Fe}^{2+}) > E(\text{H}^+/\text{H}_2) / E(\text{hydrogen})$	1	Allow emf/ E_{cell} +ve/0.77V Allow Fe ³⁺ better oxidising agent than H ⁺ Allow H ₂ better reducing agent than Fe ²⁺ Only award this explanation mark if previous mark given

7	(d)	<p>Moles $\text{Cr}_2\text{O}_7^{2-} = \underline{23.7 \times 0.01/1000} = 2.37 \times 10^{-4}$</p> <p>1 mol $\text{Cr}_2\text{O}_7^{2-}$ reacts with 6 mol Fe^{2+} so moles Fe^{2+} in 25 $\text{cm}^3 = 6 \times 2.37 \times 10^{-4} = 1.422 \times 10^{-3}$</p> <p>Moles Fe^{2+} in 250 $\text{cm}^3 = 1.422 \times 10^{-2}$</p> <p>Original moles $\text{Fe}^{2+} = \underline{10.00/277.9} = 0.0360$</p> <p>Moles Fe^{2+} oxidised = $0.0360 - 0.0142 = 0.0218$</p> <p>% oxidised = $(0.0218 \times 100)/0.0360 = 60.5\%$</p>	1 1 1 1 1 1	<p>M1 x 6</p> <p>M2 x 10 or M4/10</p> <p>Independent mark</p> <p>M4 – M3</p> <p>(M5 x 100)/M4 Allow 60 to 61 Note Max 3 if mol ratio for M2 wrong eg 1:5 gives 67.1% 1:1 gives 93.4%</p> <p>Note also, 39.5% (39-40) scores M1, M2, M3 and M4 (4 marks)</p>
---	-----	---	----------------------------	---