M1. (a) <u>Functional group</u> (isomerism)

(b)

M1 Tollens' (reagent)	M1 Fehling's (solution) or
(Credit ammoniacal silver nitrate OR	Benedict's solution
a description of making Tollens')	(Ignore Cu² (aq) or
(Ignore either $AgNO_3$ or $[Ag(NH_3)_2^+]$	CuSO₄ on their own, but mark on
or "the silver mirror test" on their	to M2 and M3)
own, but mark M2 and M3)	

M2 silver mirror

M2 <u>Red solid/precipitate</u> (Credit orange or brown <u>solid</u>)

OR

<u>black solid/precipitate</u> (NOT silver precipitate)

M3 (stays) colourless	M3 (stays) blue
or no change or no reaction	or no change or no reaction

Mark on from an incomplete/incorrect attempt at the correct reagent, penalising M1

No reagent, CE=0 Allow the following alternatives **M1** (acidified) potassium dichromate(VI) (solution) **M2** (turns) green **M3** (stays) orange/no change OR **M1** (acidified) potassium manganate(VII) (solution) **M2** (turns) colourless **M3** (stays) purple/no change For M3 Ignore "nothing (happens)" Ignore "no observation"

3

1

- (c) (Both have) C=O OR a carbonyl (group)
 (d) (i) (Free-) radical substitution ONLY Penalise "(free) radical mechanism"
 - (ii) Initiation

 $\mathsf{CI}_{\scriptscriptstyle 2} \to 2\mathsf{CI} \bullet$

Penalise absence of dot once only.

First propagation

 $\label{eq:closed} \begin{array}{l} \mathsf{Cl} \bullet + \mathsf{CH}_3\mathsf{CH}_2\mathsf{CH}_3 \rightarrow \bullet\mathsf{CH}_2\mathsf{CH}_2\mathsf{CH}_3 + \mathsf{HCl} \\ \mathsf{OR} \ \mathsf{C}_3\mathsf{H}_8 \end{array}$

Penalise incorrect position of dot on propyl radical once only. Penalise C_3H_7 • once only

Second propagation

 $\mathsf{Cl}_{\scriptscriptstyle 2} + {}^{\bullet}\mathsf{CH}_{\scriptscriptstyle 2}\mathsf{CH}_{\scriptscriptstyle 2}\mathsf{CH}_{\scriptscriptstyle 3} \to \mathsf{CH}_{\scriptscriptstyle 3}\mathsf{CH}_{\scriptscriptstyle 2}\mathsf{CH}_{\scriptscriptstyle 2}\mathsf{CI} + \mathsf{Cl}{}^{\bullet}$

OR

C₃H₇Cl

Accept $CH_3CH_2CH_2$ • with the radical dot above/below/to the side of <u>the last carbon</u>.

Termination (must make C₆H₁₄)

2 •CH₂CH₂CH₃ \rightarrow C₆H₁₄ or CH₃CH₂CH₂CH₂CH₂CH₃

Use of the secondary free radical might gain 3 of the four marks

4

(e) $M_r = \underline{44.06352}$ (for propane) $M_r = \underline{43.98982}$ (for carbon dioxide) *Mark independently*

M1 a correct value for <u>both</u> of these <u>*M*, values</u>.

M2 a statement or idea that two peaks appear (in the mass spectrum)

OR

two molecular ions are seen (in the mass spectrum).

2

1

[12]

M2. (a) (i) M1 pentan-3-one only

M2 CH₃CH₂CH₂COCH₃ (insist on C=O being drawn out) (penalise use of C₃H₇)


1 (ii) aldehyde (CH₃)₂CHCH₂CHO 1 ketone (CH₃)₂CH<u>CO</u>CH₃ 1 (insist on a clear structure for the C=O of the functional groups, but do not be too harsh on the vertical bonds between carbon atom son this occasion) (If both structures correct, but wrong way around, award one mark) (ignore names) $CH_{3}CH_{2}CH_{2}CH_{2}CHO + [O] \rightarrow CH_{3}CH_{2}CH_{2}CH_{2}COOH$ (b) (i) (accept C_4H_9CHO going to C_4H_9COOH) (insist on a balanced equation – for example do not credit [O] over the arrow alone) 1 (ii) pentanoic acid (credit pentan-1-oic acid) 1 CH₃CH₂CH₂CH₂CH₂OH OR pentan-1-ol (c) (i) (If both a structure and a formula are given, credit either correct one of these provided the other is a good, if *imperfect, attempt)* 1 (ii) Primary (credit 1° or 1) 1

[8]


(b) Orange to green;


Need full colour change to score mark.

M4. (a) M1 <u>Displayed formula</u> for butan-2-ol



M1 displayed formula <u>must</u> have all bonds drawn out, including the O—H but ignore angles Penalise "sticks"





M2 structure must be clearly identifiable as 2-methylpropan-2-ol and may be drawn in a variety of ways.

M3 Alcohol **Y** is named <u>(2)-methylpropan-1-ol</u> ONLY *M3 <u>must be correct name</u>, but ignore structures*

3

- (b) M1 The infrared spectrum shows an <u>absorption/peak in the range</u> <u>3230 to 3550</u> (cm⁻¹)(which supports the idea that an alcohol is present) In M1, allow the words "dip", "spike", "low transmittance" and "trough" as alternatives for absorption.
 - M2 Reference to the 'fingerprint region' or below 1500 (cm⁻¹)

M3 <u>Match with</u> or <u>same as</u> known sample/database spectra Check the spectrum to see if alcohol OH is labelled and credit.

OR

M2 Run infrared spectra (of the alcohols)

M3 Find which one matches or is the same as this spectrum.

(c) **M1** balanced equation

 $C_6H_{12}O_6 \rightarrow CH_3CH_2CH_2CH_2OH + 2CO_2 + H_2O$ or C_4H_9OH Or multiples for M1 and M3 In M1 and M3 penalise use of $C_4H_{10}O$ or butan-2-ol once only

M2 Any one from

- <u>excess/adequate/sufficient/correct amount of/enough/plenty/</u> <u>a good supply</u> of oxygen or air
- good mixing of the fuel and air/oxygen For M2, do <u>not</u> accept simply "oxygen" or "air" alone Ignore reference to "temperature"

M3 CH₃CH₂CH₂CH₂OH + $6O_2 \rightarrow 4CO_2 + 5H_2O$ or C₄H₉OH

M4 A biofuel is a fuel produced from (renewable) biological (re)source(s)

OR

(renewable) (re)source(s) <u>from</u> (a specified) <u>plant(s)/fruit(s)/tree(s)</u> In M4 Ignore references to "carbon neutral" Ignore "sugar" and "glucose"

(d) **M1** butan-1-ol is a <u>primary or 1°</u> (alcohol)

M2 Displayed formula (ONLY) for butanal CH₃CH₂CH₂CHO

M3 Displayed formula (ONLY) for butanoic acid CH₃CH₂CH₂COOH

M2 and M3 displayed formula must have all bonds drawn out including the O—H but ignore angles. If butanal and butanoic acid formulae are <u>both</u> correctly given but not displayed, credit one mark out of two.


M4 Oxidation (oxidised) OR Redox

M5 orange to green


Both colours required for M5 Ignore states


[15]

4









[12]