

Q1. Which of these atoms has the largest atomic radius?

- A Ar
- B Cl
- C Mg
- D Na

(Total 1 mark)

Q2.(a) Table 1 shows some data about fundamental particles in an atom.

Table 1

Particle	proton	neutron	electron
Mass / g	1.6725×10^{-24}	1.6748×10^{-24}	0.0009×10^{-24}

(i) An atom of hydrogen can be represented as H

Use data from Table 1 to calculate the mass of this hydrogen atom.

.....

(1)

(ii) Which **one** of the following is a fundamental particle that would **not** be deflected by an electric field?

- A electron
- B neutron
- C proton

Write the correct letter, **A**, **B** or **C**, in the box.

(1)

(b) A naturally occurring sample of the element boron has a relative atomic mass of 10.8.

In this sample, boron exists as two isotopes, ^{10}B and ^{11}B

(i) Calculate the percentage abundance of ^{10}B in this naturally occurring sample of boron.

.....
.....
.....
.....
.....

(2)

(ii) State, in terms of fundamental particles, why the isotopes ^{10}B and ^{11}B have similar chemical reactions.

.....
.....
.....

(1)

(c) Complete **Table 2** by suggesting a value for the third ionisation energy of boron.

Table 2

	First	Second	Third	Fourth	Fifth
Ionisation energy / kJ mol^{-1}	799	2420		25 000	32 800

(1)

(d) Write an equation to show the process that occurs when the **second** ionisation energy of boron is measured. Include state symbols in your equation.

.....

(1)

(e) Explain why the second ionisation energy of boron is higher than the first ionisation energy of boron.

.....
.....

(1)
(Total 8 marks)

Q3. In 1913 Niels Bohr proposed a model of the atom with a central nucleus, made up of protons and neutrons, around which electrons moved in orbits. After further research, the model was refined when the existence of energy levels and sub-levels was recognised.

(a) Complete the following table for the particles in the nucleus.

Particle	Relative charge	Relative mass
proton		
neutron		

(2)

(b) State the block in the Periodic Table to which the element tungsten, W, belongs.

.....

(1)

(c) Isotopes of tungsten include ^{82}W and ^{86}W

(i) Deduce the number of protons in ^{82}W

.....

(1)

(ii) Deduce the number of neutrons in ^{86}W

.....

(1)

(d) In order to detect the isotopes of tungsten using a mass spectrometer, a sample containing the isotopes must be vaporised and then ionised.

(i) Give **two** reasons why the sample must be ionised.

1

2

(2)

- (ii) State what can be adjusted in the mass spectrometer to enable ions formed by the different isotopes to be directed onto the detector.

.....

(1)

- (e) State and explain the difference, if any, between the chemical properties of the isotopes ^{82}W and ^{86}W

Difference

Explanation

.....

(2)

- (f) The table below gives the relative abundance of each isotope in the mass spectrum of a sample of tungsten.

m/z	182	183	184	186
Relative abundance /%	26.4	14.3	30.7	28.6

Use the data above to calculate a value for the relative atomic mass of this sample of tungsten. Give your answer to 2 decimal places.

.....

.....

.....

(2)

(Total 12 marks)

Q4. In one model of atomic structure, the atom has a nucleus surrounded by electrons in levels and sub-levels.

(a) Define the term *atomic number*.

.....

(1)

(b) Explain why atoms of an element may have different mass numbers.

.....

(1)

(c) The table below refers to a sample of krypton.

Relative <i>m/z</i>	82	83	84	86
Relative abundance / %	12	12	50	26

(i) Name an instrument which is used to measure the relative abundance of isotopes.

.....

(ii) Define the term *relative atomic mass*.

.....

(iii) Calculate the relative atomic mass of this sample of krypton.

.....

.....

.....

(5)

(d) Give the complete electronic configuration of krypton in terms of s, p and d sub-levels.

.....

(1)

- (e) In 1963, krypton was found to react with fluorine. State why this discovery was unexpected.

.....

(1)

- (f) Use a suitable model of atomic structure to explain the following experimental observations.

- (i) The first ionisation energy of krypton is greater than that of bromine.

.....

.....

- (ii) The first ionisation energy of aluminium is less than the first ionisation energy of magnesium.

.....

.....

(4)

(Total 13 marks)

- Q5.** (a) Complete the following table.

	Relative mass	Relative charge
Neutron		
Electron		

(2)

- (b) An atom has twice as many protons as, and four more neutrons than, an atom of ${}^9\text{Be}$. Deduce the symbol, including the mass number, of this atom.

.....

(2)


- (c) Draw the shape of a molecule of BeCl_2 and the shape of a molecule of Cl_2O . Show any lone pairs of electrons on the central atom. Name the shape of each molecule.



Name of shape Name of shape

(4)

- (d) The equation for the reaction between magnesium hydroxide and hydrochloric acid is shown below.



Calculate the volume, in cm^3 , of 1.00 mol dm^{-3} hydrochloric acid required to react completely with 1.00 g of magnesium hydroxide.

.....
.....
.....
.....
.....
.....

(4)

(Total 12 marks)