### physicsandmathstutor.com

Centre No.					Pape	r Refer	ence			Surname	Initial(s)
Candidate No.			6	6	6	5	/	0	1	Signature	

Paper Reference(s)

### 6665/01

# **Edexcel GCE**

## **Core Mathematics C3**

### **Advanced**

Thursday 15 January 2009 – Morning

Time: 1 hour 30 minutes

Materials required for examination
Mathematical Formulae (Green)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

### **Instructions to Candidates**

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer for each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

#### **Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

#### **Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy. ©2009 Edexcel Limited.

Printer's Log. No. H31123A

W850/R6665/57570 3/3/3/3



Examiner's use only

Team Leader's use only

Question Number Leave

Turn over

Total



1. (a) Find the value of  $\frac{dy}{dx}$  at the point where x = 2 on the curve with equation

 $y = x^2 \sqrt{(5x - 1)}.$ 

(b) Differentiate  $\frac{\sin 2x}{x^2}$  with respect to x.

**(4)** 

**(6)** 

2.

$$f(x) = \frac{2x+2}{x^2-2x-3} - \frac{x+1}{x-3}$$

(a) Express f(x) as a single fraction in its simplest form.

**(4)** 

(b) Hence show that  $f'(x) = \frac{2}{(x-3)^2}$ 

**(3)** 

4

**3.** 




Figure 1

Figure 1 shows the graph of y = f(x), 1 < x < 9. The points T(3, 5) and S(7, 2) are turning points on the graph.

Sketch, on separate diagrams, the graphs of

(a) 
$$y = 2f(x) - 4$$
,

**(3)** 

(b) 
$$y = |f(x)|$$
.

**(3)** 

Indicate on each diagram the coordinates of any turning points on your sketch.

1 1111	d the equation of the tangent to the curve $x = \cos(2y + \pi)$ at $\left(0, \frac{\pi}{4}\right)$ .	
Giv	e your answer in the form $y = ax + b$ , where a and b are constants to be found.	(6)

The functions f and g are defined by

$$f: x \mapsto 3x + \ln x, \quad x > 0, \quad x \in \mathbb{R}$$
  
 $g: x \mapsto e^{x^2}, \quad x \in \mathbb{R}$ 

(a) Write down the range of g.

**(1)** 

(b) Show that the composite function fg is defined by

fg: 
$$x \mapsto x^2 + 3e^{x^2}$$
,  $x \in \mathbb{R}$ .

**(2)** 

(c) Write down the range of fg.

**(1)** 

(d) Solve the equation  $\frac{d}{dx} [fg(x)] = x(xe^{x^2} + 2)$ .

**(6)** 




(a) (i) By writing  $3\theta = (2\theta + \theta)$ , show that

$$\sin 3\theta = 3\sin \theta - 4\sin^3\theta.$$

**(4)** 

(ii) Hence, or otherwise, for  $0 < \theta < \frac{\pi}{3}$ , solve

$$8\sin^3\theta - 6\sin\theta + 1 = 0.$$

Give your answers in terms of  $\pi$ .

**(5)** 

(b) Using  $\sin(\theta - \alpha) = \sin\theta\cos\alpha - \cos\theta\sin\alpha$ , or otherwise, show that

$$\sin 15^{\circ} = \frac{1}{4}(\sqrt{6} - \sqrt{2}).$$

**(4)** 

uestion 6 continued	



7.

$$f(x) = 3xe^x - 1$$

The curve with equation y = f(x) has a turning point P.

(a) Find the exact coordinates of P.

**(5)** 

The equation f(x) = 0 has a root between x = 0.25 and x = 0.3

(b) Use the iterative formula

$$x_{n+1} = \frac{1}{3} e^{-x_n}$$

with  $x_0 = 0.25$  to find, to 4 decimal places, the values of  $x_1$ ,  $x_2$  and  $x_3$ .

**(3)** 

(c) By choosing a suitable interval, show that a root of f(x) = 0 is x = 0.2576 correct to 4 decimal places.

**(3)** 


nestion 7 continued	



**8.** (a) Express  $3 \cos \theta + 4 \sin \theta$  in the form  $R \cos(\theta - \alpha)$ , where R and  $\alpha$  are constants, R > 0 and  $0 < \alpha < 90^{\circ}$ .

**(4)** 

(b) Hence find the maximum value of  $3\cos\theta + 4\sin\theta$  and the smallest positive value of  $\theta$  for which this maximum occurs.

**(3)** 

The temperature, f(t), of a warehouse is modelled using the equation

$$f(t) = 10 + 3\cos(15t)^{\circ} + 4\sin(15t)^{\circ}$$
,

where *t* is the time in hours from midday and  $0 \le t < 24$ .

(c) Calculate the minimum temperature of the warehouse as given by this model.

**(2)** 

(d) Find the value of t when this minimum temperature occurs.

(3)


Question 8 continued		blan
		Q
	(Total 12 marks)	
	TOTAL FOR PAPER: 75 MARKS	
	END	

