

# Mark Scheme (Results)

## Summer 2009

GCE

**GCE Mathematics (6665/01)**

**June 2009**  
**6665 Core Mathematics C3**  
**Mark Scheme**



Question Number	Scheme	Marks
Q1 (a)	<p>Iterative formula: <math>x_{n+1} = \frac{2}{(x_n)^2} + 2</math>, <math>x_0 = 2.5</math></p> <p><math>x_1 = \frac{2}{(2.5)^2} + 2</math></p> <p><math>x_1 = 2.32</math></p> <p><math>x_2 = 2.371581451\dots</math></p> <p><math>x_3 = 2.355593575\dots</math></p> <p><math>x_4 = 2.360436923\dots</math></p> <p>An attempt to substitute <math>x_0 = 2.5</math> into the iterative formula. Can be implied by <math>x_1 = 2.32</math> or 2.320 Both <math>x_1 = 2.32(0)</math> and <math>x_2 = \text{awrt } 2.372</math> Both <math>x_3 = \text{awrt } 2.356</math> and <math>x_4 = \text{awrt } 2.360</math> or 2.36</p>	M1  A1  A1 cso (3)
(b)	<p>Let <math>f(x) = -x^3 + 2x^2 + 2 = 0</math></p> <p><math>f(2.3585) = 0.00583577\dots</math></p> <p><math>f(2.3595) = -0.00142286\dots</math></p> <p>Sign change (and <math>f(x)</math> is continuous) therefore a root <math>\alpha</math> is such that <math>\alpha \in (2.3585, 2.3595) \Rightarrow \alpha = 2.359</math> (3 dp)</p> <p>Choose suitable interval for <math>x</math>, e.g. [2.3585, 2.3595] or tighter any one value awrt 1 sf or truncated 1 sf both values correct, sign change and conclusion</p> <p>At a minimum, both values must be correct to 1sf or truncated 1sf, candidate states "change of sign, hence root".</p>	M1  dM1  A1 (3)  [6]

Question Number	Scheme	Marks
Q2 (a)	$\cos^2 \theta + \sin^2 \theta = 1 \quad (\div \cos^2 \theta)$ $\frac{\cos^2 \theta}{\cos^2 \theta} + \frac{\sin^2 \theta}{\cos^2 \theta} = \frac{1}{\cos^2 \theta}$ <p style="text-align: right;">Dividing <math>\cos^2 \theta + \sin^2 \theta = 1</math> by <math>\cos^2 \theta</math> to give <u>underlined</u> equation.</p> $1 + \tan^2 \theta = \sec^2 \theta$ $\tan^2 \theta = \sec^2 \theta - 1 \quad (\text{as required}) \quad \mathbf{AG}$ <p style="text-align: right;">Complete proof. No errors seen.</p>	M1 A1 cso (2)
(b)	$2\tan^2 \theta + 4\sec \theta + \sec^2 \theta = 2, \quad (\text{eqn } *) \quad 0 \leq \theta < 360^\circ$ $2(\sec^2 \theta - 1) + 4\sec \theta + \sec^2 \theta = 2$ <p style="text-align: right;">Substituting <math>\tan^2 \theta = \sec^2 \theta - 1</math> into eqn * to get a quadratic in <math>\sec \theta</math> only</p> $2\sec^2 \theta - 2 + 4\sec \theta + \sec^2 \theta = 2$ $\underline{3\sec^2 \theta + 4\sec \theta - 4 = 0}$ <p style="text-align: right;">Forming a three term “one sided” quadratic expression in <math>\sec \theta</math>.</p> $(\sec \theta + 2)(3\sec \theta - 2) = 0$ <p style="text-align: right;">Attempt to factorise or solve a quadratic.</p> $\sec \theta = -2 \quad \text{or} \quad \sec \theta = \frac{2}{3}$ $\frac{1}{\cos \theta} = -2 \quad \text{or} \quad \frac{1}{\cos \theta} = \frac{2}{3}$ $\underline{\cos \theta = -\frac{1}{2}}; \quad \text{or} \quad \cos \theta = \frac{3}{2}$ <p style="text-align: right;"><math>\underline{\cos \theta = -\frac{1}{2}}</math> A1;</p> $\alpha = 120^\circ \quad \text{or} \quad \alpha = \text{no solutions}$ $\theta_1 = \underline{120^\circ} \quad \underline{120^\circ} \quad \mathbf{A1}$ $\theta_2 = 240^\circ$ <p style="text-align: right;"><math>\underline{240^\circ}</math> or <math>\theta_2 = 360^\circ - \theta_1</math> when solving using <math>\cos \theta = \dots</math> B1 ✓</p> <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> Note the final A1 mark has been changed to a B1 mark. </div>	M1 M1 M1 M1 M1 A1; A1 B1 ✓ (6) [8]

Question Number	Scheme	Marks
Q3	$P = 80e^{\frac{t}{5}}$	
(a)	$t = 0 \Rightarrow P = 80e^{\frac{0}{5}} = 80(1) = \underline{80}$	<u>80</u> B1 (1)
(b)	$P = 1000 \Rightarrow 1000 = 80e^{\frac{t}{5}} \Rightarrow \frac{1000}{80} = e^{\frac{t}{5}}$ $\therefore t = 5 \ln\left(\frac{1000}{80}\right)$ $t = 12.6286\dots$	Substitutes $P = 1000$ and rearranges equation to make $e^{\frac{t}{5}}$ the subject.  <div style="border: 1px solid black; padding: 2px; display: inline-block;">awrt 12.6 or 13 years</div> Note $t = 12$ or $t = \text{awrt } 12.6 \Rightarrow t = 12$ will score A0
(c)	$\frac{dP}{dt} = 16e^{\frac{t}{5}}$	$ke^{\frac{1}{5}t}$ and $k \neq 80$ . $16e^{\frac{1}{5}t}$
(d)	$50 = 16e^{\frac{t}{5}}$ $\therefore t = 5 \ln\left(\frac{50}{16}\right)$ $\{= 5.69717\dots\}$ $P = 80e^{\frac{1}{5}\left(5 \ln\left(\frac{50}{16}\right)\right)}$ or $P = 80e^{\frac{1}{5}(5.69717\dots)}$ $P = \frac{80(50)}{16} = \underline{250}$	Using $50 = \frac{dP}{dt}$ and an attempt to solve to find the value of $t$ or $\frac{t}{5}$ .  Substitutes their value of $t$ back into the equation for $P$ .  <u>250</u> or awrt 250

Question Number	Scheme	Marks
Q4 (i)(a)	$y = x^2 \cos 3x$ Apply product rule: $\left\{ \begin{array}{l} u = x^2 \quad v = \cos 3x \\ \frac{du}{dx} = 2x \quad \frac{dv}{dx} = -3 \sin 3x \end{array} \right\}$ $\frac{dy}{dx} = 2x \cos 3x - 3x^2 \sin 3x$ <p style="text-align: right;">(3)</p>	M1 A1 A1
(b)	$y = \frac{\ln(x^2 + 1)}{x^2 + 1}$ $u = \ln(x^2 + 1) \Rightarrow \frac{du}{dx} = \frac{2x}{x^2 + 1}$ Apply quotient rule: $\left\{ \begin{array}{l} u = \ln(x^2 + 1) \quad v = x^2 + 1 \\ \frac{du}{dx} = \frac{2x}{x^2 + 1} \quad \frac{dv}{dx} = 2x \end{array} \right\}$ $\frac{dy}{dx} = \frac{\left(\frac{2x}{x^2 + 1}\right)(x^2 + 1) - 2x \ln(x^2 + 1)}{(x^2 + 1)^2}$ $\left\{ \frac{dy}{dx} = \frac{2x - 2x \ln(x^2 + 1)}{(x^2 + 1)^2} \right\}$ <p style="text-align: right;">(4)</p>	M1 A1 M1 A1 M1 A1

Question Number	Scheme	Marks
(ii)	$y = \sqrt{4x+1}, \quad x > -\frac{1}{4}$ At $P$ , $y = \sqrt{4(2)+1} = \underline{\sqrt{9}} = 3$ $\frac{dy}{dx} = \frac{1}{2}(4x+1)^{-\frac{1}{2}}(4)$ $\frac{dy}{dx} = \frac{2}{(4x+1)^{\frac{1}{2}}}$ At $P$ , $\frac{dy}{dx} = \frac{2}{(4(2)+1)^{\frac{1}{2}}}$ Hence $m(T) = \frac{2}{3}$  Either T: $y - 3 = \frac{2}{3}(x - 2)$ ; or $y = \frac{2}{3}x + c$ and $3 = \frac{2}{3}(2) + c \Rightarrow c = 3 - \frac{4}{3} = \frac{5}{3}$ ;  Either T: $3y - 9 = 2(x - 2)$ ; T: $3y - 9 = 2x - 4$ T: $2x - 3y + 5 = 0$  or T: $y = \frac{2}{3}x + \frac{5}{3}$ T: $3y = 2x + 5$ T: $2x - 3y + 5 = 0$	B1 M1* A1 aef M1 dM1*; A1 (6) [13]

Question Number	Scheme	Marks
Q5 (a)	 <p>Curve retains shape when <math>x &gt; \frac{1}{2} \ln k</math></p> <p>Curve reflects through the <math>x</math>-axis when <math>x &lt; \frac{1}{2} \ln k</math></p> <p><math>(0, k-1)</math> and <math>(\frac{1}{2} \ln k, 0)</math> marked in the correct positions.</p>	B1 B1 B1
(b)	 <p>Correct shape of curve. The curve should be contained in quadrants 1, 2 and 3 (Ignore asymptote)</p> <p><math>(1-k, 0)</math> and <math>(0, \frac{1}{2} \ln k)</math></p>	B1 B1
(c)	<p>Range of <math>f</math>: <math>f(x) &gt; -k</math> or <math>y &gt; -k</math> or <math>(-k, \infty)</math></p> <p>Either <math>f(x) &gt; -k</math> or <math>y &gt; -k</math> or <math>(-k, \infty)</math> or <math>f &gt; -k</math> or Range <math>&gt; -k</math>.</p>	B1
(d)	$\begin{aligned} y &= e^{2x} - k \Rightarrow y + k = e^{2x} \\ &\Rightarrow \ln(y + k) = 2x \\ &\Rightarrow \frac{1}{2} \ln(y + k) = x \end{aligned}$ <p>Hence <math>f^{-1}(x) = \frac{1}{2} \ln(x + k)</math></p> <p>Attempt to make <math>x</math> (or swapped <math>y</math>) the subject</p> <p>Makes <math>e^{2x}</math> the subject and takes <math>\ln</math> of both sides</p>	M1 M1 A1 cao
(e)	<p><math>f^{-1}(x)</math>: Domain: <math>x &gt; -k</math> or <math>(-k, \infty)</math></p> <p>Either <math>x &gt; -k</math> or <math>(-k, \infty)</math> or Domain <math>&gt; -k</math> or <math>x</math> “ft one sided inequality” their part (c) RANGE answer</p>	B1 ✓ (1) [10]

Question Number	Scheme	Marks
Q6 (a)	$A = B \Rightarrow \cos(A + A) = \cos 2A = \underline{\cos A \cos A - \sin A \sin A}$ $\cos 2A = \cos^2 A - \sin^2 A$ and $\cos^2 A + \sin^2 A = 1$ gives $\underline{\cos 2A} = 1 - \sin^2 A - \sin^2 A = \underline{1 - 2\sin^2 A}$ (as required) Complete proof, with a link between LHS and RHS. No errors seen.	M1 A1 AG (2)
(b)	$C_1 = C_2 \Rightarrow 3\sin 2x = 4\sin^2 x - 2\cos 2x$ $3\sin 2x = 4\left(\frac{1 - \cos 2x}{2}\right) - 2\cos 2x$ $3\sin 2x = 2(1 - \cos 2x) - 2\cos 2x$ $3\sin 2x = 2 - 2\cos 2x - 2\cos 2x$ $3\sin 2x + 4\cos 2x = 2$ Rearranges to give correct result	M1 M1 A1 AG (3)
(c)	$3\sin 2x + 4\cos 2x = R\cos(2x - \alpha)$ $3\sin 2x + 4\cos 2x = R\cos 2x \cos \alpha + R\sin 2x \sin \alpha$ Equate $\sin 2x$ : $3 = R \sin \alpha$ Equate $\cos 2x$ : $4 = R \cos \alpha$ $R = \sqrt{3^2 + 4^2} ;= \sqrt{25} = 5$ $\tan \alpha = \frac{3}{4} \Rightarrow \alpha = 36.86989765...^\circ$ Hence, $3\sin 2x + 4\cos 2x = 5\cos(2x - 36.87)$	B1 M1 A1 (3)

Question Number	Scheme	Marks
(d)	$3\sin 2x + 4\cos 2x = 2$ $5\cos(2x - 36.87) = 2$ $\cos(2x - 36.87) = \frac{2}{5}$ $(2x - 36.87) = 66.42182\dots^\circ$ $(2x - 36.87) = 360 - 66.42182\dots^\circ$ <p>Hence, <math>x = 51.64591\dots^\circ, 165.22409\dots^\circ</math></p> <p style="text-align: right;">If there are any EXTRA solutions inside the range <math>0 \leq x &lt; 180^\circ</math> then withhold the final accuracy mark. Also ignore EXTRA solutions outside the range <math>0 \leq x &lt; 180^\circ</math>.</p>	<span style="float: right;">M1</span> <span style="float: right;">A1</span> <span style="float: right;">A1</span> <span style="float: right;">A1</span> <span style="float: right;">(4)</span> <span style="float: right;">[12]</span>

Question Number	Scheme	Marks
Q7	$f(x) = 1 - \frac{2}{(x+4)} + \frac{x-8}{(x-2)(x+4)}$ $x \in \mathbb{R}, x \neq -4, x \neq 2.$	
(a)	$f(x) = \frac{(x-2)(x+4) - 2(x-2) + x-8}{(x-2)(x+4)}$ $= \frac{x^2 + 2x - 8 - 2x + 4 + x - 8}{(x-2)(x+4)}$ $= \frac{x^2 + x - 12}{[(x+4)(x-2)]}$ $= \frac{(x+4)(x-3)}{[(x+4)(x-2)]}$ $= \frac{(x-3)}{(x-2)}$	An attempt to combine to one fraction M1 Correct result of combining all three fractions A1 Simplifies to give the correct numerator. Ignore omission of denominator A1 An attempt to factorise the numerator. dM1 Correct result A1 cso AG (5)
(b)	$g(x) = \frac{e^x - 3}{e^x - 2} \quad x \in \mathbb{R}, x \neq \ln 2.$ <p>Apply quotient rule: <math>\left\{ \begin{array}{l} u = e^x - 3 \\ \frac{du}{dx} = e^x \end{array} \quad \begin{array}{l} v = e^x - 2 \\ \frac{dv}{dx} = e^x \end{array} \right\}</math></p> $g'(x) = \frac{e^x(e^x - 2) - e^x(e^x - 3)}{(e^x - 2)^2}$ $= \frac{e^{2x} - 2e^x - e^{2x} + 3e^x}{(e^x - 2)^2}$ $= \frac{e^x}{(e^x - 2)^2}$	Applying $\frac{vu' - uv'}{v^2}$ M1 Correct differentiation A1 Correct result A1 AG cso (3)

Question Number	Scheme	Marks
(c)	$g'(x) = 1 \Rightarrow \frac{e^x}{(e^x - 2)^2} = 1$ $e^x = (e^x - 2)^2$ $e^x = e^{2x} - 2e^x - 2e^x + 4$ $\underline{e^{2x} - 5e^x + 4 = 0}$ $(e^x - 4)(e^x - 1) = 0$ $e^x = 4 \text{ or } e^x = 1$ $x = \ln 4 \text{ or } x = 0$ <p style="text-align: right;">Puts their differentiated numerator equal to their denominator.</p> <p style="text-align: right;">Attempt to factorise or solve quadratic in <math>e^x</math></p> <p style="text-align: right;">both <math>x = 0, \ln 4</math></p>	M1 A1 M1 A1 (4) [12]

Question Number	Scheme	Marks
Q8 (a)	$\sin 2x = \underline{2 \sin x \cos x}$	B1 aef (1)
(b)	$\text{cosec } x - 8 \cos x = 0, \quad 0 < x < \pi$ $\frac{1}{\sin x} - 8 \cos x = 0$ $\frac{1}{\sin x} = 8 \cos x$ $1 = 8 \sin x \cos x$ $1 = 4(2 \sin x \cos x)$ $1 = 4 \sin 2x$ $\underline{\sin 2x = \frac{1}{4}}$	M1
	$\sin 2x = k, \text{ where } -1 < k < 1 \text{ and}$ $k \neq 0$ $\underline{\sin 2x = \frac{1}{4}}$	M1
Radians	$2x = \{0.25268..., 2.88891...\}$	A1
Degrees	$2x = \{14.4775..., 165.5225...\}$	
Radians	$x = \{0.12634..., 1.44445...\}$	A1
Degrees	$x = \{7.23875..., 82.76124...\}$	
	Either arwt 7.24 or 82.76 or 0.13 or 1.44 or 1.45 or awrt $0.04\pi$ or awrt 0.46. Both <u>0.13</u> and <u>1.44</u>	
	Solutions for the final two A marks must be given in $x$ only. If there are any EXTRA solutions inside the range $0 < x < \pi$ then withhold the final accuracy mark. Also ignore EXTRA solutions outside the range $0 < x < \pi$ .	A1 cao (5)