

**C3****TRIGONOMETRY****Worksheet D**

- 1**    **a** Write down the identities for  $\sin(A + B)$  and  $\cos(A + B)$ .  
**b** Use these identities to obtain similar identities for  $\sin(A - B)$  and  $\cos(A - B)$ .  
**c** Use the above identities to obtain similar identities for  $\tan(A + B)$  and  $\tan(A - B)$ .
- 2**    Express each of the following in the form  $\sin \alpha$ , where  $\alpha$  is acute.
- |  |  |
|--|--|
| <b>a</b> $\sin 10^\circ \cos 30^\circ + \cos 10^\circ \sin 30^\circ$ | <b>b</b> $\sin 67^\circ \cos 18^\circ - \cos 67^\circ \sin 18^\circ$ |
| <b>c</b> $\sin 62^\circ \cos 74^\circ + \cos 62^\circ \sin 74^\circ$ | <b>d</b> $\cos 14^\circ \cos 39^\circ - \sin 14^\circ \sin 39^\circ$ |
- 3**    Express as a single trigonometric ratio
- |  |  |
|--|--|
| <b>a</b> $\cos A \cos 2A - \sin A \sin 2A$               | <b>b</b> $\sin 4A \cos B - \cos 4A \sin B$ |
| <b>c</b> $\frac{\tan 2A + \tan 5A}{1 - \tan 2A \tan 5A}$ | <b>d</b> $\cos A \cos 3A + \sin A \sin 3A$ |
- 4**    Find in exact form, with a rational denominator, the value of
- |                          |                           |                                   |                                    |
|--------------------------|---------------------------|-----------------------------------|------------------------------------|
| <b>a</b> $\sin 15^\circ$ | <b>b</b> $\sin 165^\circ$ | <b>c</b> $\text{cosec } 15^\circ$ | <b>d</b> $\cos 75^\circ$           |
| <b>e</b> $\cos 15^\circ$ | <b>f</b> $\sec 195^\circ$ | <b>g</b> $\tan 75^\circ$          | <b>h</b> $\text{cosec } 105^\circ$ |
- 5**    Find the maximum value that each expression can take and the smallest positive value of  $x$ , in degrees, for which this maximum occurs.
- |  |  |
|--|--|
| <b>a</b> $\cos x \cos 30^\circ + \sin x \sin 30^\circ$ | <b>b</b> $3 \sin x \cos 45^\circ + 3 \cos x \sin 45^\circ$   |
| <b>c</b> $\sin x \cos 67^\circ - \cos x \sin 67^\circ$ | <b>d</b> $4 \sin x \sin 108^\circ - 4 \cos x \cos 108^\circ$ |
- 6**    Find the minimum value that each expression can take and the smallest positive value of  $x$ , in radians in terms of  $\pi$ , for which this minimum occurs.
- |  |  |
|--|--|
| <b>a</b> $\sin x \cos \frac{\pi}{3} - \cos x \sin \frac{\pi}{3}$ | <b>b</b> $2 \cos x \cos \frac{\pi}{6} - 2 \sin x \sin \frac{\pi}{6}$ |
| <b>c</b> $\cos 4x \cos x + \sin 4x \sin x$                       | <b>d</b> $6 \sin 2x \cos 3x - 6 \sin 3x \cos 2x$                     |
- 7**    Given that  $\sin A = \frac{4}{5}$ ,  $0 < A < 90^\circ$  and that  $\cos B = \frac{2}{3}$ ,  $0 < B < 90^\circ$ , find without using a calculator the value of
- |                   |                   |                        |                        |
|-------------------|-------------------|------------------------|------------------------|
| <b>a</b> $\tan A$ | <b>b</b> $\sin B$ | <b>c</b> $\cos(A + B)$ | <b>d</b> $\sin(A + B)$ |
|-------------------|-------------------|------------------------|------------------------|
- 8**    Given that  $\text{cosec } C = \frac{5}{3}$ ,  $0 < C < 90^\circ$  and that  $\sin D = \frac{5}{13}$ ,  $90^\circ < D < 180^\circ$ , find without using a calculator the value of
- |                   |                   |                        |                        |
|-------------------|-------------------|------------------------|------------------------|
| <b>a</b> $\cos C$ | <b>b</b> $\cos D$ | <b>c</b> $\sin(C - D)$ | <b>d</b> $\sec(C - D)$ |
|-------------------|-------------------|------------------------|------------------------|
- 9**    Solve each equation for  $\theta$  in the interval  $0 \leq \theta \leq 360$ .  
Give your answers to 1 decimal place where appropriate.
- |  |  |
|--|--|
| <b>a</b> $\sin \theta^\circ \cos 15^\circ + \cos \theta^\circ \sin 15^\circ = 0.4$ | <b>b</b> $\frac{\tan 2\theta^\circ - \tan 60^\circ}{1 + \tan 2\theta^\circ \tan 60^\circ} = 1$ |
| <b>c</b> $\cos(\theta - 60)^\circ = \sin \theta^\circ$                             | <b>d</b> $2 \sin \theta^\circ + \sin(\theta + 45)^\circ = 0$                                   |
| <b>e</b> $\sin(\theta + 30)^\circ = \cos(\theta - 45)^\circ$                       | <b>f</b> $3 \cos(2\theta + 60)^\circ - \sin(2\theta - 30)^\circ = 0$                           |

**C3 TRIGONOMETRY***Worksheet D continued*

- 10** Find the value of  $k$  such that for all real values of  $x$

$$\cos(x + \frac{\pi}{3}) - \cos(x - \frac{\pi}{3}) \equiv k \sin x.$$

- 11** Prove each identity.

a  $\cos x - \cos(x - \frac{\pi}{3}) \equiv \cos(x + \frac{\pi}{3})$

b  $\sin(x - \frac{\pi}{6}) + \cos x \equiv \sin(x + \frac{\pi}{6})$

- 12** a Use the identity for  $\sin(A + B)$  to express  $\sin 2A$  in terms of  $\sin A$  and  $\cos A$ .  
 b Use the identity for  $\cos(A + B)$  to express  $\cos 2A$  in terms of  $\sin A$  and  $\cos A$ .  
 c Hence, express  $\cos 2A$  in terms of  
 i  $\cos A$       ii  $\sin A$   
 d Use the identity for  $\tan(A + B)$  to express  $\tan 2A$  in terms of  $\tan A$ .

- 13** Solve each equation for  $x$  in the interval  $0 \leq x \leq 360^\circ$ .

Give your answers to 1 decimal place where appropriate.

a  $\cos 2x + \cos x = 0$

b  $\sin 2x + \cos x = 0$

c  $2 \cos 2x = 7 \sin x$

d  $11 \cos x = 4 + 3 \cos 2x$

e  $\tan 2x - \tan x = 0$

f  $\sec x - 4 \sin x = 0$

g  $5 \sin 4x = 2 \sin 2x$

h  $2 \sin^2 x - \cos 2x - \cos x = 0$

- 14** Prove each identity.

a  $(\cos x + \sin x)^2 \equiv 1 + \sin 2x$

b  $\tan x (1 + \cos 2x) \equiv \sin 2x$

c  $\frac{2 \sin x}{2 \cos x - \sec x} \equiv \tan 2x$

d  $\tan x + \cot x \equiv 2 \operatorname{cosec} 2x$

e  $\operatorname{cosec} 2x - \cot 2x \equiv \tan x$

f  $(\cos x + \sin x)(\operatorname{cosec} x - \sec x) \equiv 2 \cot 2x$

g  $\frac{1 - \sin 2x}{\operatorname{cosec} x - 2 \cos x} \equiv \sin x$

h  $\cos 3x \equiv 4 \cos^3 x - 3 \cos x$

- 15** Use the double angle identities to prove that

a  $\cos x \equiv 2 \cos^2 \frac{x}{2} - 1$

b  $\sin^2 \frac{x}{2} \equiv \frac{1}{2}(1 - \cos x)$

- 16** a Given that  $\cos A = \frac{7}{9}$ ,  $0 < A < 90^\circ$ , find the exact value of  $\sin \frac{A}{2}$  without using a calculator.

- b Given that  $\cos B = -\frac{3}{8}$ ,  $90^\circ < B < 180^\circ$ , find the value of  $\cos \frac{B}{2}$ , giving your answer in the form  $k\sqrt{5}$ .

- 17** Prove each identity.

a  $\frac{2}{1 + \cos x} \equiv \sec^2 \frac{x}{2}$

b  $\frac{1 + \cos x}{1 - \cos x} \equiv \cot^2 \frac{x}{2}$