

Worked Solutions

Edexcel C4 Paper G

1. (a) $\frac{dy}{dx} = \frac{2\cos t}{-\sin t}$
when $t = \frac{\pi}{2}$, gradient = 0

(b) using $\cos^2 t + \sin^2 t = 1$,

$$x^2 + \left(\frac{y}{2}\right)^2 = 1$$

$$x^2 + \frac{y^2}{4} = 1$$

2. (a) $\frac{1}{x-2} + \frac{3}{2x+1} + \frac{1}{x+2}$ (using ‘cover up’ rule)

(b) $\int_3^4 \left(\frac{1}{x-2} + \frac{3}{2x+1} + \frac{1}{x+2} \right) dx$

$$= \left[\ln(x-2) + \frac{3}{2} \ln(2x+1) + \ln(x+2) \right]_3^4$$

$$= \ln 2 + \frac{3}{2} \ln 9 + \ln 6$$

$$- \left(\ln 1 + \frac{3}{2} \ln 7 + \ln 5 \right)$$

$$= \ln \left(\frac{2 \times 6}{5} \right) + \frac{3}{2} \ln \left(\frac{9}{7} \right)$$

$$= \ln \left(\frac{12}{5} \right) + \frac{3}{2} \ln \frac{9}{7}$$

3. (a) $(1+ax)^6 = 1 + 6ax + 15a^2x^2$

(2)

$$(b) (1+bx)(1+6ax+15a^2x^2) = 1 + 6ax + 15a^2x^2 + bx + 6abx^2$$

we have $6a + b = -9$...[A]

$$15a^2 + 6ab = 24 \quad \dots [B]$$

from equation [A] $b = -9 - 6a$

substitute in [B] $15a^2 + 6a(-9 - 6a) = 24$

Hence $a = -2, b = 3$

(3)

4. (a) $= \int x \frac{d}{dx} (\tan x) dx$ [By parts]

$$= x \tan x - \int \tan x \, dx = x \tan x + \ln \cos x + c$$

(b) $\int y^{-\frac{1}{2}} \, dy = \int x \sec^2 x \, dx \quad 2y^{\frac{1}{2}} = x \tan x + \ln \cos x + c$
 $y = 4, x = 0: 2\sqrt{4} = 0 + \ln 1 + c$

$$c = 4$$

$$2\sqrt{y} = x \tan x + \ln \cos x + 4$$

when $x = \frac{\pi}{4}, 2\sqrt{y} = \frac{\pi}{4} \cdot \tan \frac{\pi}{4} + \ln \frac{1}{\sqrt{2}} + 4$

$$2\sqrt{y} = \frac{\pi}{4} + \ln 2^{-\frac{1}{2}} + 4$$

$$2\sqrt{y} = \frac{\pi}{4} - \frac{1}{2} \ln 2 + 4$$

$$\sqrt{y} = \frac{\pi}{8} - \frac{1}{4} \ln 2 + 2$$

$$y = \left(\frac{\pi}{8} - \frac{1}{4} \ln 2 + 2 \right)^2$$

(5)

(3)

x	1	$1\frac{1}{2}$	2	$2\frac{1}{2}$	3
$\frac{12}{x}$	12	8	6	4.8	4

5. (a)

$$\text{(i) using two trapeziums, } I = \frac{1}{2} [12 + 4 + (2 \times 6)] = 14$$

(2)

$$\text{(ii) using four trapeziums, } I = \frac{1}{4} [12 + 4 + 2(8 + 6 + 4.8)] = 13.4$$


(2)

$$\text{(b) } \int_1^3 \frac{12}{x} dx = [12 \ln x]_1^3 = 12 \ln 3 = 13.1833\dots$$

$$\text{in (i) \% error} = \frac{14 - 13.1833}{13.1833} \times 100 = 6.2\%$$

$$\text{in (ii) \% error} = \frac{13.4 - 13.1833}{13.1833} \times 100 = 1.6\% \quad (5)$$

$$6. \text{ (a) } \vec{AB} = \begin{pmatrix} 2 \\ 5 \\ m-8 \end{pmatrix}$$



$$\vec{AO} \cdot \vec{AB} = 0$$

$$\begin{pmatrix} -1 \\ 2 \\ -8 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 5 \\ m-8 \end{pmatrix} = -2 + 10 - 8m + 64 = 0$$

$$m = 9$$

(4)

$$\text{(b) } \vec{OC} = \vec{AB} \quad (\text{OABC is a rectangle})$$

$$= \begin{pmatrix} 2 \\ 5 \\ 1 \end{pmatrix}$$

(1)

$$\text{(c) } \vec{AC} = \begin{pmatrix} 1 \\ 7 \\ -7 \end{pmatrix}$$

$$\text{equation of line } AC \text{ is } r = \begin{pmatrix} 1 \\ -2 \\ 8 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 7 \\ -7 \end{pmatrix}$$

(2)

$$\text{(d) } \vec{OB} = \begin{pmatrix} 3 \\ 3 \\ 9 \end{pmatrix}$$

let angle between diagonals be θ

$$\vec{AC} \cdot \vec{OB} = |\vec{AC}| \times |\vec{OB}| \cos \theta$$

$$\begin{pmatrix} 1 \\ 7 \\ -7 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 3 \\ 9 \end{pmatrix} = \sqrt{99}\sqrt{99} \cos \theta$$

$$3 + 21 - 63 = 99 \cos \theta$$

$$\theta = 113.2^\circ$$

acute angle between diagonals = 67° (nearest degree)

(3)

7. (a) $y = 12x - e^{\frac{1}{2}x}$

$$\frac{dy}{dx} = 12 - \frac{1}{2} e^{\frac{1}{2}x}$$

$$\frac{dy}{dx} = 0, \quad 24 = e^{\frac{1}{2}x}$$

$$x = 2 \ln 24$$

$$(b) y = 12 \times 2 \ln 24 - 24 = 24 \ln 24 - 24$$

$$(c) \frac{d^2y}{dx^2} = -\frac{1}{4} e^{\frac{1}{2}x}$$

so $\frac{d^2y}{dx^2} < 0$ and the stationary point is a maximum.

$$(d) \text{ area} = \int_2^4 \left(12x - e^{\frac{1}{2}x} \right) dx = \left[6x^2 - 2e^{\frac{1}{2}x} \right]_2^4$$

$$= 96 - 2e^2 - (24 - 2e) = 72 + 2e(1 - e)$$

(3)

(1)

(1)

(5)

8. (a) $4x - \left(x \frac{dy}{dx} + y \cdot 1 \right) + 2y \frac{dy}{dx} = 0$

$$\frac{dy}{dx}(2y - x) = y - 4x$$

$$\frac{dy}{dx} = \frac{y - 4x}{2y - x}$$

(4)

$$(b) \text{ at } (2, -6) \text{ gradient of curve} = \frac{-6 - 8}{-12 - 2} = 1$$

\therefore gradient of normal = -1

equation of normal is $y + 6 = -1(x - 2)$

$$x + y + 4 = 0$$

(3)

(c) gradient = 0 where $y = 4x$

substitute $y = 4x$ into equation of curve.

$$2x^2 - x \cdot 4x + 16x^2 = 56 \Rightarrow x = \pm 2$$

gradient = 0 at $(2, 8)$ and $(-2, -8)$

(5)