

FPI (int) June 13

1. The complex numbers z and w are given by

$$z = 8 + 3i, \quad w = -2i$$

Express in the form $a + bi$, where a and b are real constants,

(a) $z - w$,

(1)

(b) zw .

(2)

a) $8 + 5i$

b) $(8+3i)(-2i) = -16i - 6i^2 = 6 - 16i$

2. (i)

$$\mathbf{A} = \begin{pmatrix} 2k+1 & k \\ -3 & -5 \end{pmatrix}, \text{ where } k \text{ is a constant}$$

Given that

$$\mathbf{B} = \mathbf{A} + 3\mathbf{I}$$

where $\mathbf{I}$ is the 2×2 identity matrix, find

(a) $\mathbf{B}$ in terms of k ,

(2)

(b) the value of k for which $\mathbf{B}$ is singular.

(2)

(ii) Given that

$$\mathbf{C} = \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix}, \quad \mathbf{D} = \begin{pmatrix} 2 & -1 & 5 \end{pmatrix}$$

and

$$\mathbf{E} = \mathbf{CD}$$

find $\mathbf{E}$.

(2)

$$a) \mathbf{A} + 3\mathbf{I} = \begin{pmatrix} 2k+1 & k \\ -3 & -5 \end{pmatrix} + \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix} = \begin{pmatrix} 2k+4 & k \\ -3 & -2 \end{pmatrix}$$

$$b) \text{ Singular } \Rightarrow \det = ad - bc = 0$$

$$(2k+4)(-2) - (-3)(k) = 0 \Rightarrow -4k - 8 + 3k = 0 \\ \Rightarrow -k = 8 \Rightarrow \underline{k = -8}$$

$$ii) \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \begin{pmatrix} 2 & -1 & 5 \end{pmatrix} = \begin{pmatrix} 4 & -2 & 10 \\ -6 & 3 & -15 \\ 8 & -4 & 20 \end{pmatrix}$$

3.

$$f(x) = \frac{1}{2}x^4 - x^3 + x - 3$$


(a) Show that the equation $f(x) = 0$ has a root α between $x = 2$ and $x = 2.5$ (2)

(b) Starting with the interval $[2, 2.5]$ use interval bisection twice to find an interval of width 0.125 which contains α . (3)

The equation $f(x) = 0$ has a root β in the interval $[-2, -1]$.

(c) Taking -1.5 as a first approximation to β , apply the Newton-Raphson process once to $f(x)$ to obtain a second approximation to β .
Give your answer to 2 decimal places. (5)

a) $f(2) = -1$ $\therefore$ by sign change rule root $\alpha \in (2, 2.5]$
 $f(2.5) = 3.41$



c) $f'(x) = 2x^3 - 3x^2 + 1$ $x_1 = -1.5$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = -1.5 - \frac{\frac{45}{32}}{-\frac{25}{2}} = -1.39 \text{ (2dp)}$$

$\beta = \underline{\underline{-1.39}}$

4.

$$f(x) = (4x^2 + 9)(x^2 - 2x + 5)$$

(a) Find the four roots of $f(x) = 0$

(4)

(b) Show the four roots of $f(x) = 0$ on a single Argand diagram.

(2)

$$a) \quad 4x^2 + 9 = 0 \Rightarrow x^2 = -\frac{9}{4} \Rightarrow x = \pm \frac{3}{2}i$$


$$x^2 - 2x + 5 = 0 \Rightarrow (x-1)^2 - 1 = -5$$

$$\Rightarrow (x-1)^2 = -4$$

$$\Rightarrow x-1 = \pm 2i$$

$$\Rightarrow x = 1 \pm 2i$$

$$\therefore x = \frac{3}{2}i, -\frac{3}{2}i, 1+2i, 1-2i$$



5.




Figure 1

Figure 1 shows a rectangular hyperbola H with parametric equations

$$x = 3t, \quad y = \frac{3}{t}, \quad t \neq 0$$

The line L with equation $6y = 4x - 15$ intersects H at the point P and at the point Q as shown in Figure 1.

(a) Show that L intersects H where $4t^2 - 5t - 6 = 0$ (3)

(b) Hence, or otherwise, find the coordinates of points P and Q . (5)

$$a) \quad 6\left(\frac{3}{t}\right) = 4(3t) - 15 \Rightarrow \frac{18}{t} = 12t - 15 \quad (\times t)$$

$$\Rightarrow 18 = 12t^2 - 15t \quad (\div 3) \Rightarrow 4t^2 - 5t - 6 = 0 \quad \#$$

$$b) \quad (4t+3)(t-2) = 0 \Rightarrow t = -\frac{3}{4}, \quad t = 2.$$

$$t = 2, \quad P = \left(6, \frac{3}{2}\right) \quad t = -\frac{3}{4}, \quad Q = \left(-\frac{9}{4}, -4\right)$$

6.

$$\mathbf{A} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix}$$

The transformation represented by $\mathbf{B}$ followed by the transformation represented by $\mathbf{A}$ is equivalent to the transformation represented by $\mathbf{P}$.

(a) Find the matrix $\mathbf{P}$.

(2)

Triangle T is transformed to the triangle T' by the transformation represented by $\mathbf{P}$.

Given that the area of triangle T' is 24 square units,

(b) find the area of triangle T .

(3)

Triangle T' is transformed to the original triangle T by the matrix represented by $\mathbf{Q}$.

(c) Find the matrix $\mathbf{Q}$.

(2)

$$a) \mathbf{AB} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 4 \\ -2 & -3 \end{pmatrix}$$

$$b) \det \mathbf{P} = 1 \times 3 - 4 \times 2 = -3 + 8 = 5$$

$$\text{area of } T \times 5 = \text{area of } T' \therefore \text{area of } T = 4.8$$

$$c) \mathbf{Q} = \mathbf{P}^{-1} = \frac{1}{5} \begin{pmatrix} 3 & -4 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} \frac{3}{5} & -\frac{4}{5} \\ \frac{2}{5} & \frac{1}{5} \end{pmatrix}$$

7. The parabola C has equation $y^2 = 4ax$, where a is a positive constant.

The point $P(at^2, 2at)$ is a general point on C .

- (a) Show that the equation of the tangent to C at $P(at^2, 2at)$ is

$$ty = x + at^2 \quad (4)$$

The tangent to C at P meets the y -axis at a point Q .

- (b) Find the coordinates of Q . (1)

Given that the point S is the focus of C ,

- (c) show that PQ is perpendicular to SQ . (3)


$$2y \frac{dy}{dx} = 4a \Rightarrow \frac{dy}{dx} = \frac{2a}{y} \quad m_t \Big|_{y=2at} = \frac{2a}{2at} = \frac{1}{t}$$

$$(at^2, 2at): \quad y - 2at = \frac{1}{t}(x - at^2) \Rightarrow ty - 2at^2 = x - at^2 \\ \therefore ty = x + at^2 \quad \#$$

$$b) \quad x=0, \quad y=at^2 \Rightarrow y=at \quad Q(0, at)$$

$$c) \quad S(a, 0)$$

$$Q(0, at)$$



$$\therefore m = \frac{-at}{a} = -t$$

$$m_{QS} \times m_{PQ} = -t \times \frac{1}{t} = -1$$

$\therefore QS$ and PQ are perp.

8. (a) Prove by induction, that for $n \in \mathbb{Z}^+$,

$$\sum_{r=1}^n r(2r-1) = \frac{1}{6} n(n+1)(4n-1)$$

(6)

(b) Hence, show that

$$\sum_{r=n+1}^{3n} r(2r-1) = \frac{1}{3} n(an^2 + bn + c)$$

where a, b and c are integers to be found.

(4)

a) $\sum r(2r-1) = 1(2-1) = 1$ when $n=1$

$$\frac{1}{6} n(n+1)(4n-1) = \frac{1}{6} (1)(2)(3) = 1 \text{ when } n=1$$

$\therefore$ true for $n=1$

assume true when $n=k$.

$$\begin{aligned} \sum_{r=1}^{k+1} r(2r-1) &= \frac{1}{6} (k+1)(k+1+1)(4(k+1)-1) \\ &= \frac{1}{6} (k+1)(k+2)(4k+3) \end{aligned}$$

$$\begin{aligned} \sum_{r=1}^{k+1} r(2r-1) &= (k+1)(2(k+1)-1) + \sum_{r=1}^k r(2r-1) \\ &= (k+1)(2k+1) + \frac{1}{6} k(k+1)(4k-1) \\ &= \frac{1}{6} (k+1) [6(2k+1) + k(4k-1)] \\ &= \frac{1}{6} (k+1) [4k^2 + 11k + 6] = \frac{1}{6} (k+1)(k+2)(4k+3) \end{aligned}$$

$\therefore$ true for $n=1$, true for $n=k+1$ if true for $n \leq k$

$\therefore$ by induction true for all $n \in \mathbb{Z}^+$

$$b) \sum_{r=n+1}^{3n} r(2r-1) = \sum_{r=1}^{3n} r(2r-1) - \sum_{r=1}^n r(2r-1)$$

$$= \frac{1}{6}(3n)(3n+1)(4(3n)-1) - \frac{1}{6}n(n+1)(4n-1)$$

$$= \frac{1}{6}(3n)(3n+1)(12n-1) - \frac{1}{6}n(n+1)(4n-1)$$

$$= \frac{1}{6}n[(9n+3)(12n-1) - (n+1)(4n-1)]$$

$$= \frac{1}{6}n[108n^2 + 27n - 3 - 4n^2 - 3n + 1]$$

$$= \frac{1}{6}n[104n^2 + 24n - 2]$$

$$= \frac{2}{6}n[52n^2 + 12n - 1]$$

$$= \frac{1}{3}n(52n^2 + 12n - 1)$$

9. The complex number w is given by

$$w = 10 - 5i$$

(a) Find $|w|$.

(1)

(b) Find $\arg w$, giving your answer in radians to 2 decimal places.

(2)

The complex numbers z and w satisfy the equation

$$(2 + i)(z + 3i) = w$$

(c) Use algebra to find z , giving your answer in the form $a + bi$, where a and b are real numbers.

(4)


Given that

$$\arg(\lambda + 9i + w) = \frac{\pi}{4}$$

where λ is a real constant,

(d) find the value of λ .

a) $|w| = \sqrt{10^2 + 5^2} = \sqrt{125}$

b)  $\arg w = -\tan^{-1}\left(\frac{5}{10}\right)$
 $\therefore \arg w = -0.46$

c) $(2+i)(a+(3+b)i) = 2a + (a+6+2b)i - (3+b)$

$$\Rightarrow (2a-3-b) + (a+6+2b)i = 10-5i$$

$$\Rightarrow \begin{aligned} 2a-b &= 13 \\ a+2b &= -11 \end{aligned} \Rightarrow \begin{aligned} 4a-2b &= 26 \\ a+2b &= -11 \end{aligned}$$


$$\therefore a = 3$$

$$b = -7$$

$$\underline{5a = 15}$$

d) $\arg(\lambda + 9i + 10 - 5i)$

$$= \arg(\lambda + 10 + 4i) = \frac{\pi}{4}$$



$$\therefore \lambda + 10 = 4 \Rightarrow \lambda = -6$$

10. (i) Use the standard results for $\sum_{r=1}^n r^3$ and $\sum_{r=1}^n r$ to evaluate

$$\sum_{r=1}^{24} (r^3 - 4r)$$

(2)

- (ii) Use the standard results for $\sum_{r=1}^n r^2$ and $\sum_{r=1}^n r$ to show that

$$\sum_{r=0}^n (r^2 - 2r + 2n + 1) = \frac{1}{6}(n+1)(n+a)(bn+c)$$

for all integers $n \geq 0$, where a , b and c are constant integers to be found.

(6)

$$\begin{aligned} \text{a) } \sum_1^{24} r^3 - 4 \sum_1^{24} r &= \frac{1}{4}(24)^2(24+1)^2 - 4 \times \frac{1}{2}(24)(24+1) \\ &= 90000 - 1200 = \underline{\underline{88800}} \end{aligned}$$

$$\begin{aligned}
b) \quad \sum_0^n r^2 - 2r + 2n + 1 &= (2n+1) + \sum_1^n r^2 - 2r + 2n + 1 \\
&= (2n+1) + \sum r^2 - 2 \sum r + (2n+1) \sum 1 \\
&= (2n+1) + \frac{1}{6} n(n+1)(2n+1) - 2 \times \frac{1}{2} n(n+1) + (2n+1) \times n \\
&= \frac{1}{6} [n(n+1)(2n+1) + 6(2n+1) - 6n(n+1) + 6n(2n+1)] \\
&= \frac{1}{6} [(n+1)[n(2n+1) - 6n] + 6(2n+1)(n+1)] \\
&= \frac{1}{6} [(n+1)(n(2n+1) - 6n + 6(2n+1))] \\
&= \frac{1}{6} [(n+1)(2n^2 + n - 6n + 12n + 6)] \\
&= \frac{1}{6} [(n+1)(2n^2 + 7n + 6)] \\
&= \frac{1}{6} (n+1)(n+2)(2n+3)
\end{aligned}$$