



General Certificate of Education
Advanced Level Examination
June 2012

Mathematics

MFP2

Unit Further Pure 2

Thursday 31 May 2012 9.00 am to 10.30 am

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.
You may use a graphics calculator.

Time allowed

- 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

1 (a) Sketch the curve $y = \cosh x$. *(1 mark)*

(b) Solve the equation

$$6 \cosh^2 x - 7 \cosh x - 5 = 0$$

giving your answers in logarithmic form. *(6 marks)*



2 (a) Draw on the Argand diagram below:

(i) the locus of points for which

$$|z - 2 - 3i| = 2 \quad (3 \text{ marks})$$

(ii) the locus of points for which


$$|z + 2 - i| = |z - 2| \quad (3 \text{ marks})$$

(b) Indicate on your diagram the points satisfying both

$$|z - 2 - 3i| = 2$$

and

$$|z + 2 - i| \leq |z - 2| \quad (1 \text{ mark})$$



Turn over ►



3 (a) Show that

$$\frac{2^{r+1}}{r+2} - \frac{2^r}{r+1} = \frac{r2^r}{(r+1)(r+2)} \quad (3 \text{ marks})$$

(b) Hence find

$$\sum_{r=1}^{30} \frac{r2^r}{(r+1)(r+2)}$$

giving your answer in the form $2^n - 1$, where n is an integer. (3 marks)

4 The cubic equation

$$z^3 + pz + q = 0$$

has roots α , β and γ .

(a) (i) Write down the value of $\alpha + \beta + \gamma$. (1 mark)

(ii) Express $\alpha\beta\gamma$ in terms of q . (1 mark)

(b) Show that

$$\alpha^3 + \beta^3 + \gamma^3 = 3\alpha\beta\gamma \quad (3 \text{ marks})$$

(c) Given that $\alpha = 4 + 7i$ and that p and q are real, find the values of:

(i) β and γ ; (2 marks)

(ii) p and q . (3 marks)

(d) Find a cubic equation with integer coefficients which has roots $\frac{1}{\alpha}$, $\frac{1}{\beta}$ and $\frac{1}{\gamma}$. (3 marks)

5 The function f , where $f(x) = \sec x$, has domain $0 \leq x < \frac{\pi}{2}$ and has inverse function f^{-1} , where $f^{-1}(x) = \sec^{-1} x$.

(a) Show that

$$\sec^{-1} x = \cos^{-1} \frac{1}{x} \quad (2 \text{ marks})$$

(b) Hence show that

$$\frac{d}{dx}(\sec^{-1} x) = \frac{1}{\sqrt{x^4 - x^2}} \quad (4 \text{ marks})$$



6 (a) Show that

$$\frac{1}{4}(\cosh 4x + 2 \cosh 2x + 1) = \cosh^2 x \cosh 2x \quad (3 \text{ marks})$$

(b) Show that, if $y = \cosh^2 x$, then

$$1 + \left(\frac{dy}{dx}\right)^2 = \cosh^2 2x \quad (3 \text{ marks})$$

(c) The arc of the curve $y = \cosh^2 x$ between the points where $x = 0$ and $x = \ln 2$ is rotated through 2π radians about the x -axis. Show that the area S of the curved surface formed is given by

$$S = \frac{\pi}{256}(a \ln 2 + b)$$

where a and b are integers.

(7 marks)

7 (a) Prove by induction that, for all integers $n \geq 1$,

$$\frac{3}{1^2 \times 2^2} + \frac{5}{2^2 \times 3^2} + \frac{7}{3^2 \times 4^2} + \dots + \frac{2n+1}{n^2(n+1)^2} = 1 - \frac{1}{(n+1)^2} \quad (7 \text{ marks})$$

(b) Find the smallest integer n for which the sum of the series differs from 1 by less than 10^{-5} .

(2 marks)

Turn over ►



8 (a) Use De Moivre's Theorem to show that, if $z = \cos \theta + i \sin \theta$, then

$$z^n + \frac{1}{z^n} = 2 \cos n\theta \quad (3 \text{ marks})$$

(b) (i) Expand $\left(z^2 + \frac{1}{z^2}\right)^4$. (1 mark)

(ii) Show that

$$\cos^4 2\theta = A \cos 8\theta + B \cos 4\theta + C$$

where A , B and C are rational numbers. (4 marks)

(c) Hence solve the equation

$$8 \cos^4 2\theta = \cos 8\theta + 5$$

for $0 \leq \theta \leq \pi$, giving each solution in the form $k\pi$. (3 marks)

(d) Show that

$$\int_0^{\frac{\pi}{2}} \cos^4 2\theta \, d\theta = \frac{3\pi}{16} \quad (3 \text{ marks})$$

