

1. 2 is a ‘critical value’, e.g. used in solution, or  $x = 2$  seen as an asymptote

$$x^2 = 2x^2 - 4x \Rightarrow x^2 - 4x = 0$$

$$x = 0, \quad x = 4 \quad \text{M1: two other critical values}$$

$$x < 0 \quad \text{B1}$$

$$2 < x < 4 \quad \text{M1: An inequality using the critical value } 2 \quad \text{M1 A1} \quad 6$$

[6]

First M mark can be implied by the two correct values, but otherwise a Method must be seen.

≤ appearing: maximum 1 mark penalty (at first occurrence).

2. (a)  $m^2 + 2m + 5 = 0 \Rightarrow = -1 \pm 2i \quad \text{M1 A1}$

$$x = e^{-t} (A \cos 2t + B \sin 2t)$$

M: Correct form (needs the two different constants)

M1 A1 4

(b)  $(1, 0) \Rightarrow A = 1 \quad \text{dB1}$

$$\dot{x} = -e^{-t} (A \cos 2t + B \sin 2t) + e^{-t} (-2A \sin 2t + 2B \cos 2t)$$

M: Product diff. attempt  $\quad \text{dM1}$


$$\text{With } A = 1, e^{-t} \{ \cos 2t(-1 + 2B) + \sin 2t(-B - 2) \}$$

$$\dot{x} = 1, t = 0 \Rightarrow 1 = -A + 2B \quad \text{M1}$$

$$B = 1 \quad (x = e^{-t} (\cos 2t + \sin 2t))$$

M: Use value of  $A$  to find  $B$ .  $\quad \text{dM1 A1cso} \quad 5$

(c)



‘Single oscillation’ between 0 and  $\pi$   $\quad \text{B1}$

Decreasing amplitude (dep. on a turning point)  $\quad \text{B1ft}$

Initially increasing to maximum  $\quad \text{B1ft}$

Any one correct intercept, whether in terms of  $\pi$  or not: 1 or  $\frac{3\pi}{8}$  or  $\frac{7\pi}{8}$  B1 4

(Allow degrees:  $67.5^\circ$  or  $157.5^\circ$ ) (Allow awrt  $0.32\pi$  or  $1.18$  or  $2.75$ )

[13]

- (a) First M: Form and attempt to solve auxiliary equation.  
 $2^{\text{nd}} \text{ M: } Ae^{(-1+2i)t} + 5e^{(-1+2i)t}$  scores M1, as does  $Ae^{m_1 t} + Be^{m_2 t}$  for real  $m_1, m_2$ .

- (b) B mark and first and third M marks are dependent on the M's in part (a).

- (c) First B1: Starts on positive  $x$ -axis, dips below  $t$ -axis, above  $t$ -axis at  $t = \pi$ , and no more than 2 turning points between 0 and  $\pi$   
 (Assume 0 to  $\pi$  if axis is not labelled).

Second B1ft: Increasing amplitude for positive real part of  $m$ .

Third B 1ft: Initially decreasing to minimum for negative  $B$ .

Initially at maximum for  $B = 0$ .

Final B1: Dependent on a sketch attempt.

Confusion of variables: Can lose the final A mark in (a).

3. (a)  $\frac{dy}{dx} = v + x \frac{dv}{dx}$  B1

$$v + x \frac{dv}{dx} = \frac{3x - 4vx}{4x + 3vx} \quad (\text{all in terms of } v \text{ and } x) \quad \text{M1}$$

$$x \frac{dv}{dx} = \frac{3 - 4v - v(4 + 3v)}{4 + 3v}$$

(Requires  $x \frac{dv}{dx} = f(v)$ , 2 terms over common denom.) M1

$$x \frac{dv}{dx} = \frac{3v^2 + 8v - 3}{3v + 4} \quad \text{A1 cso} \quad 4$$

(b)  $\frac{3v + 4}{3v^2 + 8v - 3} dv = -\frac{1}{x} dx$  Separating variables M1

$$\pm \ln x \quad \text{B1}$$

$$\frac{1}{2} \ln(3v^2 + 8v - 3) \quad M: k \ln(3v^2 + 8v - 3) \quad \text{M1 A1}$$

$$\frac{1}{2} \ln \left( \frac{3y^2}{x^2} + \frac{8y}{x} - 3 \right) = -\ln x + C \quad \text{Or any equivalent form} \quad \text{A1} \quad 5$$

(c)  $\frac{3y^2}{x^2} + \frac{8y}{x} - 3 = \frac{A}{x^2}$

Removing ln's correctly at any stage, dep. on having C. M1

Using (1, 7) to form an equation in A (need not be  $A = \dots$ ) M1

$$(1,7) \Rightarrow 3 \times 49 + 56 - 3 = A \Rightarrow A = 200 \text{ (or equiv., can still be ln) A1}$$

$$3y^2 + 8yx - 3x^2 = 200$$

$$(3y - x)(y + 3x) = 200 \quad (\text{M dependent on the 2 previous M's}) \text{ M1 A1 cso} \quad 5$$

[14]

Parts (b) and (c) may well merge.

(b) Partial fractions may be used  $\left( A = \frac{3}{2}, B = \frac{1}{2} \right)$ , giving  $\frac{1}{2} \ln(3v - 1) + \frac{1}{2} \ln(v + 3)$ .

(c) Final M requires formation and factorisation of the quadratic.

4. (a) (i)  $r^2 \sin^2 \theta = a^2 \cos 2\theta \sin^2 \theta = a^2(1 - 2 \sin^2 \theta) \sin^2 \theta$  B1 1  
 $(= a^2(\sin^2 \theta - 2 \sin^4 \theta))$

(ii)  $\frac{d}{d\theta}(a^2(\sin^2 \theta - 2 \sin^4 \theta)) = a^2(2 \sin \theta \cos \theta - 8 \sin^2 \theta \cos \theta), = 0$

M1, A1, M1

$$2 = 8 \sin^2 \theta \quad (\text{Proceed to a } \sin^2 \theta = b) \quad \text{M1}$$

$$\sin \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}, r = \frac{a}{\sqrt{2}} \quad \text{A1, A1 cso} \quad 6$$

(b)  $\frac{a^2}{2} \int \cos 2\theta d\theta = \frac{a^2}{4} \sin 2\theta \quad \text{M: Attempt } \frac{1}{2} \int r^2 d\theta, \text{ to get } k \sin 2\theta \quad \text{M1 A1}$

$$[\dots]_{\pi/6}^{\pi/4} = \frac{a^2}{4} \left[ 1 - \frac{\sqrt{3}}{2} \right] \quad \text{M: Using correct limits} \quad \text{M1 A1}$$

$$\Delta = \frac{1}{2} \left( \frac{a}{\sqrt{2}} \cdot \frac{1}{2} \right) \times \left( \frac{a}{\sqrt{2}} - \frac{\sqrt{3}}{2} \right) = \frac{\sqrt{3}a^2}{16}$$

M: Full method for rectangle or triangle M1 A1

$$R = \frac{\sqrt{3}a^2}{16} - \frac{a^2}{4} \left[ 1 - \frac{\sqrt{3}}{2} \right] = \frac{a^2}{16} (3\sqrt{3} - 4)$$

M: Subtracting, either way round dM1 A1 cso 8

[15]

- (a) (ii) First A1: Correct derivative of a correct expression for  $r^2 \sin^2 \theta$  or  $r \sin \theta$ .  
 (b) Final M mark is dependent on the first and third M's.

Attempts at the triangle area by integration: a full method is required for M1.

Missing a factors: (or  $a^2$ ) Maximum one mark penalty in the question.

5.  $\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}$  B1

$\cos \frac{\pi}{10} + i \sin \frac{\pi}{10}$  B1

$\cos\left(\frac{(4k+1)\pi}{10}\right) + i \sin\left(\frac{(4k+1)\pi}{10}\right), k = 2, 3, 4 (\text{or equiv.})$  M1 A2, 1, 0 5

$\left[\cos\left(\frac{9\pi}{10}\right) + i \sin\left(\frac{9\pi}{10}\right), \cos\left(\frac{13\pi}{10}\right) + i \sin\left(\frac{13\pi}{10}\right), \cos\left(\frac{17\pi}{10}\right) + i \sin\left(\frac{17\pi}{10}\right)\right]$

[Degrees : 18, 90, 162, 234, 306]

[5]

6.  $\left(\frac{dy}{dx}\right)_0 \approx \frac{y_1 - y_{-1}}{2h} \Rightarrow 2 \approx \frac{y_1 - y_{-1}}{0.2} \Rightarrow y_1 - y_{-1} \approx 0.4$  M1 A1

$\left(\frac{d^2y}{dx^2}\right)_0 \approx \frac{y_1 - 2y_0 + y_{-1}}{2h} \Rightarrow 8 \approx \frac{y_1 - 2y_0 + y_{-1}}{0.01}$

[For M1, an attempt at evaluating  $\left(\frac{d^2y}{dx^2}\right)_0$  is required.]

$\Rightarrow y_1 + y_{-1} \approx 2.08$  A1

Subtracting to give  $y_{-1} \approx 0.84$  M1 A1 6

[6]

7. (a) Correct method for producing 2<sup>nd</sup> order differential equation M1

e.g.  $\frac{d}{dx} \left\{ (1+2x) \frac{dy}{dx} \right\} = \frac{d}{dx} \left\{ x + 4y^2 \right\}$  attempted

$(1+2x) \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} = 1 + 8y \frac{dy}{dx}$  seen + conclusion AG A1 2

(b) Differentiating again w.r.t.  $x$ :

$$(1 + 2x)\frac{d^3y}{dx^3} + 2\frac{d^2y}{dx^2} = 8y\frac{d^2y}{dx^2} + 8\left(\frac{dy}{dx}\right)^2 - 2\frac{d^2y}{dx^2} \text{ or equiv.} \quad \text{M1 A2, 1, 0} \quad 3$$

$$[\text{e.g. } (1 + 2x)\frac{d^3y}{dx^3} = 8\left(\frac{dy}{dx}\right)^2 + 4(2y - 1)\frac{d^2y}{dx^2}$$

$$(c) \frac{dy}{dx} \text{ (at } x = 0) = 1 \quad \text{B1}$$

$$\text{Finding } \frac{d^2y}{dx^2} \text{ (at } x = 0) \quad (= 3) \quad \text{M1}$$

$$\text{Finding } \frac{d^3y}{dx^3}, \text{ at } x = 0; \quad = 8 \text{ [A1 f.t. is on part (c) values only]} \quad \text{M1 A1ft}$$

$$y = \frac{1}{2} + x + \frac{3}{2}x^2 + \frac{4}{3}x^3 + \dots \quad \text{M1 A1} \quad 6$$

[11]

[Alternative (c):

$$\text{Polynomial for } y: y = \frac{1}{2} + ax + bx^2 + cx^3 + \dots \quad \text{M1}$$

In given d.e.:

$$(1 + 2x)(a + 2bx + 3cx^2 + \dots) \equiv x + 4(\frac{1}{2} + ax + bx^2 + cx^3 + \dots)^2 \quad \text{M1A1}$$

$$a = 1 \quad \text{B1,} \quad \text{Complete method for other coefficients M1, answer A1}$$

8. (a) Relating lines and angle (generous) M1

*[angle between  $\pm 2i$  to  $P$  and  $\pm 2$  to  $P$ ]* A1

$$\text{Angle between correct lines is } \frac{\pi}{2} \quad \text{M1 A1} \quad 4$$

Circle

Selecting correct ("top half") semi-circle.

[If algebraic approach:

Method for finding Cartesian equation M1

Correct equation, any form,  $\Rightarrow x(x + 2) + y(y - 2) = 0$  A1

Sketch: showing circle M1


Correct circle { centre  $(-1, 1)$  }, choosing only "top half" A1]

(b)  $|z + 1 - i|$  is radius;  $= \sqrt{2}$  M1 A1 2

$$(c) \quad z = \frac{2(1+i) - 2\omega}{\omega} \quad \left( = \frac{2(1+i)}{\omega} - 2 \right) \quad M1$$

$$\frac{z - 2i}{z + 2} = \frac{2(1+i) - 2(1+i)\omega}{2(1+i)} \quad (= -\omega) \quad M1 A1$$

$\text{Arg}(1 - \omega) = \frac{\pi}{2}$  is line segment, passing through (1,0) A1, A1



A1 6

[12]

$$\text{Alt (c): } u + iv = \frac{2 + 2i}{(x+2) + iy} = \frac{(2x + 2y + 4) + i(x + 2 - y)}{(x+2)^2 + y^2} \quad M1$$

$$x = -1 + \sqrt{2} \cos \theta, y = 1 + \sqrt{2} \sin \theta \quad M1$$

$$\Rightarrow w = \frac{(2\sqrt{2} \cos \theta + 2\sqrt{2} \sin \theta + 4) + i.....}{(2\sqrt{2} \cos \theta + 2\sqrt{2} \sin \theta + 4)} \{= 1 + i f(\theta)\} \quad A1,$$

$\Rightarrow$  part of line  $u = 1$ , show lower “half” of line A1, A1