

1. Given that $3x \sin 2x$ is a particular integral of the differential equation

$$\frac{d^2y}{dx^2} + 4y = k \cos 2x,$$

where k is a constant,

- (a) calculate the value of k ,

(4)

- (b) find the particular solution of the differential equation for which at $x = 0$, $y = 2$, and for which at $x = \frac{\pi}{4}$, $y = \frac{\pi}{2}$.

(4)(Total 8 marks)

$$y = 3x \sin 2x$$

$$y' = 3 \sin 2x + 6x \cos 2x$$

$$y'' = 6 \cos 2x + 6 \cos 2x + -12x \sin 2x$$

$$y'' = 12 \cos 2x - 12x \sin 2x$$

$$\begin{aligned} y'' &= -12x \sin 2x + 12 \cos 2x \\ + 4y &\quad \underline{12x \sin 2x} \\ u \cos 2x &\equiv 12 \cos 2x \therefore u = 12 \end{aligned}$$

$$y_{PI} = 3x \sin 2x \Rightarrow u = 12$$

$$\begin{aligned} y &= Ae^{mx} & y'' &= A m^2 e^{mx} \Rightarrow M = \pm 2i \Rightarrow y = A \cos 2x + B \sin 2x \\ y' &= Ame^{mx} & \cancel{+ 4y} &= Am^2 e^{mx} \\ y'' &= Am^2 e^{mx} & \cancel{O} &= \cancel{4e^{mx}} / (m^2 + 4) \quad \therefore y = A \cos 2x + (B + 3x) \sin 2x \end{aligned}$$

$$(0, 2) \Rightarrow 2 = A \quad y = 2 \cos 2x + (B + 3x) \sin 2x$$

$$y' = -4 \sin 2x + 3 \sin 2x + 2(B + 3x) \cos 2x \quad [\text{not required}]$$

$$(\frac{\pi}{4}, \frac{\pi}{2}) \Rightarrow \frac{\pi}{2} = (0) + (B + \frac{3\pi}{4})(1) \quad \therefore B = -\frac{\pi}{4}.$$

$$\therefore y = 2 \cos 2x + (3x - \frac{\pi}{4}) \sin 2x$$

2

2. Given that for all real values of r ,

$$(2r+1)^3 - (2r-1)^3 = Ar^2 + B,$$

where A and B are constants,

- (a) find the value of A and the value of B .

(2)

$$(b) \text{ Hence, or otherwise, prove that } \sum_{r=1}^n r^2 = \frac{1}{6} n(n+1)(2n+1).$$

(5)

$$(c) \text{ Calculate } \sum_{r=1}^{40} (3r-1)^2.$$

(3)(Total 10 marks)

$$(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

$$(2r+1)^3 = (2r)^3 + 3(2r)^2 + 3(2r) + 1$$

$$(2r+1)^3 = 8r^3 + 12r^2 + 6r + 1$$

$$(2r-1)^3 = 8r^3 - 12r^2 + 6r - 1$$

$$24r^2 + 2 \quad A=24 \quad B=2$$

$$\sum_{r=1}^n 24r^2 + 2 = 24 \sum_{r=1}^n r^2 + 2n$$

$$\sum_{r=1}^n (3^3 - 1^3) + (5^3 - 3^3) + (7^3 - 5^3) + \dots + ((2n+1)^3 - (2n-1)^3)$$

$$\Rightarrow 24 \sum_{r=1}^n r^2 + 2n = (2n+1)^3 - 1 = 8n^3 + 12n^2 + 6n + 1 - 1$$

$$\Rightarrow 24 \sum_{r=1}^n r^2 = 8n^3 + 12n^2 + 4n = 4n(2n^2 + 3n + 1)$$

$$\therefore \sum_{r=1}^n r^2 = \frac{1}{6} n(2n^2 + 3n + 1) \Rightarrow \sum_{r=1}^n r^2 = \frac{1}{6} n(2n+1)(n+1)$$

$$(c) \sum_{r=1}^{40} (3r-1)^2 = \sum_{r=1}^{40} 9r^2 - 6 \sum_{r=1}^{40} r + \sum_{r=1}^{40} 1 = \frac{3}{2}(40)(81)(40) - 6 \left(\frac{40(41)}{2} \right) + 40$$

$$= 194380$$

3. (a) Use algebra to find the exact solutions of the equation

$$|2x^2 + x - 6| = 6 - 3x.$$

(6)


- (b) On the same diagram, sketch the curve with equation $y = |2x^2 + x - 6|$ and the line with equation $y = 6 - 3x$.

(3)

- (c) Find the set of values of x for which

$$|2x^2 + x - 6| > 6 - 3x.$$

(3)(Total 12 marks)



$$2x^2 + x - 6 = 6 - 3x$$

$$2x^2 + 4x - 12 = 0$$

$$x^2 + 2x - 6 = 0$$

$$(x+1)^2 = 7$$

$$x = -1 \pm \sqrt{7}$$

$$a = -1 - \sqrt{7}$$

$$b = -1 + \sqrt{7}$$

$$2x^2 + x - 6 = 3x - 6$$

$$2x^2 - 2x = 0$$

$$2x(x-1) = 0$$

$$x = 0 \quad x = 1$$

$$\begin{aligned} c) \quad & x < -1 - \sqrt{7} \\ & \text{or } x > -1 + \sqrt{7} \\ & \text{or } 0 < x < 1 \end{aligned}$$

4. During an industrial process, the mass of salt, S kg, dissolved in a liquid t minutes after the process begins is modelled by the differential equation

$$\frac{dS}{dt} + \frac{2S}{120-t} = \frac{1}{4}, \quad 0 \leq t < 120.$$

Given that $S = 6$ when $t = 0$,

- (a) find S in terms of t ,

(8)

- (b) calculate the maximum mass of salt that the model predicts will be dissolved in the liquid at any one time during the process.

(4)(Total 12 marks)

$$\frac{1}{(120-t)^2} \frac{dS}{dt} + \frac{2S}{(120-t)^3} = \frac{1}{4(120-t)^2} \Rightarrow \frac{d}{dt} \left[\frac{S}{(120-t)^2} \right] = \frac{1}{4(120-t)^2}$$

$$\Rightarrow \frac{S}{(120-t)^2} = \frac{1}{4} \int (120-t)^{-2} dt = \frac{1}{4} (120-t)^{-1} + C$$

$$\therefore S = \frac{1}{4}(120-t) + C(120-t)^2$$

$$S=6, t=0 \quad 6 = 30 + C(120)^2 \Rightarrow C = -\frac{1}{600}$$

$$\therefore S = 30 - \frac{1}{4}t - \frac{(120-t)^2}{600}$$

$$b) \text{ Max when } \frac{dS}{dt} = 0 \Rightarrow -\frac{1}{4} - \frac{(120-t)}{300} = 0 \Rightarrow 120-t = 75 \Rightarrow t = 45$$

$$\therefore \text{Max } S = \frac{1}{4}(120-45) + \frac{(120-45)^2}{600}$$

$$\text{Max } S = 9.38$$

$$|(2x-3)(x+2)|$$

1.5 -2

5. (a) Find the Taylor expansion of $\cos 2x$ in ascending powers of $\left(x - \frac{\pi}{4}\right)$ up to and including

the term in $\left(x - \frac{\pi}{4}\right)^5$.

(5)

- (b) Use your answer to (a) to obtain an estimate of $\cos 2$, giving your answer to 6 decimal places.

(3)(Total 8 marks)

$$\begin{aligned}
 y &= \cos 2x & f\left(\frac{\pi}{4}\right) &= 0 & \therefore f(x) &= -2\left(x - \frac{\pi}{4}\right) + \frac{4}{3}\left(x - \frac{\pi}{4}\right)^3 - \frac{4}{15}\left(x - \frac{\pi}{4}\right)^5 \\
 y' &= -2\sin 2x & f'\left(\frac{\pi}{4}\right) &= -2 & f(1) &\approx \cos 2 \\
 y'' &= -4\cos 2x & f''\left(\frac{\pi}{4}\right) &= 0 & f(1) &\approx -0.416147 \\
 y''' &= 8\sin 2x & f'''\left(\frac{\pi}{4}\right) &= 8 & \\
 y'''' &= 16\cos 2x & f''''\left(\frac{\pi}{4}\right) &= 0 & \\
 y''''' &= -32\sin 2x & f'''''\left(\frac{\pi}{4}\right) &= -32 &
 \end{aligned}$$

6. (a) Use de Moivre's theorem to show that $\sin 5\theta = \sin \theta (16\cos^4 \theta - 12\cos^2 \theta + 1)$.

(5)

- (b) Hence, or otherwise, solve, for $0 \leq \theta < \pi$

$$\sin 5\theta + \cos \theta \sin 2\theta = 0.$$

(6)(Total 11 marks)

$$\begin{array}{ccccccc}
 & & & 1 & 2 & 1 & \\
 & & & 1 & 3 & 2 & 1 \\
 & & & 1 & 4 & 6 & 4 & 1 \\
 & & & 1 & 5 & 10 & 10 & 5 & 1
 \end{array}$$

$$(\cos \theta + i \sin \theta)^5 = \cos 5\theta + i \sin 5\theta$$

$$= \cos^5 \theta + 5i \cos^4 \theta \sin \theta - 10 \cos^3 \theta \sin^2 \theta - 10i \cos^2 \theta \sin^3 \theta + 5 \cos \theta \sin^4 \theta + i \sin^5 \theta$$

$$\text{equating imaginary parts} \Rightarrow \sin 5\theta = 5\cos^4 \theta \sin \theta - 10 \cos^2 \theta \sin^3 \theta + \sin^5 \theta$$

$$\Rightarrow \sin 5\theta = \sin \theta [5\cos^4 \theta - 10 \cos^2 \theta (1 - \cos^2 \theta) + (1 - 2\cos^2 \theta + \cos^4 \theta)]$$

$$\Rightarrow \sin 5\theta = \sin \theta [5\cos^4 \theta - 10 \cos^2 \theta + 10 \cos^4 \theta + 1 - 2\cos^2 \theta + (\cos^4 \theta)]$$

$$\therefore \sin 5\theta = \sin \theta [16\cos^4 \theta - 12\cos^2 \theta + 1] \quad *$$

$$b) \sin 5\theta + \cos \theta \sin 2\theta = 0$$

$$= \sin \theta [16\cos^4 \theta - 12\cos^2 \theta + 1] + 2\sin \theta \cos^2 \theta = 0$$

$$= \sin \theta [16\cos^4 \theta - 10\cos^2 \theta + 1] = 0$$

$$= \sin \theta (8\cos^2 \theta - 1)(2\cos^2 \theta - 1) = 0$$

$$\sin \theta = 0 \\ \therefore \theta = 0$$

$$\cos \theta = \sqrt{\frac{1}{8}}$$

$$\theta = 1.209^\circ$$

$$\cos \theta = -\sqrt{\frac{1}{8}}$$

$$\theta = 1.932^\circ$$

$$\cos \theta = \frac{1}{\sqrt{2}}$$

$$\theta = \frac{\pi}{4}$$

$$\cos \theta = -\frac{1}{\sqrt{2}}$$

$$\theta = \frac{3}{4}\pi$$

7.

$$\frac{d^2x}{dt^2} + 3\sin x = 0 \quad \text{At } t=0, \quad x=0 \quad \text{and} \quad \frac{dx}{dt}=0.4$$

$$t=0$$

$$x=0$$

$$x'=0.4$$

- (b) Find a series solution for x , in ascending powers of t , up to and including the term in t^3 .

(4)

- (c) Use your answer to (b) to obtain an estimate of x at $t=0.3$.

(2)(Total 11 marks)

$$\frac{d^3x}{dt^3} + \frac{d}{dt}(3\sin x) = 0 \Rightarrow \frac{d^3x}{dt^3} + (3\cos x)\frac{dx}{dt} = 0$$

$$x'' + 3\sin x = 0 \Rightarrow x'' + 3(0) \Rightarrow x'' = 0$$

$$x''' + (3\cos(0))0 \cdot 4 = 0 \Rightarrow x''' = -1.2$$

$$x = 0.4t - 0.2t^3 \quad t=0.3 \quad x \approx 0.1146$$

8. The point P represents a complex number z on an Argand diagram, where

$$|z - 6 + 3i| = 3|z + 2 - i|$$

- (a) Show that the locus of P is a circle, giving the coordinates of the centre and the radius of this circle.

(7)

- The point Q represents a complex number z on an Argand diagram, where

$$\tan[\arg(z+6)] = \frac{1}{2}$$

- (b) On the same Argand diagram, sketch the locus of P and the locus of Q .

(5)

- (c) On your diagram, shade the region which satisfies both

$$|z - 6 + 3i| > 3|z + 2 - i| \quad \text{and} \quad \tan[\arg(z+6)] > \frac{1}{2}$$

(2)(Total 14 marks)

$$a) |(x-6)+i(y+3)| = 3|(x+2)+i(y-1)|$$

$$\Rightarrow (x-6)^2 + (y+3)^2 = 9(x+2)^2 + 9(y-1)^2$$

$$\Rightarrow x^2 - 12x + 36 + y^2 + 6y + 9 = 9x^2 + 36x + 36 + 9y^2 - 18y + 9$$

$$\Rightarrow 8x^2 + 48x + 8y^2 - 24y = 0 \quad (\div 8) \quad x^2 + 6x + y^2 - 3y = 0$$

$$(x+3)^2 + (y-2)^2 = 13$$

$$\text{radius } (-3, 2) \quad r = \sqrt{13}$$

$$b) \arg(z+6) = \tan^{-1}\left(\frac{1}{2}\right) \Rightarrow$$

$$x=0 \quad 9+(y-2)^2=13 \quad y-2=\pm 2 \\ \therefore y=0 \quad y=4$$

passes through origin

$$y=0 \quad (x+3)^2=9 \\ x=-3 \pm 3$$

$$\arg(z+6) > \tan^{-1}\left(\frac{1}{2}\right) \text{ i.e. above angle}$$

