

1. Obtain the general solution of the differential equation

$$x \frac{dy}{dx} + 2y = \cos x, \quad x > 0,$$

giving your answer in the form $y = f(x)$.

(Total 8 marks)

$$\frac{dy}{dx} + \frac{2}{x}y = \frac{\cos x}{x}$$


$$e^{\int \frac{2}{x} dx} \Rightarrow (e^{\ln x})^2 = x^2$$

$$x^2 \frac{dy}{dx} + 2xy = x \cos x \Rightarrow \frac{d}{dx}(x^2 y) = x \cos x \Rightarrow x^2 y = \int x \cos x dx$$

$$\begin{aligned} \downarrow \quad u=x \quad v=\sin x \quad \uparrow \\ u'=1 \quad v'=\cos x \end{aligned} \Rightarrow x^2 y = x \sin x - \int \sin x dx$$

$$\Rightarrow x^2 y = x \sin x + \cos x + c \quad \therefore y = \frac{\sin x}{x} + \frac{\cos x}{x^2} + \frac{c}{x^2}$$

2.



The diagram above shows a sketch of the curve with equation

$$y = \frac{x^2 - 1}{x + 2}, \quad x \neq -2.$$

The curve crosses the x-axis at $x = 1$ and $x = -1$ and the line $x = -2$ is an asymptote of the curve.

(a) Use algebra to solve the equation $\frac{x^2 - 1}{x + 2} = 3(1 - x)$.

(b) Hence, or otherwise, find the set of values of x for which

$$\frac{x^2 - 1}{x + 2} < 3(1 - x).$$

(Total 9 marks)

$$a) (x^2 - 1) = 3[(x + 2)(1 - x)]$$

$$\Rightarrow x^2 - 1 = 3(x + 2)(1 - x)$$

$$\Rightarrow x^2 - 1 = -3x^2 - 3x + 6$$

$$\Rightarrow 4x^2 + 3x - 7 = 0$$

$$\Rightarrow (4x + 7)(x - 1) = 0$$

$$x = -\frac{7}{4} \quad x = 1$$

$$\Rightarrow x^2 - 1 = -3(x + 2)(1 - x)$$

$$x^2 - 1 = +3x^2 + 3x - 6$$

$$0 = 2x^2 + 3x - 5$$

$$0 = (2x + 5)(x - 1)$$

$$x = -\frac{5}{2} \quad x = 1$$

$$x < -\frac{5}{2} \quad \text{or} \quad -\frac{7}{4} < x < 1$$

3. A scientist is modelling the amount of a chemical in the human bloodstream. The amount x of the chemical, measured in mg l^{-1} , at time t hours satisfies the differential equation

$$2x \frac{d^2x}{dt^2} - 6 \left(\frac{dx}{dt} \right)^2 = x^2 - 3x^4, \quad x > 0.$$

- (a) Show that the substitution $y = \frac{1}{x^2}$ transforms this differential equation into

$$\frac{d^2y}{dt^2} + y = 3. \quad [1]$$

- (b) Find the general solution of differential equation [1].

Given that at time $t = 0$, $x = \frac{1}{2}$ and $\frac{dx}{dt} = 0$,

- (c) find an expression for x in terms of t ,

- (d) write down the maximum value of x as t varies.

(1)
(Total 14 marks)

$$\frac{d^2x}{dt^2} = \frac{d}{dt} \left(-\frac{x^3}{2} \frac{dy}{dt} \right) = -\frac{x^3}{2} \frac{d^2y}{dt^2} - \frac{3x^2}{2} \frac{dx}{dt} \frac{dy}{dt}$$

$$\therefore \frac{d^2x}{dt^2} = -\frac{x^3}{2} \frac{d^2y}{dt^2} + \frac{3x^5}{4} \left(\frac{dy}{dt} \right)^2$$

$$2x \frac{d^2x}{dt^2} - 6 \left(\frac{dx}{dt} \right)^2 = x^2 - 3x^4$$

$$2x \left(-\frac{x^3}{2} \frac{d^2y}{dt^2} + \frac{3x^5}{4} \left(\frac{dy}{dt} \right)^2 \right) - 6 \left(-\frac{x^3}{2} \frac{dy}{dt} \right)^2 = x^2 - 3x^4$$

$$-x^4 \frac{d^2y}{dt^2} + \frac{3x^6}{2} \left(\frac{dy}{dt} \right)^2 - \frac{3x^6}{2} \left(\frac{dy}{dt} \right)^2 = x^2 - 3x^4$$

$$\div -x^2 \quad x^2 \frac{d^2y}{dt^2} = 3x^2 - 1 \Rightarrow \frac{1}{y} \frac{d^2y}{dt^2} = \frac{3}{y} - 1$$

$$(xy) \quad \frac{dy}{dt} = 3 - y \quad \therefore \frac{dy}{dt} + y = 3$$

hmm!

$$\text{alt } y = x^{-2} \quad \frac{dy}{dt} = -2x^{-3} \frac{dx}{dt} \Rightarrow \frac{dx}{dt} = -\frac{x^3}{2} \frac{dy}{dt}$$

$$\frac{d^2x}{dt^2} = -\frac{3x^2}{2} \frac{dx}{dt} \frac{dy}{dt} - \frac{x^3}{2} \frac{d^2y}{dt^2} \Rightarrow \frac{d^2x}{dt^2} = \frac{3x^5}{4} \left(\frac{dy}{dt} \right)^2 - \frac{x^3}{2} \frac{d^2y}{dt^2}$$

then sub in as previously.

$$x^2 y = 1$$

$$\frac{d}{dt}(x^2 y) = 0$$

$$\left[\frac{d}{dt}(x^2) \right] y + x^2 \left(\frac{d}{dt} y \right) = 0$$

$$(5) \quad 2x \frac{dx}{dt} y + x^2 \frac{dy}{dt} = 0$$

$$(4) \quad 2xy \frac{dx}{dt} + x^2 \frac{dy}{dt} = 0$$

$$(4) \quad \Rightarrow \frac{dx}{dt} = -\frac{x}{2y} \frac{dy}{dt}$$

$$\Rightarrow \frac{dx}{dt} = -\frac{x^3}{2} \frac{dy}{dt}$$

$$\frac{d^2y}{dt^2} + y = 3$$

$$y = Ae^{mt}$$

$$y' = Ame^{mt}$$

$$y'' = Am^2e^{mt}$$

$$y'' + y = 0$$

$$Ae^{mt}(m^2 + 1) = 0$$

$$\neq 0 \quad = 0 \Rightarrow m = \pm i$$

$$\therefore y_{cf} = A\cos t + B\sin t$$

$$y = \lambda$$

$$y' = 0$$

$$y'' = 0$$

$$y'' + y = 3$$

$$\Rightarrow \lambda = 3$$

$$\therefore y_{PI} = 3$$

$$\therefore y = A\cos t + B\sin t + 3$$

$$c) \quad \frac{1}{x^2} = A\cos t + B\sin t + 3$$

$$\Rightarrow x = \sqrt{\frac{1}{A\cos t + B\sin t + 3}}$$

$$x = \frac{1}{2}, t = 0 \quad \frac{1}{2} = \sqrt{\frac{1}{A+3}} \quad \therefore A = 1$$

$$\therefore y = \cos t + B\sin t + 3$$

$$\frac{dy}{dt} = -\sin t + B\cos t \Rightarrow \frac{-2}{x^3} \frac{dx}{dt} = -\sin t + B\cos t$$

$$x = \frac{1}{2}, t = 0, \frac{dx}{dt} = 0 \Rightarrow -16 \times 0 = B \quad \therefore B = 0$$

$$\therefore x = \sqrt{\frac{1}{\cos t + 3}}$$


$$\max x \Rightarrow \frac{dx}{dt} = 0 \Rightarrow \frac{dy}{dt} = 0 \Rightarrow \sin t = 0$$

$$\Rightarrow t = 0, \pi, \dots$$

$$\therefore \max \text{ when } \cos t = -1 \text{ (at } t = \pi)$$

$$\Rightarrow x = \sqrt{\frac{1}{2}} \quad \therefore \max x = \frac{\sqrt{2}}{2}$$

4.



The diagram above shows a sketch of the curve C with polar equation

$$r = 4\sin\theta\cos^2\theta, \quad 0 \leq \theta < \frac{\pi}{2}.$$

The tangent to C at the point P is perpendicular to the initial line.

- (a) Show that P has polar coordinates $\left(\frac{3}{2}, \frac{\pi}{6}\right)$.

The point Q on C has polar coordinates $\left(\sqrt{2}, \frac{\pi}{4}\right)$.

The shaded region R is bounded by OP , OQ and C , as shown in the diagram above.

- (b) Show that the area of R is given by

$$\int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \left(\sin^2 2\theta \cos 2\theta + \frac{1}{2} - \frac{1}{2} \cos 4\theta \right) d\theta$$

- (c) Hence, or otherwise, find the area of R , giving your answer in the form $a + b\pi$, where a and b are rational numbers.

tangent perp to initial line $\Rightarrow \frac{dx}{d\theta} = 0$

$$x = r \cos \theta \quad x = 4\sin\theta \cos^3\theta$$

$$\frac{dx}{d\theta} = 4\cos^4\theta - 12\sin^2\theta \cos^2\theta$$

$$12(1 - \cos^2\theta)\cos^2\theta = 4\cos^4\theta$$

$$12\cos^2\theta = 16\cos^4\theta$$

$$4\cos^2\theta(4\cos^2\theta - 3) = 0$$

$$\cos\theta = 0 \quad \cos\theta = \pm \frac{\sqrt{3}}{2}$$

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2} \quad \theta = \frac{\pi}{6}, \frac{5\pi}{6}$$

$$r = 4\sin\theta \cos^2\theta \quad \therefore \cos\theta = \frac{\sqrt{3}}{2}$$

$$r = 4\left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right)^2 \quad \therefore \sin\theta = \frac{1}{2}$$

$$r = \frac{3}{2} \quad \therefore P\left(\frac{3}{2}, \frac{\pi}{6}\right)$$

(5)
(Total 14 marks)

$$R = \frac{1}{2} \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} 16\sin^2\theta \cos^4\theta d\theta = 2 \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} (2\sin\theta \cos\theta)^2 \left(\frac{1}{2}(\cos 2\theta + 1)\right) d\theta$$

$$R = \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \sin^2 2\theta (\cos 2\theta + 1) d\theta = \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \sin^2 2\theta \cos 2\theta + \sin^2 2\theta d\theta$$

$$R = \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \sin^2 2\theta \cos 2\theta + \frac{1}{2} - \frac{1}{2} \cos 4\theta d\theta$$

$$\therefore R = \left[\frac{1}{6} \sin^3 2\theta + \frac{1}{2} \theta - \frac{1}{8} \sin 4\theta \right]_{\frac{\pi}{6}}^{\frac{\pi}{4}} = \left[\left(\frac{1}{6} + \frac{\pi}{8} \right) - \left(\frac{1}{6} \left(\frac{\sqrt{3}}{2} \right)^3 + \frac{\pi}{12} - \frac{\sqrt{3}}{16} \right) \right] = \frac{1}{6} + \frac{\pi}{24}$$


5. Find the set of values of x for which

$$\frac{x+1}{2x-3} < \frac{1}{x-3}$$

(Total 7 marks)

$$\begin{aligned} (x+1)(2x-3)(x-3)^2 &< (x-3)(2x-3)^2 \\ \Rightarrow (x+1)(2x-3)(x-3)^2 - (x-3)(2x-3)^2 &< 0 \\ \Rightarrow (x-3)(2x-3)[(x+1)(x-3) - (2x-3)] &< 0 \\ \Rightarrow (x-3)(2x-3)[x^2 - 2x - 3 - 2x + 6] &< 0 \\ \Rightarrow (x-3)(2x-3)[x^2 - 4x + 3] &< 0 \\ \Rightarrow (x-3)^2(2x-3)(x-1) &< 0 \end{aligned}$$

3, 3 1.5 1



6.

$$\frac{dy}{dx} - y \tan x = 2 \sec^3 x.$$

Given that $y = 3$ at $x = 0$, find y in terms of x

(Total 7 marks)

$$\text{IF } f(x) = e^{-\int \tan x dx} = e^{-\int \frac{\sin x}{\cos x} dx} = e^{\ln \cos x} = \cos x$$

$$\cos x \frac{dy}{dx} - y \cos x \tan x = 2 \sec^2 x \Rightarrow \frac{d}{dx}(y \cos x) = 2 \sec^2 x$$

$$\Rightarrow y \cos x = 2 \int \sec^2 x dx = 2 \tan x + C \quad \therefore y = \frac{2 \tan x + C}{\cos x}$$

$$(0, 3) \quad 3 = \frac{2+C}{1} \quad \therefore C = 1 \quad \Rightarrow y = \frac{2 \tan x + 1}{\cos x}$$

7. For the differential equation

$$\frac{d^2 y}{dx^2} + 3 \frac{dy}{dx} + 2y = 2x(x+3),$$

find the solution for which at $x = 0$, $\frac{dy}{dx} = 1$ and $y = 1$.

(Total 12 marks)

$$\begin{aligned} y &= Ae^{mx} \\ y' &= Am e^{mx} \\ y'' &= Am^2 e^{mx} \end{aligned}$$

$$\begin{aligned} y'' + 3y' + 2y &= 0 \\ Ae^{mx}(m^2 + 3m + 2) &= 0 \\ \neq 0 \quad \Rightarrow (m+2)(m+1) &= 0 \\ \quad \quad \quad -2 \quad \quad \quad -1 \end{aligned}$$

$$y_{cf} = Ae^{-x} + Be^{-2x}$$

$$\begin{aligned} y &= a + bx + cx^2 \\ y' &= b + 2cx \\ y'' &= 2c \end{aligned}$$

$$\begin{aligned} y'' &= 2c \\ + 3y' &= 3b + 6cx \\ + 2y &= 2a + 2bx + 2cx^2 \end{aligned}$$

$$2x^2 + 6x = (2a + 3b + 2c) + (2b + 6c)x + 2cx^2$$

$$\therefore \underline{c=1} \quad 2b+6=6 \quad \therefore \underline{b=0} \quad 2a+2=0 \quad \therefore \underline{a=-1}$$

$$y_{PI} = x^2 - 1$$

$$y' = -Ae^{-x} - 2Be^{-2x} + 2x$$

$$\begin{aligned} x=0, y'=1 \quad 1 &= -A - 2B \quad \Rightarrow \begin{aligned} A+2B &= -1 \\ A+B &= 2 \end{aligned} \\ \underline{B=-3} \quad \therefore \underline{A=5} \end{aligned}$$

$$\begin{aligned} \therefore y &= Ae^{-x} + Be^{-2x} + x^2 - 1 \\ x=0, y=1 \quad 1 &= A + B - 1 \quad \therefore A+B=2 \end{aligned}$$

$$\therefore y = 5e^{-x} - 3e^{-2x} + x^2 - 1$$

8. (a) Sketch the curve C with polar equation

$$r = 5 + \sqrt{3} \cos \theta,$$

$$0 \leq \theta \leq 2\pi.$$

(2)

(b) Find the polar coordinates of the points where the tangents to C are parallel to the initial line $\theta = 0$. Give your answers to 3 significant figures where appropriate.

(6)

(c) Using integration, find the area enclosed by the curve C , giving your answer in terms of π .

(6)
(Total 14 marks)




$$\theta = 0 \quad r_{\max} = 5 + \sqrt{3}$$

$$\theta = \pi \quad r_{\min} = 5 - \sqrt{3}$$

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2} \quad r = 5$$

$$\text{Parallel to initial line } \frac{dy}{d\theta} = 0$$



$$y = r \sin \theta = 5 \sin \theta + \sqrt{3} \sin \theta \cos \theta$$

$$\frac{dy}{d\theta} = 5 \cos \theta + \sqrt{3} (\cos^2 \theta - \sin^2 \theta)$$

$$5 \cos \theta + \sqrt{3} (\cos^2 \theta - \sin^2 \theta) = 0$$

$$2\sqrt{3} \cos^2 \theta + 5 \cos \theta - \sqrt{3} = 0$$

$$(2\sqrt{3} \cos \theta - 1)(\cos \theta + \sqrt{3}) = 0$$

$$\therefore \cos \theta = \frac{1}{2\sqrt{3}} \quad \cos \theta = -\sqrt{3} \quad \text{no solution}$$

$$\theta = \underline{1.28^\circ}, \underline{5.01^\circ}$$

$$\cos \theta = \frac{1}{2\sqrt{3}} \Rightarrow r = 5 + \frac{\sqrt{3}}{2\sqrt{3}} \therefore r = 5.5 \quad (5.5, 1.28); (5.5, 5.01)$$

$$\text{Area} = 2 \times \frac{1}{2} \int_0^\pi (5 + \sqrt{3} \cos \theta)^2 d\theta = \int_0^\pi 25 + 10\sqrt{3} \cos \theta + 3 \cos^2 \theta d\theta$$

$$= \int_0^\pi 25 + 10\sqrt{3} \cos \theta + 3 \left(\frac{1}{2} \cos 2\theta + \frac{1}{2} \right) d\theta = \int_0^\pi \frac{53}{2} + 10\sqrt{3} \cos \theta + \frac{3}{2} \cos 2\theta d\theta$$

$$= \frac{1}{2} \int_0^\pi 53 + 20\sqrt{3} \cos \theta + 3 \cos 2\theta d\theta$$

$$= \frac{1}{2} [53\theta + 20\sqrt{3} \sin \theta + \frac{3}{2} \sin 2\theta]_0^\pi = \frac{1}{2} [53\pi] = \frac{53}{2} \pi$$

10.

$$(1-x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} + 2y = 0.$$

At $x=0$, $y=2$ and $\frac{dy}{dx} = -1$.

(a) Find the value of $\frac{d^3y}{dx^3}$ at $x=0$.

(3)

(b) Express y as a series in ascending powers of x , up to and including the term in x^3 .

(4)

(Total 7 marks)

$$\frac{d}{dx}\left[(1-x^2)\frac{d^2y}{dx^2}\right] - \frac{d}{dx}\left(x\frac{dy}{dx}\right) + \frac{d}{dx}(2y) = 0 \quad (1-2x)\frac{d^2y}{dx^2} - 2x\frac{d^2y}{dx^2} - x\frac{d^2y}{dx^2} - \frac{dy}{dx} + 2\frac{dy}{dx} = 0$$

$$\Rightarrow (1-2x)\frac{d^2y}{dx^2} - 3x\frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$$

$$(1)(y'') - 0 + 2(2) = 0 \therefore y'' = -4$$

$$(1)y''' - 0 - 1 = 0 \therefore y''' = 1$$

$$\therefore y = 2 - x - 2x^2 + \frac{1}{6}x^3$$

11. (a) Given that $z = \cos \theta + i \sin \theta$, use de Moivre's theorem to show that

$$z^n + \frac{1}{z^n} = 2 \cos n\theta.$$

(2)

(b) Express $32 \cos^6 \theta$ in the form $p \cos 6\theta + q \cos 4\theta + r \cos 2\theta + s$, where p , q , r and s are integers.

(5)

(c) Hence find the exact value of

$$\int_0^{\frac{\pi}{3}} \cos^6 \theta d\theta.$$

(4)

(Total 11 marks)

$$z^6 + \frac{1}{z^6} = 2 \cos 6\theta$$

$$\left(z + \frac{1}{z}\right)^6 = (2 \cos \theta)^6 = 32 \cos^6 \theta$$

$$\left(z + \frac{1}{z}\right)^6 = z^6 + 6z^5\left(\frac{1}{z}\right) + 15z^4\left(\frac{1}{z^2}\right) + 20z^3\left(\frac{1}{z^3}\right) + 15z^2\left(\frac{1}{z^4}\right) + 6z\left(\frac{1}{z^5}\right) + \left(\frac{1}{z^6}\right)$$

$$\left(z + \frac{1}{z}\right)^6 = \left(z^6 + \frac{1}{z^6}\right) + 6\left(z^4 + \frac{1}{z^4}\right) + 15\left(z^2 + \frac{1}{z^2}\right) + 20$$

$$\therefore 32 \cos^6 \theta = 2 \cos 6\theta + 12 \cos 4\theta + 30 \cos 2\theta + 20$$

$$\therefore \int_0^{\frac{\pi}{3}} \cos^6 \theta d\theta = \frac{1}{16} \int_0^{\frac{\pi}{3}} \cos 6\theta + 6 \cos 4\theta + 15 \cos 2\theta + 10 d\theta$$

$$= \frac{1}{16} \left[\frac{1}{6} \sin 6\theta + \frac{3}{2} \sin 4\theta + \frac{15}{2} \sin 2\theta + 10\theta \right]_0^{\frac{\pi}{3}}$$

$$= \frac{1}{16} \left(-\frac{3\sqrt{3}}{4} + \frac{15\sqrt{3}}{4} + \frac{10\pi}{3} \right) = \frac{3\sqrt{3}}{16} + \frac{5\pi}{24}$$

$$z^n = \cos n\theta + i \sin n\theta$$

$$z^{-n} = \cos(-n\theta) + i \sin(-n\theta)$$

$$z^{-n} = \cos(n\theta) - i \sin(n\theta)$$

$$\therefore z^n + \frac{1}{z^n} = \frac{\cos n\theta + i \sin n\theta + \cos n\theta - i \sin n\theta}{2 \cos n\theta}$$

$$\begin{array}{ccccccc} & & 1 & & & & \\ & & 2 & & 1 & & \\ & 1 & 3 & 1 & & & \\ & 4 & 6 & 4 & 1 & & \\ 1 & 5 & 10 & 10 & 5 & 1 & \\ 6 & 15 & 20 & & & & \end{array}$$

12. The transformation T from the z -plane, where $z = x + iy$, to the w -plane, where

$w = u + iv$, is given by

$$w = \frac{z+1}{z}, \quad z \neq 0.$$

- (a) The transformation T maps the points on the line with equation $y = x$ in the z -plane, other than $(0, 0)$, to points on a line l in the w -plane. Find a cartesian equation of l .

(5)

- (b) Show that the image, under T , of the line with equation $x + y + 1 = 0$ in the z -plane is a circle C in the w -plane, where C has cartesian equation

$$u^2 + v^2 - u + v = 0.$$

(7)

- (c) On the same Argand diagram, sketch l and C .

(3)

(Total 15 marks)

$$wz = z + i \Rightarrow wz - z = i \Rightarrow z(w - 1) = i \Rightarrow z = \frac{i}{w - 1}$$

$$z = \frac{i}{(u-1)+iv} \cdot \frac{(u-1)-iv}{(u-1)-iv} = \frac{v + i(u-1)}{(u-1)^2 + v^2}$$

$$y = x \Rightarrow \frac{v}{(u-1)^2 + v^2} = \frac{u-1}{(u-1)^2 + v^2} \Rightarrow v = u - 1$$

$$b) \quad y = x - 1 \Rightarrow \frac{u-1}{(u-1)^2 + v^2} = \frac{v}{(u-1)^2 + v^2} - \frac{1}{(u-1)^2 + v^2}$$

$$\Rightarrow (u-1) = -v - (u-1)^2 - v^2$$

$$\Rightarrow u-1 = -v - u^2 + 2u - 1 - v^2$$

$$\Rightarrow u^2 - u + v^2 + v = 0 \quad \#$$

$$\Rightarrow \left(u - \frac{1}{2}\right)^2 + \left(v + \frac{1}{2}\right)^2 = \frac{1}{2}$$

$$u=0, v=0$$

$$C\left(\frac{1}{2}, -\frac{1}{2}\right) \quad r = \frac{\sqrt{2}}{2}$$

$$u=0 \quad \left(v + \frac{1}{2}\right)^2 = \frac{1}{4}$$

$$v + \frac{1}{2} = \pm \frac{1}{2} \Rightarrow v=0, v=1$$

