

1. Attempt to arrange in correct form $\frac{dy}{dx} + \frac{2}{x}y = \frac{\cos x}{x}$ M1

Integrating Factor: $= e^{\int \frac{2}{x} dx}, \left[(= e^{2 \ln x} = e^{\ln x^2}) = x^2 \right]$ M1, A1

$[x^2 \frac{dy}{dx} + 2xy = x \cos x \text{ implies M1M1A1}]$

$\therefore x^2 y = \int x^2 \cdot \frac{\cos x}{x} dx \text{ or equiv.}$ M1ft

[IF. $y = \int I.F. (\text{candidate's RHS}) dx$]

By Parts: $(x^2 y) = x \sin x - \int \sin x dx$ M1

i.e. $(x^2 y) = x \sin x, + \cos x (+ c)$ A1, A1cao

$y = \frac{\sin x}{x} + \frac{\cos x}{x^2} + \frac{c}{x^2}$ A1ft8

First M: At least two terms divided by x .

“By parts” M: Must be complete method, e.g $\int x^2 \cos x dx$ requires **two** applications

Because of functions involved, **be generous with sign**, but $x \sin x \pm \int \cos x dx$ is M0

(S.C. “Loop” integral like

$\int e^x \cos x dx$, allow M1 if two applications of “by parts”, despite incomplete method)

Final A ft for dividing all terms by candidates IF., providing “c” used.

[8]

2. (a) $[x > -2]$: Attempt to solve $x^2 - 1 = 3(1 - x)(x + 2)$ M1
 $[4x^2 + 3x - 7 = 0]$
 $x = 1$, or $\frac{5}{2}$ B1, A1

$[x < -2]$: Attempt to solve $x^2 - 1 = -3(1 - x)(x + 2)$ M1
Solving $x + 1 = 3x + 6$ (2 $x^2 + 3x - 5 = 0$) M1dep
 $x = -\frac{5}{2}$ A16

“Squaring”

If candidates do not notice the factor of $(x - 1)^2$ they have quartic to solve;

Squaring and finding quartic = 0 [8 $x^4 + 18x^3 - 25x^2 - 36x + 35 = 0$]

Finding one factor and factorising $(x - 1)(8x^3 + 26x^2 + x - 35) = 0$ M1

Finding one other factor and reducing other factor to quadratic,
likely to be $(x - 1)^2(8x^2 + 34x + 35) = 0$ M1

Complete factorisation $(x - 1)^2(2x + 5)(4x + 7) = 0$ M1

[Second M1 implies the first, if candidate starts there or cancels $(x - 1)^2$]

$x = 1$ B1, $x = -7/4$ A1, $x = -5/2$ A1

$x = 1$ allowed anywhere, no penalty in (b)

(b) $-\frac{7}{4} < x < 1$ One part M1
Both correct and enclosed A1
 $x < -\frac{5}{2}$ {Must be for $x < -2$ and only one value} B1ft3

Correct answers seen with no working is independent of (a)
(graphical calculator) mark as scheme.

Only allow the accuracy mark if no other interval, in both parts
≤ used penalise first time used

3. (a) $y = x^{-2} \Rightarrow \frac{dy}{dt} = -2x^{-3} \frac{dx}{dt}$ [Use of chain rule; need $\frac{dx}{dt}$] M1

$$\Rightarrow \frac{d^2y}{dt^2} = -2x^{-3} \frac{d^2x}{dt^2}, \quad + 6x^{-4} \left(\frac{dx}{dt} \right)^2$$

(÷ given d.e. by x^4) $\frac{2}{x^3} \frac{d^2x}{dt^2} - \frac{6}{x^4} \left(\frac{dx}{dt} \right)^2 = \frac{1}{x^2} - 3$

becomes $\left(-\frac{d^2y}{dt^2} = y - 3 \right)$ $\frac{d^2y}{dt^2} + y = 3$ AG A1ft, M1A1

A1 cs05

Second M1 is for attempt at product rule. (be generous)
Final A1 requires all working correct and sufficient “substitution” work

(b) Auxiliary equation: $m^2 + 1 = 0$ and produce
Complementary Function $y = \dots$ M1
 $(y) = A \cos t + B \sin t$ A1cao
Particular integral: $y = 3$ B1
 \therefore General solution: $(y) = A \cos t + B \sin t + 3$ A1ft4

Answer can be stated; M1 is implied by correct C.F. stated
(allow θ for t)

A1 f.t. for candidates CF + PI

Allow $m^2 + m = 0$ and $m^2 - 1 = 0$ for M1. Marks for (b) can be gained in (c)

(c) $\frac{1}{x^2} = A \cos t + B \sin t + 3$
 $x = \frac{1}{2}, t = 0 \Rightarrow (4 = A + 3) A = 1$ B1
Differentiating (to include $\frac{dx}{dt}$): $-2x^{-3} \frac{dx}{dt} = -A \sin t + B \cos t$ M1
 $\frac{dx}{dt} = 0, t = 0 \Rightarrow (0 = 0 + B) B = 0$ M1
 $\therefore \frac{1}{x^2} = 3 + \cos t$ so $x = \frac{1}{\sqrt{3 + \cos t}}$ A1 cao4

Second M : complete method to find other constant
(This may involve solving two equations in A and B)

(d) (Max. value of x when $\cos t = -1$) so $\max x = \frac{1}{\sqrt{2}}$ or AWRT 0.707 B11

4. (a) $\frac{x \frac{dx}{d\theta} r \cos \theta - 4 \sin \theta \cos^3 \theta}{d\theta} = 4 \cos^4 \theta - 12 \cos^2 \theta \sin^2 \theta$ M1

any correct expression

Solving $\frac{dx}{d\theta} = 0 \quad \left[\frac{dx}{d\theta} = 0 \Rightarrow 4 \cos^2 \theta (\cos^2 \theta - 3 \sin^2 \theta) = 0 \right]$ M1A1

$\sin \theta = \frac{1}{2}$ or $\cos \theta = \frac{\sqrt{3}}{2}$ or $\tan \theta = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6}$ M1

$r = 4 \sin \frac{\pi}{6} \cos^2 \frac{\pi}{6} = \frac{3}{2}$ AG A1 cso

AG A1cs06

So many ways x may be expressing e.g.

$$2 \sin 2\theta \cos^2 \theta, \sin 2\theta(1 + \cos 2\theta), \sin 2\theta + (1/2) \sin 4\theta$$

leading to many results for $\frac{dx}{d\theta}$

Some relevant equations in solving

$$[(1 - 4 \sin^2 \theta) = 0, (4 \cos^2 \theta - 3) = 0, (1 - 3 \tan^2 \theta) = 0, \cos 3\theta = 0]$$

Showing that $\theta = \frac{\pi}{6}$ satisfies $\frac{dx}{d\theta} = 0$, allow M1 A1

providing $\frac{dx}{d\theta}$ correct

Starting with $x = r \sin \theta$ can gain M0M1M1

(b) $A = \frac{1}{2} \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} r^2 d\theta = \frac{1}{2} \cdot 16 \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \sin^2 \theta \cos^4 \theta d\theta$

$$8 \sin^2 \theta \cos^4 \theta = 2 \cos^2 \theta (4 \sin^2 \theta \cos^2 \theta) = 2 \cos^2 \theta \sin^2 2\theta$$
 M1
$$= (\cos 2\theta + 1) \sin^2 2\theta$$
 M1
$$= \cos 2\theta \sin^2 2\theta + \frac{1 - \cos 4\theta}{2} = \text{Answer}$$
 AG A1 cs03

First M1 for use of double angle formula for $\sin 2A$

Second M1 for use of $\cos 2A = 2 \cos^2 A - 1$

Answer given: must be intermediate step, as shown, and no incorrect work

$$\begin{aligned}
 (c) \quad \text{Area} &= \left[\frac{1}{6} \sin^3 2\theta + \frac{\theta}{2} - \frac{\sin 4\theta}{8} \right]_{\left(\frac{\pi}{6}\right)}^{\left(\frac{\pi}{4}\right)} && \text{ignore limits} && \text{M1A1} \\
 &= \left(\frac{1}{6} \sin^3 \frac{\pi}{2} + \frac{\pi}{8} - \frac{\sin \pi}{8} \right) - \left(\frac{1}{6} \sin^3 \frac{\pi}{3} + \frac{\pi}{12} - \frac{\sin \frac{2\pi}{3}}{8} \right) && \text{(sub. limits)} && \text{M1} \\
 &= \left(\frac{1}{6} + \frac{\pi}{8} \right) - \left(\frac{\sqrt{3}}{16} + \frac{\pi}{12} - \frac{\sqrt{3}}{16} \right) = \frac{1}{6}, + \frac{\pi}{24} && \text{both cao} && \text{A1, A15}
 \end{aligned}$$

For first M, of the form $a \sin^3 2\theta + \frac{\theta}{2} \pm b \sin 4\theta$ (Allow if two of correct form)

On ePen the order of the As in answer is as written

[14]

5. $1\frac{1}{2}$ and 3 are ‘critical values’, e.g. used in solution, or both seen as asymptotes. B1
- $$(x+1)(x-3) = 2x - 3 \Rightarrow x(x-4) = 0$$
- $$x = 4, x = 0$$
- M1A1, A1

M1: Attempt to find at least one other critical value

$$0 < x < 1\frac{1}{2}, 3 < x < 4 \quad \text{M1A1, A17}$$

M1: An inequality using $1\frac{1}{2}$ or 3

First M mark can be implied by the two correct values, but otherwise a method must be seen. (The method may be graphical, but either $(x =) 4$ or $(x =) 0$ needs to be clearly written or used in this case).
Ignore ‘extra values’ which might arise through ‘squaring both sides’ methods.

\leq appearing: maximum one A mark penalty (final mark).

[7]

6. Integrating factor $e^{\int -\tan x dx} = e^{\ln(\cos x)}$ (or $e^{-\ln(\sec x)}$), $= \cos x$ (or $\frac{1}{\sec x}$) M1, A1
 $\left(\cos x \frac{dy}{dx} - y \sin x = 2 \sec^2 x \right)$
 $y \cos x = \int 2 \sec^2 x dx$ (or equiv.) $\left(\text{Or: } \frac{d}{dx}(y \cos x) = 2 \sec^2 x \right)$ M1A1(ft)
 $y \cos x = 2 \tan x (+C)$ (or equiv.) A1
 $y = 3 \tan x + C$ M1
 $y = \frac{\cos x}{\cos x}$ (Or equiv. in the form $y = f(x)$) A17

1st M: Also scored for $e^{\int \tan x dx} = e^{-\ln(\cos x)}$ (or $e^{\ln(\sec x)}$), then A0 for $\sec x$.

2nd M: Attempt to use their integrating factor (requires one side of the equation 'correct' for their integrating factor).

2nd A: The follow-through is allowed only in the case where the integrating factor used is $\sec x$ or $-\sec x$. $\left(y \sec x = \int 2 \sec^4 x dx \right)$

3rd M: Using $y = 3$ at $x = 0$ to find a value for C (dependent on an integration attempt, however poor, on the RHS).

Alternative

1st M: Multiply through the given equation by $\cos x$.

1st A: Achieving $\cos x \frac{dy}{dx} - y \sin x = 2 \sec^2 x$. (Allowing the possibility of integrating by inspection). [7]

7. C.F. $m^2 + 3m + 2 = 0$ $m = -1$ and $m = -2$ M1
 $y = Ae^{-x} + Be^{-2x}$ A12
- P.I. $y = cx^2 + dx + e$ B1
 $\frac{dy}{dx} = 2cx + d, \frac{d^2y}{dx^2} = 2c$ $2c + 3(2cx + d) + 2(cx^2 + dx + e) \equiv 2x^2 + 6x$ M1
 $2c = 2$ $c = 1$ (One correct value) A1
 $6c + 2d = 6$ $d = 0$
 $2c + 3d + 2e = 0$ $e = -1$ (Other two correct values) A1
General soln: $y = Ae^{-x} + Be^{-2x} + x^2 - 1$ (Their C.F. + their P.I.) A1ft5

$$\frac{x \frac{dy}{dx}}{dx} = 0, y = 1; 1 = A + B - 1 \\ -Ae^{-x} - 2Be^{-2x} + 2x, x = 0, \frac{dy}{dx} = 1$$

(A + B = 2) M1
1 = -A - 2B M1

Solving simultaneously: $A = 5$ and $B = -3$ M1A1
 Solution: $y = 5e^{-x} - 3e^{-2x} + x^2 - 1$ A15

1st M: Attempt to solve auxiliary equation.

2nd M: Substitute their $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ into the D>E> to form an identity in x with unknown constants.


3rd M: Using $y = 1$ at $x = 0$ in their general solution to find an equation in A and B .

4th M: Differentiating their general solution (condone ‘slips’, but the powers of each term must be correct) and using $\frac{dy}{dx} = 1$ at $x = 0$ to find an equation in A and B .

5th M: Solving simultaneous equations to find both a value of A and a value of B .

[12]

8. (a)



Shape (close curve, approx. symmetrical about the initial line, in all ‘quadrants’ and ‘centred’ to the right of the pole/origin).

B1

Shape (at least one correct ‘intercept’ r value... shown on sketch or perhaps seen in a table).

B12

(Also allow awrt 3.27 or awrt 6.73).

(b) $\frac{dy}{d\theta} r \sin \theta = 5 \sin \theta + \sqrt{3} \sin \theta \cos \theta$ M1
 $\frac{d\theta}{d\theta} = 5 \cos \theta - \sqrt{3} \sin^2 \theta + \sqrt{3} \cos^2 \theta (= 5 \cos \theta + \sqrt{3} \cos 2\theta)$ A1
 $5 \cos \theta - \sqrt{3}(1 - \cos^2 \theta) + \sqrt{3} \cos^2 \theta = 0$ M1
 $2\sqrt{3} \cos^2 \theta + 5 \cos \theta - \sqrt{3} = 0$
 $(2\sqrt{3} \cos \theta - 1)(\cos \theta + \sqrt{3}) = 0$ $\cos \theta = \dots (0.288\dots)$ M1
 $\theta = 1.28$ and 5.01 (awrt) (Allow ± 1.28 awrt) $\cos \theta = \text{Also allow } \pm \arccos \frac{1}{2\sqrt{3}}$ A1
 $r = 5 + \sqrt{3} \left(\frac{1}{2\sqrt{3}} \right) = \frac{11}{2}$ (Allow awrt 5.50) A16

2nd M: Forming a quadratic in $\cos \theta$.

3rd M: Solving a 3 term quadratic to find a value of $\cos \theta$ (even if called θ).

Speacial case: Working with $r \cos \theta$ instead of $r \sin \theta$.

1st M1 for $r \cos \theta = 5 \cos \theta + \sqrt{3} \cos^2 \theta$

1st A1 for derivative – $5 \sin \theta - 2\sqrt{3} \sin \theta \cos \theta$, then no further marks.

(c) $r^2 = 25 + 10\sqrt{3} \cos \theta + 3 \cos^2 \theta$ B1
 $\int 25 + 10\sqrt{3} \cos \theta + 3 \cos^2 \theta d\theta = \frac{53\theta}{2} + 10\sqrt{3} \sin \theta + 3 \left(\frac{\sin 2\theta}{4} \right)$ M1 A1ft A1ft
(ft for integration of $(a + b \cos \theta)$ and $c \cos 2\theta$ respectively)
 $\frac{1}{2} \left[25\theta + 10\sqrt{3} \sin \theta + \frac{3 \sin 2\theta}{4} + \frac{3\theta}{2} \right]_0^{2\pi} = \dots$ M1
 $= \frac{1}{2} (50\pi + 3\pi) = \frac{53\pi}{2}$ or equiv. in terms of π . A16

1st M: Attempt to integrate at least one term.

2nd M: Requires use of the $\frac{1}{2}$, correct limits (which could be 0

to 2π , or $-\pi$ to π , or ‘double’ 0 to π), and subtraction (which could be implied).

[14]

9. (a) $\frac{y_1 - 0.2}{0.1} \approx \left(\frac{dy}{dx} \right)_0 = 0.2 \times e^0 (= 0.2)$ M1
 $y_1 \approx 0.22$ A12

$$(b) \quad \left(\frac{dy}{dx} \right)_0 = 0.22 \times e^{0.01} \approx 0.2222... \quad B1$$

$$\frac{y_2 - 0.2}{0.2} \approx 0.2222... \quad M1$$

$$y_2 \approx 0.2444 \quad cao \quad A13$$

[5]

10. (a) $(1-x^2) \frac{d^3y}{dx^3} - 2x \frac{d^2y}{dx^2} - x \frac{d^2y}{dx^2} - \frac{dy}{dx} + 2 \frac{dy}{dx} = 0 \quad M1$

At $x=0$, $\frac{d^3y}{dx^3} = -\frac{dy}{dx} = 1 \quad M1A1cso3$

(b) $\left(\frac{d^2y}{dx^2} \right)_0 = -4 \quad Allow \text{ anywhere} \quad B1$

$$y = f(0) + f'(0)x + \frac{f''(0)}{2}x^2 + \frac{f'''(0)}{6}x^3 + ...$$

$$= 2 - x - 2x^2, + \frac{1}{6}x^3 + ... \quad M1A1ft, A1 (\text{dep})4$$

[7]

11. (a) $z^n = (\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$
 $z^{-n} = (\cos \theta + i \sin \theta)^{-n} = \cos(-n\theta) + i \sin(-n\theta) = \cos n\theta - i \sin n\theta \quad \text{both} \quad M1$

Adding $z^n + \frac{1}{z^n} = 2 \cos n\theta * \quad cso \quad A12$

(b) $\left(z + \frac{1}{z} \right)^6 = z^6 + 6z^4 + 15z^2 + 20 + 15z^{-2} + 6z^{-4} + z^{-6} \quad M1$

$$= z^6 + z^{-6} + 6(z^4 + z^{-4}) + 15(z^2 + z^{-2}) + 20 \quad M1$$

$$64 \cos^6 \theta = 2 \cos 6\theta + 12 \cos 4\theta + 30 \cos 2\theta + 20 \quad M1$$

$$32 \cos^6 \theta = \cos 6\theta + 6 \cos 4\theta + 15 \cos 2\theta + 10 \quad A1, A1$$

$(p=1, q=6, r=15, s=10) \quad A1 \text{ any two correct}$ 5

$$(c) \quad \int \cos^6 \theta d\theta = \left(\frac{1}{32} \right) \int (\cos 6\theta + 6 \cos 4\theta + 15 \cos 2\theta + 10) d\theta$$

$$= \left(\frac{1}{32} \right) \left[\frac{\sin 6\theta}{6} + \frac{6 \sin 4\theta}{4} + \frac{15 \sin 2\theta}{2} + 10\theta \right]$$

$$\left[\dots \right]_0^{\frac{\pi}{3}} = \frac{1}{32} \left[-\frac{3}{2} \times \frac{\sqrt{3}}{2} + \frac{15}{2} \times \frac{\sqrt{3}}{2} + \frac{10\pi}{3} \right] = \frac{5\pi}{48} + \frac{3\sqrt{3}}{32}$$

M1A1ft
M1A14

or exact equivalent

[11]

12. (a) Let $z = \lambda + \lambda i$; $w = \frac{\lambda + (\lambda + 1)i}{\lambda(1+i)}$

$$= \frac{\lambda + (\lambda + 1)i}{\lambda(1+i)} \times \frac{1-i}{1-i}$$

$$u + iv = \frac{(2\lambda + 1) + i}{2\lambda}$$

$$u = 1 + \frac{1}{2\lambda}, v = \frac{1}{2\lambda}$$

Eliminating λ gives a line with equation $v = u - 1$ or equivalent

M1
M1
A1
M1
A15

(b) Let $z = \lambda - (\lambda + 1)i$; $w = \frac{\lambda - \lambda i}{\lambda - (\lambda + 1)i}$

$$= \frac{\lambda - \lambda i}{\lambda - (\lambda + 1)i} \times \frac{\lambda + (\lambda + 1)i}{\lambda + (\lambda + 1)i}$$

$$u + iv = \frac{\lambda(2\lambda + 1) + \lambda i}{2\lambda^2 + 2\lambda + 1}$$

$$u = \frac{\lambda(2\lambda + 1)}{2\lambda^2 + 2\lambda + 1}, v = \frac{\lambda}{2\lambda^2 + 2\lambda + 1}$$

$$\frac{u}{v} = 2\lambda + 1$$

$$v = \frac{2\lambda}{4\lambda^2 + 4\lambda + 2} = \frac{(2\lambda + 1) - 1}{(2\lambda + 1)^2 + 1} = \frac{\frac{u}{v} - 1}{\left(\frac{u}{v}\right)^2 + 1}$$

Reducing to the circle with equation $u^2 + v^2 - u + v = 0$ * cso

M1
M1
A1
M1
M1
M1
M1
M1
M1A17

Alternative 1

$$\text{Let } z = \lambda - (\lambda + 1)i; w = \frac{\lambda - \lambda i}{\lambda - (\lambda + 1)i} \quad \text{M1}$$

$$= \frac{\lambda - \lambda i}{\lambda - (\lambda + 1)i} \times \frac{\lambda + (\lambda + 1)i}{\lambda + (\lambda + 1)i} \quad \text{M1}$$

$$u + iv = \frac{\lambda(2\lambda + 1) + \lambda i}{2\lambda^2 + 2\lambda + 1} \quad \text{A1}$$

$$u = \frac{\lambda(2\lambda + 1)}{2\lambda^2 + 2\lambda + 1}, v = \frac{\lambda}{2\lambda^2 + 2\lambda + 1} \quad \text{M1}$$

$$\begin{aligned} u^2 + v^2 - u + v &= \left(\frac{\lambda(2\lambda + 1)}{2\lambda^2 + 2\lambda + 1} \right)^2 + \left(\frac{\lambda}{2\lambda^2 + 2\lambda + 1} \right)^2 - \frac{\lambda(2\lambda + 1)}{2\lambda^2 + 2\lambda + 1} + \frac{\lambda}{2\lambda^2 + 2\lambda + 1} \\ &= \frac{(4\lambda^4 + 4\lambda^3 + \lambda^2) + \lambda^2 - 2\lambda^2(2\lambda^2 + 2\lambda + 1)}{(2\lambda^2 + 2\lambda + 1)^2} \quad \text{M1} \\ &= 0^* \quad \text{M1A1} \end{aligned}$$

Alternative 2

$$\text{Let } z = \lambda - (\lambda + 1)i; u + iv = \frac{\lambda - \lambda i}{\lambda - (\lambda + 1)i} \quad \text{M1}$$

$$(u + iv)(\lambda - (\lambda + 1)i) = \lambda - \lambda i \quad \text{M1}$$

$$u\lambda + v(\lambda + 1) + [v\lambda - u(\lambda + 1)]i = \lambda - \lambda i \quad \text{A1}$$

Equating real & imaginary parts


$$u\lambda + v(\lambda + 1) = \lambda \text{ (i)} \quad v\lambda - \lambda u - u = -\lambda \text{ (ii)} \quad \text{M1}$$

$$\text{From (i) } \lambda = \frac{v}{1-u-v} \quad \text{From (ii) } \lambda = \frac{u}{1-u+v}$$

$$\frac{v}{1-u-v} = \frac{u}{1-u+v} \quad \text{M1}$$

$$\text{Reducing to the circle with equation } u^2 + v^2 - u + v = 0^* \quad \text{M1A1}$$

(c)



ft their line
Circle through origin, centre in correct quadrant
Intersection correctly placed

B1ft
B1
B13
[15]