

Mark Scheme (Results)

Summer 2009

GCE

GCE Mathematics (6668/01)

June 2009
6668 Further Pure Mathematics FP2 (new)
Mark Scheme

Question Number	Scheme	Marks
Q1 (a)	$\frac{1}{r(r+2)} = \frac{1}{2r} - \frac{1}{2(r+2)}$	$\frac{1}{2r} - \frac{1}{2(r+2)}$ B1 aef (1)
(b)	$\begin{aligned} \sum_{r=1}^n \frac{4}{r(r+2)} &= \sum_{r=1}^n \left(\frac{2}{r} - \frac{2}{r+2} \right) \\ &= \left(\frac{2}{1} - \frac{2}{3} \right) + \left(\frac{2}{2} - \frac{2}{4} \right) + \dots \\ &\quad \dots + \left(\frac{2}{n-1} - \frac{2}{n+1} \right) + \left(\frac{2}{n} - \frac{2}{n+2} \right) \end{aligned}$ <p style="text-align: right;">List the first two terms and the last two terms</p> $\begin{aligned} &= \frac{2}{1} + \frac{2}{2} ; - \frac{2}{n+1} - \frac{2}{n+2} \\ &= 3 - \frac{2}{n+1} - \frac{2}{n+2} \\ &= \frac{3(n+1)(n+2) - 2(n+2) - 2(n+1)}{(n+1)(n+2)} \\ &= \frac{3n^2 + 9n + 6 - 2n - 4 - 2n - 2}{(n+1)(n+2)} \\ &= \frac{3n^2 + 5n}{(n+1)(n+2)} \\ &= \frac{n(3n+5)}{(n+1)(n+2)} \end{aligned}$ <p style="text-align: right;">Attempt to combine to an at least 3 term fraction to a single fraction and an attempt to take out the brackets from their numerator.</p> <p style="text-align: right;">Correct Result</p>	M1 M1 A1 M1 M1 A1 cso AG (5) [6]

Question Number	Scheme	Marks
Q2 (a)	$z^3 = 4\sqrt{2} - 4\sqrt{2}i, \quad -\pi < \theta \leq \pi$	
	$r = \sqrt{(4\sqrt{2})^2 + (-4\sqrt{2})^2} = \sqrt{32 + 32} = \sqrt{64} = 8$ $\theta = -\tan^{-1}\left(\frac{4\sqrt{2}}{4\sqrt{2}}\right) = -\frac{\pi}{4}$	A valid attempt to find the modulus and argument of $4\sqrt{2} - 4\sqrt{2}i.$ M1
	$z^3 = 8\left(\cos\left(-\frac{\pi}{4}\right) + i\sin\left(-\frac{\pi}{4}\right)\right)$	
	$\text{So, } z = (8)^{\frac{1}{3}}\left(\cos\left(\frac{-\frac{\pi}{4}}{3}\right) + i\sin\left(\frac{-\frac{\pi}{4}}{3}\right)\right)$	Taking the cube root of the modulus and dividing the argument by 3. M1
	$\Rightarrow z = 2\left(\cos\left(-\frac{\pi}{12}\right) + i\sin\left(-\frac{\pi}{12}\right)\right)$	$2\left(\cos\left(-\frac{\pi}{12}\right) + i\sin\left(-\frac{\pi}{12}\right)\right)$ A1
	$\text{Also, } z^3 = 8\left(\cos\left(\frac{7\pi}{4}\right) + i\sin\left(\frac{7\pi}{4}\right)\right)$ $\text{or } z^3 = 8\left(\cos\left(-\frac{9\pi}{4}\right) + i\sin\left(-\frac{9\pi}{4}\right)\right)$	Adding or subtracting 2π to the argument for z^3 in order to find other roots. M1
	$\Rightarrow z = 2\left(\cos\frac{7\pi}{12} + i\sin\frac{7\pi}{12}\right)$	Any one of the final two roots A1
	$\text{and } z = 2\left(\cos\left(\frac{-3\pi}{4}\right) + i\sin\left(\frac{-3\pi}{4}\right)\right)$	Both of the final two roots. A1
	Special Case 1: Award SC: M1M1A1M1A0A0 for ALL three of $2\left(\cos\frac{\pi}{12} + i\sin\frac{\pi}{12}\right)$, $2\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right)$ and $2\left(\cos\left(\frac{-7\pi}{12}\right) + i\sin\left(\frac{-7\pi}{12}\right)\right).$	[6]
	Special Case 2: If r is incorrect (and not equal to 8) and candidate states the brackets () correctly then give the first accuracy mark ONLY where this is applicable.	

Question Number	Scheme	Marks
Q3	$\sin x \frac{dy}{dx} - y \cos x = \sin 2x \sin x$ $\frac{dy}{dx} - \frac{y \cos x}{\sin x} = \frac{\sin 2x \sin x}{\sin x}$ <p>An attempt to divide every term in the differential equation by $\sin x$. Can be implied.</p> $\frac{dy}{dx} - \frac{y \cos x}{\sin x} = \sin 2x$ <p>Integrating factor = $e^{\int -\frac{\cos x}{\sin x} dx} = e^{-\ln \sin x}$</p> $= \frac{1}{\sin x}$ $\left(\frac{1}{\sin x} \right) \frac{dy}{dx} - \frac{y \cos x}{\sin^2 x} = \frac{\sin 2x}{\sin x}$ $\frac{d}{dx} \left(\frac{y}{\sin x} \right) = \sin 2x \times \frac{1}{\sin x}$ $\frac{d}{dx} \left(\frac{y}{\sin x} \right) = 2 \cos x$ $\frac{y}{\sin x} = \int 2 \cos x \, dx$ $\frac{y}{\sin x} = 2 \sin x + K$ $y = 2 \sin^2 x + K \sin x$ <p>A credible attempt to integrate the RHS with/without $+ K$</p> $y = 2 \sin^2 x + K \sin x$	M1 dM1 A1 aef A1 aef M1 A1

Question Number	Scheme	Marks
Q4	<p> $A = \frac{1}{2} \int_0^{2\pi} (a + 3\cos\theta)^2 d\theta$ $(a + 3\cos\theta)^2 = a^2 + 6a\cos\theta + 9\cos^2\theta$ $= a^2 + 6a\cos\theta + 9\left(\frac{1 + \cos 2\theta}{2}\right)$ $\underline{\qquad\qquad\qquad}$ $A = \frac{1}{2} \int_0^{2\pi} \left(a^2 + 6a\cos\theta + \frac{9}{2} + \frac{9}{2}\cos 2\theta\right) d\theta$ $= \left(\frac{1}{2}\right) \left[a^2\theta + 6a\sin\theta + \frac{9}{2}\theta + \frac{9}{4}\sin 2\theta \right]_0^{2\pi}$ $= \frac{1}{2} [(2\pi a^2 + 0 + 9\pi + 0) - (0)]$ $= \pi a^2 + \frac{9\pi}{2}$ Hence, $\pi a^2 + \frac{9\pi}{2} = \frac{107}{2}\pi$ $a^2 + \frac{9}{2} = \frac{107}{2}$ $a^2 = 49$ As $a > 0$, $a = 7$ Some candidates may achieve $a = 7$ from incorrect working. Such candidates will not get full marks </p> <p> Applies $\frac{1}{2} \int_0^{2\pi} r^2 (d\theta)$ with correct limits. Ignore $d\theta$. $\cos^2\theta = \frac{\pm 1 \pm \cos 2\theta}{2}$ <u>Correct underlined expression.</u> Integrated expression with at least 3 out of 4 terms of the form $\pm A\theta \pm B\sin\theta \pm C\theta \pm D\sin 2\theta$. Ignore the $\frac{1}{2}$. Ignore limits. $a^2\theta + 6a\sin\theta +$ correct ft integration. Ignore the $\frac{1}{2}$. Ignore limits. $\pi a^2 + \frac{9\pi}{2}$ Integrated expression equal to $\frac{107}{2}\pi$. $a = 7$ </p>	B1 M1 A1 M1* A1 ft A1 dm1* A1 cso [8]

Question Number	Scheme	Marks
Q5	$y = \sec^2 x = (\sec x)^2$	
(a)	$\frac{dy}{dx} = 2(\sec x)^1(\sec x \tan x) = 2\sec^2 x \tan x$ <p>Apply product rule:</p> $\left\{ \begin{array}{l} u = 2\sec^2 x \\ \frac{du}{dx} = 4\sec^2 x \tan x \end{array} \right. \quad \left. \begin{array}{l} v = \tan x \\ \frac{dv}{dx} = \sec^2 x \end{array} \right\}$ $\frac{d^2y}{dx^2} = 4\sec^2 x \tan^2 x + 2\sec^4 x$ $= 4\sec^2 x(\sec^2 x - 1) + 2\sec^4 x$ <p>Hence, $\frac{d^2y}{dx^2} = 6\sec^4 x - 4\sec^2 x$</p>	Either $2(\sec x)^1(\sec x \tan x)$ or $2\sec^2 x \tan x$ B1 aef M1 A1
(b)	$y_{\frac{\pi}{4}} = (\sqrt{2})^2 = 2, \left(\frac{dy}{dx} \right)_{\frac{\pi}{4}} = 2(\sqrt{2})^2(1) = 4$ $\left(\frac{d^2y}{dx^2} \right)_{\frac{\pi}{4}} = 6(\sqrt{2})^4 - 4(\sqrt{2})^2 = 24 - 8 = 16$ $\frac{d^3y}{dx^3} = 24\sec^3 x(\sec x \tan x) - 8\sec x(\sec x \tan x)$ $= 24\sec^4 x \tan x - 8\sec^2 x \tan x$ $\left(\frac{d^2y}{dx^2} \right)_{\frac{\pi}{4}} = 24(\sqrt{2})^4(1) - 8(\sqrt{2})^2(1) = 96 - 16 = 80$ $\sec x \approx 2 + 4(x - \frac{\pi}{4}) + \frac{16}{2}(x - \frac{\pi}{4})^2 + \frac{80}{6}(x - \frac{\pi}{4})^3 + \dots$ $\left\{ \sec x \approx 2 + 4(x - \frac{\pi}{4}) + 8(x - \frac{\pi}{4})^2 + \frac{40}{3}(x - \frac{\pi}{4})^3 + \dots \right\}$	Two terms added with one of either $A \sec^2 x \tan^2 x$ or $B \sec^4 x$ in the correct form. Correct differentiation Applies $\tan^2 x = \sec^2 x - 1$ leading to the correct result. Both $y_{\frac{\pi}{4}} = 2$ and $\left(\frac{dy}{dx} \right)_{\frac{\pi}{4}} = 4$ Attempts to substitute $x = \frac{\pi}{4}$ into both terms in the expression for $\frac{d^2y}{dx^2}$. Two terms differentiated with either $24\sec^4 x \tan x$ or $-8\sec^2 x \tan x$ being correct $\left(\frac{d^3y}{dx^3} \right)_{\frac{\pi}{4}} = 80$ Applies a Taylor expansion with at least 3 out of 4 terms ft correctly. Correct Taylor series expansion. [10]

Question Number	Scheme	Marks
Q6	$w = \frac{z}{z + i}$, $z = -i$	
(a)	$w(z + i) = z \Rightarrow wz + iw = z \Rightarrow iw = z - wz$ $\Rightarrow iw = z(1 - w) \Rightarrow z = \frac{iw}{(1 - w)}$	Complete method of rearranging to make z the subject. $z = \frac{iw}{(1 - w)}$ M1 A1 aef
	$ z = 3 \Rightarrow \left \frac{iw}{1 - w} \right = 3$	Putting $ z $ in terms of their w = 3 dM1
	$\left. \begin{aligned} iw &= 3 1 - w \Rightarrow w = 3 w - 1 \Rightarrow w ^2 = 9 w - 1 ^2 \\ \Rightarrow u + iv ^2 &= 9 u + iv - 1 ^2 \end{aligned} \right\}$	
	$\Rightarrow u^2 + v^2 = 9[(u - 1)^2 + v^2]$	Applies $w = u + iv$, and uses Pythagoras correctly to get an equation in terms of u and v without any i 's. Correct equation. ddM1
	$\left. \begin{aligned} \Rightarrow u^2 + v^2 &= 9u^2 - 18u + 9 + 9v^2 \\ \Rightarrow 0 &= 8u^2 - 18u + 8v^2 + 9 \end{aligned} \right\}$	A1
	$\Rightarrow 0 = u^2 - \frac{9}{4}u + v^2 + \frac{9}{8}$	Simplifies down to $u^2 + v^2 \pm \alpha u \pm \beta v \pm \delta = 0$. dddM1
	$\Rightarrow (u - \frac{9}{8})^2 - \frac{81}{64} + v^2 + \frac{9}{8} = 0$	
	$\Rightarrow (u - \frac{9}{8})^2 + v^2 = \frac{9}{64}$	
	{Circle} centre $(\frac{9}{8}, 0)$, radius $\frac{3}{8}$	One of centre or radius correct. Both centre and radius correct. A1 A1 (8)
(b)		Circle indicated on the Argand diagram in the correct position in follow through quadrants. Ignore plotted coordinates. B1ft
	Region outside a circle indicated only.	B1 (2)
		[10]

Question Number	Scheme	Marks
Q7 (a)	$y = x^2 - a^2 , a > 1$ <p>Correct Shape. Ignore cusps. Correct coordinates.</p>	B1 B1
(b)	$ x^2 - a^2 = a^2 - x, a > 1$ $\{ x > a\}, \quad x^2 - a^2 = a^2 - x$ $\Rightarrow x^2 + x - 2a^2 = 0$ $\Rightarrow x = \frac{-1 \pm \sqrt{1 - 4(1)(-2a^2)}}{2}$ $\Rightarrow x = \frac{-1 \pm \sqrt{1 + 8a^2}}{2}$ $\{ x < a\}, \quad -x^2 + a^2 = a^2 - x$ $\{\Rightarrow x^2 - x = 0 \Rightarrow x(x - 1) = 0\}$ $\Rightarrow x = 0, 1$	(2) M1 aef M1 A1 M1 aef B1 A1
(c)	$ x^2 - a^2 > a^2 - x, a > 1$ $x < \frac{-1 - \sqrt{1 + 8a^2}}{2} \quad \text{or} \quad x > \frac{-1 + \sqrt{1 + 8a^2}}{2}$ $\{\text{or}\} \quad 0 < x < 1$ <p style="text-align: center;">x is less than their least value x is greater than their maximum value</p> <p>For $\{ x < a\}$, Lowest $< x <$ Highest $0 < x < 1$</p>	(6) B1 ft B1 ft M1 A1
		[12]

Question Number	Scheme	Marks
<p>Q8</p> <p>(a)</p> <p>$\frac{d^2x}{dt^2} + 5\frac{dx}{dt} + 6x = 2e^{-t}$, $x = 0$, $\frac{dx}{dt} = 2$ at $t = 0$.</p> <p>AE, $m^2 + 5m + 6 = 0 \Rightarrow (m+3)(m+2) = 0$ $\Rightarrow m = -3, -2$.</p> <p>So, $x_{CF} = Ae^{-3t} + Be^{-2t}$</p> <p>$\left\{ x = k e^{-t} \Rightarrow \frac{dx}{dt} = -k e^{-t} \Rightarrow \frac{d^2x}{dt^2} = k e^{-t} \right\}$</p> <p>$\Rightarrow k e^{-t} + 5(-k e^{-t}) + 6k e^{-t} = 2e^{-t} \Rightarrow 2k e^{-t} = 2e^{-t}$ $\Rightarrow k = 1$</p> <p>{ So, $x_{PI} = e^{-t}$ }</p> <p>So, $x = Ae^{-3t} + Be^{-2t} + e^{-t}$</p> <p>$\frac{dx}{dt} = -3Ae^{-3t} - 2Be^{-2t} - e^{-t}$</p> <p>$t = 0, x = 0 \Rightarrow 0 = A + B + 1$ $t = 0, \frac{dx}{dt} = 2 \Rightarrow 2 = -3A - 2B - 1$</p> <p>$\left\{ \begin{array}{l} 2A + 2B = -2 \\ -3A - 2B = 3 \end{array} \right\}$</p> <p>$\Rightarrow A = -1, B = 0$</p> <p>So, $x = -e^{-3t} + e^{-t}$</p>	<p>$Ae^{m_1 t} + Be^{m_2 t}$, where $m_1 \neq m_2$. $Ae^{-3t} + Be^{-2t}$</p> <p>Substitutes $k e^{-t}$ into the differential equation given in the question. Finds $k = 1$.</p> <p>their x_{CF} + their x_{PI}</p> <p>Finds $\frac{dx}{dt}$ by differentiating their x_{CF} and their x_{PI}</p> <p>Applies $t = 0, x = 0$ to x and $t = 0, \frac{dx}{dt} = 2$ to $\frac{dx}{dt}$ to form simultaneous equations.</p> <p>$x = -e^{-3t} + e^{-t}$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1*</p> <p>dM1*</p> <p>ddM1*</p> <p>A1 cao (8)</p>

Question Number	Scheme	Marks
(b)	<p>$x = -e^{-3t} + e^{-t}$</p> <p>$\frac{dx}{dt} = 3e^{-3t} - e^{-t} = 0$</p> <p>$3 - e^{2t} = 0$ $\Rightarrow t = \frac{1}{2}\ln 3$</p> <p>So, $x = -e^{-\frac{3}{2}\ln 3} + e^{-\frac{1}{2}\ln 3} = -e^{\ln 3^{-\frac{3}{2}}} + e^{\ln 3^{-\frac{1}{2}}}$</p> <p>$x = -3^{-\frac{3}{2}} + 3^{-\frac{1}{2}}$</p> <p>$= -\frac{1}{3\sqrt{3}} + \frac{1}{\sqrt{3}} = \frac{2}{3\sqrt{3}} = \frac{2\sqrt{3}}{9}$</p> <p>$\frac{d^2x}{dt^2} = -9e^{-3t} + e^{-t}$</p> <p>At $t = \frac{1}{2}\ln 3$, $\frac{d^2x}{dt^2} = -9e^{-\frac{3}{2}\ln 3} + e^{-\frac{1}{2}\ln 3}$</p> <p>$= -9(3)^{-\frac{3}{2}} + 3^{-\frac{1}{2}} = -\frac{9}{3\sqrt{3}} + \frac{1}{\sqrt{3}} = -\frac{3}{\sqrt{3}} + \frac{1}{\sqrt{3}}$</p> <p>As $\frac{d^2x}{dt^2} = -\frac{9}{3\sqrt{3}} + \frac{1}{\sqrt{3}} = \left\{ -\frac{2}{\sqrt{3}} \right\} < 0$ then x is maximum.</p> <p>Differentiates their x to give $\frac{dx}{dt}$ and puts $\frac{dx}{dt}$ equal to 0. A credible attempt to solve. $t = \frac{1}{2}\ln 3$ or $t = \ln\sqrt{3}$ or awrt 0.55</p> <p>Substitutes their t back into x and an attempt to eliminate out the ln's.</p> <p>uses exact values to give $\frac{2\sqrt{3}}{9}$</p> <p>Finds $\frac{d^2x}{dt^2}$ and substitutes their t into $\frac{d^2x}{dt^2}$</p> <p>$-\frac{9}{3\sqrt{3}} + \frac{1}{\sqrt{3}} < 0$ and maximum conclusion.</p>	M1 M1 dM1* A1 ddM1 A1 AG dM1* A1 (7) [15]