

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise A, Question 1

Question:

Integrate the following with respect to x .

- a $\sinh x + 3 \cosh x$
- b $5 \operatorname{sech}^2 x$
- c $\frac{1}{\sinh^2 x}$
- d $\cosh x - \frac{1}{\cosh^2 x}$
- e $\frac{\sinh x}{\cosh^2 x}$
- f $\frac{3}{\sinh x \tanh x}$
- g $\operatorname{sech} x (\operatorname{sech} x + \tanh x)$
- h $(\operatorname{sech} x + \operatorname{cosech} x)(\operatorname{sech} x - \operatorname{cosech} x)$

Solution:

- a $\int (\sinh x + 3 \cosh x) dx = \cosh x + 3 \sinh x + C$
- b $\int 5 \operatorname{sech}^2 x dx = 5 \tanh x + C$
- c $\int \frac{1}{\sinh^2 x} dx = \int \operatorname{cosech}^2 x dx = -\operatorname{coth} x + C$
- d $\int \left(\cosh x - \frac{1}{\cosh^2 x} \right) dx = \int (\cosh x - \operatorname{sech}^2 x) dx = \sinh x - \tanh x + C$
- e $\int \frac{\sinh x}{\cosh^2 x} dx = \int \frac{1}{\cosh x} \cdot \frac{\sinh x}{\cosh x} dx = \int \operatorname{sech} x \tanh x dx = -\operatorname{sech} x + C$
- f $\int \frac{3}{\sinh x \tanh x} dx = 3 \int \operatorname{cosech} x \operatorname{coth} x dx = -3 \operatorname{cosech} x + C$
- g $\int \operatorname{sech} x (\operatorname{sech} x + \tanh x) dx = \int (\operatorname{sech}^2 x + \operatorname{sech} x \tanh x) dx = \tanh x - \operatorname{sech} x + C$
- h $\int (\operatorname{sech}^2 x - \operatorname{cosech}^2 x) dx = \tanh x + \operatorname{coth} x + C$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise A, Question 2

Question:

Find

a $\int \sinh 2x \, dx$

b $\int \cosh\left(\frac{x}{3}\right) dx$

c $\int \operatorname{sech}^2(2x-1) dx$

d $\int \operatorname{cosech}^2 5x \, dx$

e $\int \operatorname{cosech} 2x \coth 2x \, dx$

f $\int \operatorname{sech}\left(\frac{x}{\sqrt{2}}\right) \tanh\left(\frac{x}{\sqrt{2}}\right) dx$

g $\int \left(5 \sinh 5x - 4 \cosh 4x + 3 \operatorname{sech}^2\left(\frac{x}{2}\right) \right) dx$

Solution:

a $\int \sinh 2x \, dx = \frac{1}{2} \cosh 2x + C$

b $\int \cosh\left(\frac{x}{3}\right) dx = \frac{1}{\left(\frac{1}{3}\right)} \sinh\left(\frac{x}{3}\right) + C = 3 \sinh\left(\frac{x}{3}\right) + C$

c $\int \operatorname{sech}^2(2x-1) dx = \frac{1}{2} \tanh(2x-1) + C$

d $\int \operatorname{cosech}^2 5x dx = -\frac{1}{5} \coth 5x + C$

e $\int \operatorname{cosech} 2x \coth 2x \, dx = -\frac{1}{2} \operatorname{cosech} 2x + C$

f $\int \operatorname{sech}\left(\frac{x}{\sqrt{2}}\right) \tanh\left(\frac{x}{\sqrt{2}}\right) dx = -\frac{1}{\left(\frac{1}{\sqrt{2}}\right)} \operatorname{sech}\left(\frac{x}{\sqrt{2}}\right) + C = \sqrt{2} \operatorname{sech}\left(\frac{x}{\sqrt{2}}\right) + C$

g $\int 5 \sinh 5x - 4 \cosh 4x + 3 \operatorname{sech}^2\left(\frac{x}{2}\right) dx = 5 \times \frac{1}{5} \cosh 5x - 4 \times \frac{1}{4} \sinh 4x + 3 \times \frac{1}{\left(\frac{1}{2}\right)} \tanh\left(\frac{x}{2}\right) + C$
 $= \cosh 5x - \sinh 4x + 6 \tanh\left(\frac{x}{2}\right) + C$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise A, Question 3

Question:

Write down the results of the following. (This is a recognition exercise and also involves some integrals from C4.)

a $\int \frac{1}{1+x^2} dx$

b $\int \frac{1}{\sqrt{1+x^2}} dx$

c $\int \frac{1}{1+x} dx$

d $\int \frac{2x}{1+x^2} dx$

e $\int \frac{1}{\sqrt{1-x^2}} dx, |x| < 1$

f $\int \frac{1}{\sqrt{x^2-1}} dx$

g $\int \frac{3x}{\sqrt{x^2-1}} dx$

h $\int \frac{3}{(1+x)^2} dx$

Solution:

a $\arctan x + C$

b $\operatorname{arsinh} x + C$

c $\ln |1+x| + C$

d $\ln(1+x^2) + C$

e $\arcsin x + C$

f $\operatorname{arcosh} x + C$

g $3\sqrt{x^2-1} + C$

h $-\frac{3}{1+x} + C$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise A, Question 4

Question:

Find

a $\int \frac{2x+1}{\sqrt{1-x^2}} dx$

b $\int \frac{1+x}{\sqrt{x^2-1}} dx$

c $\int \frac{x-3}{1+x^2} dx$

Solution:

$$\begin{aligned} \text{a } \int \frac{2x+1}{\sqrt{1-x^2}} dx &= \int \frac{2x}{\sqrt{1-x^2}} dx + \int \frac{1}{\sqrt{1-x^2}} dx \\ &= 2 \int x(1-x^2)^{-\frac{1}{2}} dx + \int \frac{1}{\sqrt{1-x^2}} dx \\ &= -2\sqrt{1-x^2} + \arcsin x + C \end{aligned}$$

$$\begin{aligned} \text{b } \int \frac{1+x}{\sqrt{x^2-1}} dx &= \int \frac{1}{\sqrt{x^2-1}} dx + \int \frac{x}{\sqrt{x^2-1}} dx \\ &= \int \frac{1}{\sqrt{x^2-1}} dx + \int x(x^2-1)^{-\frac{1}{2}} dx \\ &= \operatorname{arcosh} x + \sqrt{x^2-1} + C \end{aligned}$$

$$\begin{aligned} \text{c } \int \frac{x-3}{\sqrt{1+x^2}} dx &= \int \frac{x}{\sqrt{1+x^2}} dx - \int \frac{3}{\sqrt{1+x^2}} dx \\ &= \int x(1+x^2)^{-\frac{1}{2}} dx - \int \frac{3}{\sqrt{1+x^2}} dx \\ &= \sqrt{1+x^2} - 3\operatorname{arsinh} x + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise A, Question 5

Question:

- a Show that $\frac{x^2}{1+x^2} = 1 - \frac{1}{1+x^2}$
- b Hence find $\int \frac{x^2}{1+x^2} dx$

Solution:

$$\text{a } \frac{x^2}{1+x^2} = \frac{1+x^2-1}{1+x^2} = \frac{1+x^2}{1+x^2} - \frac{1}{1+x^2} = 1 - \frac{1}{1+x^2}$$

$$\text{b } \int \frac{x^2}{1+x^2} dx = \int \left\{ 1 - \frac{1}{1+x^2} \right\} dx \quad \leftarrow \text{Using a.}$$
$$= x - \arctan x + C$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 1

Question:

Find

a $\int \sinh^3 x \cosh x \, dx$

b $\int \tanh 4x \, dx$

c $\int \tanh^5 x \operatorname{sech}^2 x \, dx$

d $\int \operatorname{cosech}^7 x \coth x \, dx$

e $\int \sqrt{\cosh 2x} \sinh 2x \, dx$

f $\int \operatorname{sech}^{10} 3x \tanh 3x \, dx$

Solution:

$$\text{a } \int \sinh^3 x \cosh x \, dx = \int (\sinh x)^3 \cosh x \, dx = \frac{1}{4} \sinh^4 x + C$$

$$\text{b } \int \tanh 4x \, dx = \int \frac{\sinh 4x}{\cosh 4x} \, dx = \frac{1}{4} \ln \cosh 4x + C$$

$$\text{c } \int \tanh^5 x \operatorname{sech}^2 x \, dx = \int (\tanh x)^5 \operatorname{sech}^2 x \, dx = \frac{1}{6} \tanh^6 x + C$$

$$\begin{aligned} \text{d } \int \operatorname{cosech}^7 x \coth x \, dx &= \int \operatorname{cosech}^6 x (\operatorname{cosech} x \coth x) \, dx \\ &= - \int (\operatorname{cosech} x)^6 (-\operatorname{cosech} x \coth x) \, dx \\ &= -\frac{1}{7} \operatorname{cosech}^7 x + C \end{aligned}$$

$$\begin{aligned} \text{e } \int \sqrt{\cosh 2x} \sinh 2x \, dx &= \frac{1}{2} \int (\cosh 2x)^{\frac{1}{2}} (2 \sinh 2x) \, dx \\ &= \frac{1}{2} \left[\frac{(\cosh 2x)^{\frac{3}{2}}}{\left(\frac{3}{2}\right)} \right] + C \\ &= \frac{1}{3} (\cosh 2x)^{\frac{3}{2}} + C \end{aligned}$$

$$\begin{aligned} \text{f } \int \operatorname{sech}^{10} 3x \tanh 3x \, dx &= -\frac{1}{3} \int \operatorname{sech}^9 3x (-3 \operatorname{sech} 3x \tanh 3x) \, dx \\ &= -\frac{1}{3} \left[\frac{\operatorname{sech}^{10} 3x}{10} \right] + C \\ &= -\frac{1}{30} \operatorname{sech}^{10} 3x + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 2

Question:

Find

$$\text{a } \int \frac{\sinh x}{2+3 \cosh x} dx$$

$$\text{b } \int \frac{1+\tanh x}{\cosh^2 x} dx$$

$$\text{c } \int \frac{5 \cosh x + 2 \sinh x}{\cosh x} dx.$$

Solution:

$$\begin{aligned} \text{a } \int \frac{\sinh x}{2+3 \cosh x} dx &= \frac{1}{3} \int \frac{3 \sinh x}{2+3 \cosh x} dx \\ &= \frac{1}{3} \ln(2+3 \cosh x) + C \end{aligned}$$

$$\begin{aligned} \text{b } \int \frac{1+\tanh x}{\cosh^2 x} dx &= \int (1+\tanh x) \operatorname{sech}^2 x dx \\ &= \int (\operatorname{sech}^2 x + \tanh x \operatorname{sech}^2 x) dx \\ &= \tanh x + \frac{1}{2} \tanh^2 x + C \quad \text{or} \quad \tanh x - \frac{1}{2} \operatorname{sech}^2 x + C \end{aligned}$$

$$\begin{aligned} \text{c } \int \frac{5 \cosh x + 2 \sinh x}{\cosh x} dx &= \int (5 + 2 \tanh x) dx \\ &= 5x + 2 \ln \cosh x + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 3

Question:

- a Show that $\int \coth x \, dx = \ln \sinh x + C$.
- b Show that $\int_1^2 \coth 2x \, dx = \ln \sqrt{\left(e^2 + \frac{1}{e^2}\right)}$.

Solution:

$$\text{a } \int \coth x \, dx = \int \frac{\cosh x}{\sinh x} \, dx = \ln \sinh x + C$$

$$\text{b } \int \coth 2x \, dx = \frac{1}{2} \ln \sinh 2x + C$$

$$\text{So } \int_1^2 \coth 2x \, dx = \left[\frac{1}{2} \ln \sinh 2x \right]_1^2 = \frac{1}{2} (\ln \sinh 4 - \ln \sinh 2)$$

$$= \frac{1}{2} \ln \left(\frac{\sinh 4}{\sinh 2} \right)$$

$$= \frac{1}{2} \ln \left(\frac{e^4 - e^{-4}}{e^2 - e^{-2}} \right)$$

$$= \frac{1}{2} \ln(e^2 + e^{-2})$$

$$= \ln \sqrt{e^2 + \frac{1}{e^2}}$$

← Using $a^2 - b^2 = (a+b)(a-b)$
with $a = e^2, b = e^{-2}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 4

Question:

Use integration by parts to find

a $\int x \sinh 3x \, dx$

b $\int x \operatorname{sech}^2 x \, dx$.

Solution:

$$\begin{aligned} \text{a } \int x \sinh 3x \, dx &= \frac{1}{3} x \cosh 3x - \int \frac{1}{3} \cosh 3x \, dx \\ &= \frac{1}{3} x \cosh 3x - \frac{1}{9} \sinh 3x + C \end{aligned}$$

Using $\int u \frac{dv}{dx} \, dx = uv - \int v \frac{du}{dx} \, dx$ with
 $u = x$ and $\frac{dv}{dx} = \sinh 3x$

$$\begin{aligned} \text{b } \int x \operatorname{sech}^2 x \, dx &= x \tanh x - \int \tanh x \, dx \\ &= x \tanh x - \ln \cosh x + C \end{aligned}$$

Using integration by parts with
 $u = x$ and $\frac{dv}{dx} = \operatorname{sech}^2 x$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 5

Question:

Find

a $\int e^x \cosh x \, dx$

b $\int e^{-2x} \sinh 3x \, dx$

c $\int \cosh x \cosh 3x \, dx$.

Solution:

$$\begin{aligned} \text{a } \int e^x \cosh x \, dx &= \int e^x \left(\frac{e^x + e^{-x}}{2} \right) dx \\ &= \frac{1}{2} \int (e^{2x} + 1) dx \\ &= \frac{1}{4} e^{2x} + \frac{1}{2} x + C \end{aligned}$$

← You cannot use integration by parts.

$$\begin{aligned} \text{b } \int e^{-2x} \sinh 3x \, dx &= \int e^{-2x} \left(\frac{e^{3x} - e^{-3x}}{2} \right) dx \\ &= \frac{1}{2} \int (e^x - e^{-5x}) dx \\ &= \frac{1}{2} e^x + \frac{1}{10} e^{-5x} + C \end{aligned}$$

← You could use integration by parts twice.

$$\begin{aligned} \text{c } \int \cosh x \cosh 3x \, dx &= \int \left(\frac{e^x + e^{-x}}{2} \right) \left(\frac{e^{3x} + e^{-3x}}{2} \right) dx \\ &\quad \text{or write as } \frac{1}{2} (\cosh 4x + \cosh 2x) \\ &= \frac{1}{4} \int (e^{4x} + e^{-4x} + e^{2x} + e^{-2x}) dx \\ &= \frac{1}{16} e^{4x} - \frac{1}{16} e^{-4x} + \frac{1}{8} e^{2x} - \frac{1}{8} e^{-2x} + C \quad \text{or} \quad \frac{1}{8} \sinh 4x + \frac{1}{4} \sinh 2x + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 6

Question:

By writing $\cosh 3x$ in exponential form, find $\int \cosh^2 3x \, dx$ and show that it is equivalent to the result found in Example 5b.

Solution:

$$\begin{aligned}\int \cosh^2 3x \, dx &= \frac{1}{4} \int (e^{3x} + e^{-3x})^2 \, dx \\ &= \frac{1}{4} \int (e^{6x} + 2 + e^{-6x}) \, dx \\ &= \frac{1}{24} e^{6x} - \frac{1}{24} e^{-6x} + \frac{1}{2} x + C \\ &= \frac{1}{12} \sinh 6x + \frac{1}{2} x + C \quad \text{which was result in Example 5b}\end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 7

Question:

Evaluate $\int_0^1 \frac{1}{\sinh x + \cosh x} dx$, giving your answer in terms of e .

Solution:

$$\sinh x + \cosh x = \frac{1}{2}(e^x - e^{-x}) + \frac{1}{2}(e^x + e^{-x}) = e^x$$

$$\text{So } \int_0^1 \left(\frac{1}{\sinh x + \cosh x} \right) dx = \int_0^1 e^{-x} dx = [-e^{-x}]_0^1 = 1 - \frac{1}{e}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 8

Question:

Use appropriate identities to find

a $\int \sinh^2 x \, dx$

b $\int (\operatorname{sech} x - \tanh x)^2 \, dx$

c $\int \frac{\cosh^2 3x}{\sinh^2 3x} \, dx$

d $\int \sinh^2 x \cosh^2 x \, dx$

e $\int \cosh^5 x \, dx$

f $\int \tanh^3 2x \, dx.$

Solution:

$$\text{a } \int \sinh^2 x \, dx = \frac{1}{2} \int (\cosh 2x - 1) \, dx = \frac{1}{4} \sinh 2x - \frac{1}{2} x + C$$

$$\begin{aligned} \text{b } \int (\operatorname{sech} x - \tanh x)^2 \, dx &= \int (\operatorname{sech}^2 x - 2\operatorname{sech} x \tanh x + \tanh^2 x) \, dx \\ &= \int (\operatorname{sech}^2 x - 2\operatorname{sech} x \tanh x + 1 - \operatorname{sech}^2 x) \, dx \\ &= \int (1 - 2\operatorname{sech} x \tanh x) \, dx \\ &= x + 2\operatorname{sech} x + C \end{aligned}$$

$$\begin{aligned} \text{c } \int \frac{\cosh^2 3x}{\sinh^2 3x} \, dx &= \int \coth^2 3x \, dx \\ &= \int (1 + \operatorname{cosech}^2 3x) \, dx \\ &= x - \frac{1}{3} \coth 3x + C \end{aligned}$$

$$\begin{aligned} \text{d } \int \sinh^2 x \cosh^2 x \, dx &= \int \left(\frac{1}{2} \sinh 2x \right)^2 \, dx \\ &= \frac{1}{4} \int \sinh^2 2x \, dx \\ &= \frac{1}{4} \int \left(\frac{\cosh 4x - 1}{2} \right) \, dx \quad \leftarrow \text{Using } \cosh 2u = 1 + 2\sinh^2 u \\ &= -\frac{1}{8} x + \frac{1}{32} \sinh 4x + C \end{aligned}$$

$$\begin{aligned} \text{e } \int \cosh^5 x \, dx &= \int \cosh^4 x \cosh x \, dx \\ &= \int (1 + \sinh^2 x)^2 \cosh x \, dx \\ &= \int (1 + 2\sinh^2 x + \sinh^4 x) \cosh x \, dx \\ &= \int (\cosh x + 2\sinh^2 x \cosh x + \sinh^4 x \cosh x) \, dx \\ &= \sinh x + \frac{2}{3} \sinh^3 x + \frac{1}{5} \sinh^5 x + C \end{aligned}$$

$$\begin{aligned} \text{f } \int \tanh^3 2x \, dx &= \int \tanh^2 2x \tanh 2x \, dx \\ &= \int (1 - \operatorname{sech}^2 2x) \tanh 2x \, dx \\ &= \int (\tanh 2x - \tanh 2x \operatorname{sech}^2 2x) \, dx \\ &= \frac{1}{2} \ln \cosh 2x - \frac{1}{4} \tanh^2 2x + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 9

Question:

Show that $\int_0^{\ln 2} \cosh^2\left(\frac{x}{2}\right) dx = \frac{1}{8}(3 + \ln 16)$.

Solution:

$$\begin{aligned} \int_0^{\ln 2} \cosh^2\left(\frac{x}{2}\right) dx &= \int_0^{\ln 2} \left(\frac{1 + \cosh x}{2}\right) dx \\ &= \frac{1}{2} [x + \sinh x]_0^{\ln 2} \\ &= \frac{1}{2} \left[\ln 2 + \left(\frac{e^{\ln 2} - e^{-\ln 2}}{2}\right) \right] \leftarrow \boxed{e^{\ln 2} = 2, e^{-\ln 2} = e^{\frac{\ln 1}{2}} = \frac{1}{2}} \\ &= \frac{1}{2} \left[\ln 2 + \frac{3}{4} \right] \\ &= \frac{1}{8} [3 + 4 \ln 2] \\ &= \frac{1}{8} (3 + \ln 16) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 10

Question:

The region bounded by the curve $y = \sinh x$, the line $x = 1$ and the positive x -axis is rotated through 360° about the x -axis. Show that the volume of the solid of revolution formed is $\frac{\pi}{8e^2}(e^4 - 4e^2 - 1)$.

Solution:

$$\begin{aligned}\text{Volume} &= \pi \int_0^1 \sinh^2 x \, dx = \frac{\pi}{2} \int_0^1 (\cosh 2x - 1) dx \\ &= \frac{\pi}{2} \left[\frac{1}{2} \sinh 2x - x \right]_0^1 \\ &= \frac{\pi}{2} \left[\frac{1}{2} \sinh 2 - 1 \right] \\ &= \frac{\pi}{2} \left[\frac{1}{4} (e^2 - e^{-2}) - 1 \right] \\ &= \frac{\pi}{8} [e^2 - 4 - e^{-2}] \\ &= \frac{\pi}{8e^2} (e^4 - 4e^2 - 1).\end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 11

Question:

Using the result for $\int \operatorname{sech} x \, dx$ given in Example 7, find

a $\int \frac{2}{\cosh x} \, dx$

b $\int \operatorname{sech} 2x \, dx$

c $\int \sqrt{1 - \tanh^2\left(\frac{x}{2}\right)} \, dx$.

Solution:

Using $\int \operatorname{sech} x \, dx = 2 \arctan(e^x) + C$

a $\int \frac{2}{\cosh x} \, dx = \int 2 \operatorname{sech} x \, dx = 4 \arctan(e^x) + C$

b Using the substitution $u = 2x$,

$\left(\text{or using } \int f(ax+b) \, dx = \frac{1}{a} f(ax+b) + C \right)$

$$\int \operatorname{sech} 2x \, dx = \frac{1}{2} \int \operatorname{sech} u \, du = \arctan(e^u) + C = \arctan(e^{2x}) + C$$

c $\int \sqrt{1 - \tanh^2\left(\frac{x}{2}\right)} \, dx = \int \operatorname{sech}\left(\frac{x}{2}\right) \, dx = \frac{1}{\left(\frac{1}{2}\right)} 2 \arctan\left(e^{\frac{x}{2}}\right) + C$
 $= 4 \arctan\left(e^{\frac{x}{2}}\right) + C$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise B, Question 12

Question:

Using the substitution $u = x^2$, or otherwise, find

a $\int x \cosh^2(x^2) dx$

b $\int \frac{x}{\cosh^2(x^2)} dx$.

Solution:

Using the substitution $u = x^2$, $du = 2x dx$,

a So $\int x \cosh^2(x^2) dx = \frac{1}{2} \int \cosh^2 u du$

$$= \frac{1}{4} \int (\cosh 2u + 1) du$$

$$= \frac{1}{8} \sinh 2u + \frac{u}{4} + C$$

$$= \frac{1}{8} \sinh(2x^2) + \frac{x^2}{4} + C$$

b So $\int \frac{x}{\cosh^2(x^2)} dx = \int x \operatorname{sech}^2(x^2) dx$

$$= \frac{1}{2} \int \operatorname{sech}^2 u du$$

$$= \frac{1}{2} \tanh u + C$$

$$= \frac{1}{2} \tanh(x^2) + C$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 1

Question:

Use the substitution $x = a \tan \theta$ to show that $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C$.

Solution:

Using $x = a \tan \theta$, $dx = a \sec^2 \theta d\theta$

$$\begin{aligned} \text{so } \int \frac{1}{a^2 + x^2} dx &= \int \frac{1}{a^2 + a^2 \tan^2 \theta} a \sec^2 \theta d\theta \\ &= \int \frac{a \sec^2 \theta}{a^2 \sec^2 \theta} d\theta \\ &= \frac{1}{a} \int d\theta \\ &= \frac{1}{a} \theta + C \\ &= \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C \end{aligned}$$

$$\left. \begin{array}{l} \leftarrow \\ x = a \tan \theta \Rightarrow \theta = \arctan\left(\frac{x}{a}\right) \end{array} \right\}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 2

Question:

Use the substitution $x = \cos \theta$ to show that $\int \frac{1}{\sqrt{1-x^2}} dx = -\arccos x + C$.

Solution:

Using $x = \cos \theta$, $dx = -\sin \theta d\theta$

$$\begin{aligned} \text{so } \int \frac{1}{\sqrt{1-x^2}} dx &= \int \frac{1}{\sqrt{1-\cos^2 \theta}} (-\sin \theta) d\theta \\ &= -\int d\theta \\ &= -\theta + C \\ &= -\arccos x + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 3

Question:

Use suitable substitutions to find

a $\int \frac{3}{\sqrt{4-x^2}} dx$

b $\int \frac{1}{\sqrt{x^2-9}} dx$

c $\int \frac{4}{5+x^2} dx$

d $\int \frac{1}{\sqrt{4x^2+25}} dx$.

Solution:

a Let $x = 2 \sin \theta$, so $dx = 2 \cos \theta d\theta$

$$\begin{aligned} \int \frac{3}{\sqrt{4-x^2}} dx &= \int \frac{3}{\sqrt{4-4\sin^2\theta}} 2 \cos \theta d\theta \\ &= \int \frac{6 \cos \theta}{2 \cos \theta} d\theta \\ &= 3 \int d\theta \\ &= 3\theta + C \\ &= 3 \arcsin\left(\frac{x}{2}\right) + C \end{aligned}$$

b Let $x = 3 \cosh u$, so $dx = 3 \sinh u du$

$$\begin{aligned} \int \frac{1}{\sqrt{x^2-9}} dx &= \int \frac{1}{\sqrt{9 \cosh^2 u - 9}} 3 \sinh u du \\ &= \int \frac{1}{3 \sqrt{\cosh^2 u - 1}} 3 \sinh u du \\ &= \int \frac{3 \sinh u}{3 \sinh u} du \\ &= \int 1 du \\ &= u + C \\ &= \operatorname{arcosh}\left(\frac{x}{3}\right) + C \end{aligned}$$

c Let $x = \sqrt{5} \tan \theta$, so $dx = \sqrt{5} \sec^2 \theta d\theta$

$$\begin{aligned} \int \frac{4}{5+x^2} dx &= \int \frac{4}{5+5 \tan^2 \theta} \sqrt{5} \sec^2 \theta d\theta \\ &= \int \frac{4\sqrt{5} \sec^2 \theta}{5 \sec^2 \theta} d\theta \\ &= \frac{4\sqrt{5}}{5} \int d\theta \\ &= \frac{4\sqrt{5}}{5} \theta + C \\ &= \frac{4\sqrt{5}}{5} \arctan\left(\frac{x}{\sqrt{5}}\right) + C \end{aligned}$$

$5 + 5 \tan^2 \theta = 5(1 + \tan^2 \theta) = 5 \sec^2 \theta$

d You need $4x^2 = 25 \sinh^2 u$, or $2x = 5 \sinh u$, then $dx = \frac{5}{2} \cosh u \, du$

$$\begin{aligned} \int \frac{1}{\sqrt{4x^2 + 25}} \, dx &= \int \frac{1}{\sqrt{25 \sinh^2 u + 25}} \left(\frac{5}{2} \cosh u \right) du \\ &= \frac{5}{2} \int \frac{\cosh u}{5 \sqrt{\sinh^2 u + 1}} \, du \\ &= \frac{1}{2} \int \frac{\cosh u}{\cosh u} \, du \\ &= \frac{1}{2} \int 1 \, du \\ &= \frac{1}{2} u + C \\ &= \frac{1}{2} \operatorname{arsinh} \left(\frac{2x}{5} \right) + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 4

Question:

Write down the results for the following:

a $\int \frac{1}{\sqrt{25-x^2}} dx$

b $\int \frac{3}{\sqrt{x^2+9}} dx$

c $\int \frac{1}{\sqrt{x^2-2}} dx$

d $\int \frac{2}{16+x^2} dx$.

Solution:

a $\int \frac{1}{\sqrt{25-x^2}} dx = \arcsin\left(\frac{x}{5}\right) + C$ ← Using $\int \frac{1}{\sqrt{a^2-x^2}} dx = \arcsin\left(\frac{x}{a}\right) + C$

b $\int \frac{3}{\sqrt{x^2+9}} dx = 3\operatorname{arsinh}\left(\frac{x}{3}\right) + C$ ← Using $\int \frac{1}{\sqrt{x^2+a^2}} dx = \operatorname{arsinh}\left(\frac{x}{a}\right) + C$.

c $\int \frac{1}{\sqrt{x^2-2}} dx = \operatorname{arcosh}\left(\frac{x}{\sqrt{2}}\right) + C$ ← Using $\int \frac{1}{\sqrt{x^2-a^2}} dx = \operatorname{arcosh}\left(\frac{x}{a}\right) + C$

d $\int \frac{2}{16+x^2} dx = 2 \int \frac{1}{16+x^2} dx$
 $= 2 \left\{ \frac{1}{4} \arctan\left(\frac{x}{4}\right) \right\} + C$ ← Using $\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C$
 $= \frac{1}{2} \arctan\left(\frac{x}{4}\right) + C$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 5

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

Find

a $\int \frac{1}{\sqrt{4x^2 - 12}} dx$

b $\int \frac{1}{4 + 3x^2} dx$

c $\int \frac{1}{\sqrt{9x^2 + 16}} dx$

d $\int \frac{1}{\sqrt{3 - 4x^2}} dx$.

Solution:

$$\begin{aligned}
 \text{a } \int \frac{1}{\sqrt{4x^2 - 12}} dx &= \int \frac{1}{\sqrt{4(x^2 - 3)}} dx \\
 &= \frac{1}{2} \int \frac{1}{\sqrt{(x^2 - 3)}} dx \\
 &= \frac{1}{2} \operatorname{arcosh} \left(\frac{x}{\sqrt{3}} \right) + C
 \end{aligned}$$

$$\begin{aligned}
 \text{b } \int \frac{1}{4 + 3x^2} dx &= \int \frac{1}{3 \left\{ \frac{4}{3} + x^2 \right\}} dx \\
 &= \frac{1}{3} \left[\frac{1}{\left(\frac{2}{\sqrt{3}} \right)} \arctan \left(\frac{x}{\left(\frac{2}{\sqrt{3}} \right)} \right) \right] + C \\
 &= \frac{\sqrt{3}}{6} \arctan \left(\frac{\sqrt{3}x}{2} \right) + C
 \end{aligned}$$

$$\begin{aligned}
 \text{c } \int \frac{1}{\sqrt{9x^2 + 16}} dx &= \int \frac{1}{\sqrt{9 \left\{ x^2 + \left(\frac{16}{9} \right) \right\}}} dx \\
 &= \frac{1}{3} \int \frac{1}{\sqrt{\left\{ x^2 + \left(\frac{16}{9} \right) \right\}}} dx \\
 &= \frac{1}{3} \operatorname{arsinh} \left(\frac{x}{\left(\frac{4}{3} \right)} \right) + C \\
 &= \frac{1}{3} \operatorname{arsinh} \left(\frac{3x}{4} \right) + C
 \end{aligned}$$

$$\begin{aligned}
 \text{d } \int \frac{1}{\sqrt{3 - 4x^2}} dx &= \int \frac{1}{\sqrt{4 \left\{ \frac{3}{4} - x^2 \right\}}} dx \\
 &= \frac{1}{2} \int \frac{1}{\sqrt{\left\{ \frac{3}{4} - x^2 \right\}}} dx \\
 &= \frac{1}{2} \arcsin \left(\frac{x}{\left(\frac{\sqrt{3}}{2} \right)} \right) + C \\
 &= \frac{1}{2} \arcsin \left(\frac{2x}{\sqrt{3}} \right) + C
 \end{aligned}$$

$a^2 = \frac{3}{4} \text{ so } a = \frac{\sqrt{3}}{2}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 6

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

Evaluate

$$\text{a } \int_1^3 \frac{2}{1+x^2} dx$$

$$\text{b } \int_1^2 \frac{3}{\sqrt{1+4x^2}} dx$$

$$\text{c } \int_{-1}^2 \frac{1}{\sqrt{21-3x^2}} dx.$$

Solution:

$$\begin{aligned} \text{a } \int_1^3 \frac{2}{1+x^2} dx &= 2[\arctan x]_1^3 \\ &= 2(\arctan 3 - \arctan 1) \\ &= 0.927 \quad (3 \text{ s.f.}) \end{aligned}$$

Remember that you need to be in radian mode.

$$\begin{aligned} \text{b } \int_1^2 \frac{3}{\sqrt{1+4x^2}} dx &= 3 \int_1^2 \frac{1}{2\sqrt{\frac{1}{4}+x^2}} dx \\ &= \frac{3}{2} \left[\operatorname{arsinh} \frac{x}{\left(\frac{1}{2}\right)} \right]_1^2 \\ &= \frac{3}{2} [\operatorname{arsinh}(2x)]_1^2 \\ &= \frac{3}{2} [\operatorname{arsinh} 4 - \operatorname{arsinh} 2] \\ &= 0.977 \quad (3 \text{ s.f.}) \end{aligned}$$

$$\begin{aligned} \text{c } \int_{-1}^2 \frac{1}{\sqrt{21-3x^2}} dx &= \frac{1}{\sqrt{3}} \int_{-1}^2 \frac{1}{\sqrt{7-x^2}} dx \\ &= \frac{1}{\sqrt{3}} \left[\arcsin \left(\frac{x}{\sqrt{7}} \right) \right]_{-1}^2 \\ &= \frac{1}{\sqrt{3}} \left[\arcsin \left(\frac{2}{\sqrt{7}} \right) - \arcsin \left(-\frac{1}{\sqrt{7}} \right) \right] \\ &= \frac{1}{\sqrt{3}} [0.85707\dots - (-0.38759\dots)] \\ &= 0.719 \quad (3 \text{ s.f.}) \end{aligned}$$

You need to be in radian mode

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 7

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

Evaluate, giving your answers in terms of π or as a single natural logarithm, whichever is appropriate.

a $\int_0^4 \frac{1}{\sqrt{x^2 + 16}} dx$

b $\int_{13}^{15} \frac{1}{\sqrt{x^2 - 144}} dx$

c $\int_{\sqrt{2}}^{\sqrt{6}} \frac{1}{\sqrt{4 - x^2}} dx$

Solution:

Reminder: The logarithmic form of an inverse hyperbolic function is in the Edexcel formulae booklet.

$$\begin{aligned} \text{a } \int_0^4 \frac{1}{\sqrt{x^2+16}} dx &= \left[\operatorname{arsinh} \left(\frac{x}{4} \right) \right]_0^4 \\ &= \operatorname{arsinh} 1 - \operatorname{arsinh} 0 \\ &= \ln \{1 + \sqrt{2}\} \end{aligned}$$

Using $\operatorname{arsinh} x = \ln \{x + \sqrt{x^2 + 1}\}$

$$\begin{aligned} \text{b } \int_{13}^{15} \frac{1}{\sqrt{x^2-144}} dx &= \left[\operatorname{arcosh} \left(\frac{x}{12} \right) \right]_{13}^{15} \\ &= \operatorname{arcosh} \left(\frac{5}{4} \right) - \operatorname{arcosh} \left(\frac{13}{12} \right) \\ &= \ln \left\{ \frac{5}{4} + \sqrt{\frac{25}{16} - 1} \right\} - \ln \left\{ \frac{13}{12} + \sqrt{\frac{169}{144} - 1} \right\} \\ &= \ln \left\{ \frac{5}{4} + \sqrt{\frac{9}{16}} \right\} - \ln \left\{ \frac{13}{12} + \sqrt{\frac{25}{144}} \right\} \\ &= \ln 2 - \ln \left(\frac{3}{2} \right) \\ &= \ln \left(\frac{4}{3} \right) \end{aligned}$$

Using $\operatorname{arcosh} x = \ln \{x + \sqrt{x^2 - 1}\}$

Using $\ln a - \ln b = \ln \left(\frac{a}{b} \right)$

$$\begin{aligned} \text{c } \int_{\sqrt{2}}^{\sqrt{3}} \frac{1}{\sqrt{4-x^2}} dx &= \left[\arcsin \left(\frac{x}{2} \right) \right]_{\sqrt{2}}^{\sqrt{3}} \\ &= \arcsin \left(\frac{\sqrt{3}}{2} \right) - \arcsin \left(\frac{\sqrt{2}}{2} \right) \\ &= \left(\frac{\pi}{3} \right) - \left(\frac{\pi}{4} \right) \\ &= \left(\frac{\pi}{12} \right) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 8

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

The curve C has equation $y = \frac{2}{\sqrt{2x^2 + 9}}$. The region R is bounded by C , the

coordinate axes and the lines $x = -1$ and $x = 3$.

a Find the area of R .

The region R is rotated through 360° about the x -axis.

b Find the volume of the solid generated.

Solution:

$$\text{a Area of } R = \int_{-1}^3 y \, dx = \int_{-1}^3 \frac{2}{\sqrt{2x^2+9}} \, dx$$

$$= \int_{-1}^3 \frac{2}{\sqrt{2\left(x^2 + \frac{9}{2}\right)}} \, dx$$

← Curve is always above x -axis

$$= \sqrt{2} \left[\operatorname{arsinh} \frac{x}{\left(\frac{3}{\sqrt{2}}\right)} \right]_{-1}^3$$

$$= \sqrt{2} \left[\operatorname{arsinh} \left(\frac{\sqrt{2}x}{3} \right) \right]_{-1}^3$$

$$= \sqrt{2} \left[\operatorname{arsinh} \sqrt{2} - \operatorname{arsinh} \left(-\frac{\sqrt{2}}{3} \right) \right]$$

$$= 2.27 \text{ (3 s.f.)}$$

$$\text{b Volume} = \pi \int_{-1}^3 y^2 \, dx = \pi \int_{-1}^3 \frac{4}{2x^2+9} \, dx$$

$$= 2\pi \int_{-1}^3 \frac{1}{x^2 + \left(\frac{3}{2}\right)} \, dx$$

$$= 2\pi \left[\frac{1}{\left(\frac{3}{\sqrt{2}}\right)} \arctan \frac{x}{\left(\frac{3}{\sqrt{2}}\right)} \right]_{-1}^3$$

$$= \left(\frac{2\sqrt{2}\pi}{3} \right) \left[\arctan \left(\frac{\sqrt{2}x}{3} \right) \right]_{-1}^3$$

$$= \left(\frac{2\sqrt{2}\pi}{3} \right) \left[\arctan(\sqrt{2}) - \arctan \left(-\frac{\sqrt{2}}{3} \right) \right]$$

$$= 1.32\pi \text{ (3 s.f.)} = 4.13 \text{ (3 s.f.)}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 9

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

A circle C has centre the origin and radius r .

- a Show that the area of C can be written as $4 \int_0^r \sqrt{r^2 - x^2} dx$.
- b Hence show that the area of C is πr^2 .


Solution:

- a Cartesian equation of circle is $x^2 + y^2 = r^2$.

Area of C can be written as $4 \int_0^r y dx = 4 \int_0^r \sqrt{r^2 - x^2} dx$

- b Use substitution $x = r \sin \theta$, so $dx = r \cos \theta d\theta$,

$$\begin{aligned}
 4 \int_0^r \sqrt{r^2 - x^2} dx &= 4 \int_0^{\frac{\pi}{2}} \sqrt{r^2 - r^2 \sin^2 \theta} r \cos \theta d\theta \\
 &= 4r^2 \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta \\
 &= 2r^2 \int_0^{\frac{\pi}{2}} (1 + \cos 2\theta) d\theta \\
 &= 2r^2 \left[\theta + \frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{2}} \\
 &= 2r^2 \left(\frac{\pi}{2} \right) \\
 &= \pi r^2
 \end{aligned}$$



Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 10

Question:

- a Use the substitution $x = \frac{2}{3} \tan \theta$ to find $\int \frac{x^2}{9x^2 + 4} dx$.
- b Use the substitution $x = \sinh^2 u$ to find $\int \sqrt{\frac{x}{x+1}} dx, x > 0$.

Solution:

a With $x = \frac{2}{3} \tan \theta$ and $dx = \frac{2}{3} \sec^2 \theta d\theta$,

$$9x^2 + 4 = 9\left(\frac{4}{9} \tan^2 \theta\right) + 4 = 4 \tan^2 \theta + 4 = 4(\tan^2 \theta + 1) = 4 \sec^2 \theta$$

and $\frac{x^2}{9x^2 + 4} = \frac{\frac{4}{9} \tan^2 \theta}{4 \sec^2 \theta} = \frac{\tan^2 \theta}{9 \sec^2 \theta}$

so $\int \frac{x^2}{9x^2 + 4} dx = \int \frac{\tan^2 \theta}{9 \sec^2 \theta} \times \frac{2}{3} \sec^2 \theta d\theta$

$$= \frac{2}{27} \int \tan^2 \theta d\theta$$

$$= \frac{2}{27} \int (\sec^2 \theta - 1) d\theta$$

$$= \frac{2}{27} (\tan \theta - \theta) + C$$

$$= \frac{2}{27} \left(\frac{3x}{2} - \arctan \frac{3x}{2} \right) + C$$

$$= \frac{x}{9} - \frac{2}{27} \arctan \frac{3x}{2} + C$$

b With $x = \sinh^2 u$ and $dx = 2 \sinh u \cosh u du$,

and $\frac{x}{x+1} = \frac{\sinh^2 u}{\sinh^2 u + 1} = \frac{\sinh^2 u}{\cosh^2 u}$

$$\int \sqrt{\frac{x}{x+1}} dx = \int \frac{\sinh u}{\cosh u} 2 \sinh u \cosh u du$$

$$= \int 2 \sinh^2 u du$$

$$= \int (\cosh 2u - 1) du$$

$$= \frac{\sinh 2u}{2} - u + C$$

$$= \sinh u \cosh u - \operatorname{arsinh}(\sqrt{x}) + C$$

$$= \sqrt{x} \sqrt{1+x} - \operatorname{arsinh}(\sqrt{x}) + C$$

$\sinh u = \sqrt{x} \text{ and}$ $\cosh u = \sqrt{1 + \sinh^2 u}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 11

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

By splitting up each integral into two separate integrals, or otherwise, find

- a $\int \frac{x-2}{\sqrt{x^2-4}} dx$
 b $\int \frac{2x-1}{\sqrt{2-x^2}} dx$
 c $\int \frac{2+3x}{1+3x^2} dx$.

Solution:

$$\begin{aligned} \text{a } \int \frac{x-2}{\sqrt{x^2-4}} dx &= \int \frac{x}{\sqrt{x^2-4}} dx - \int \frac{2}{\sqrt{x^2-4}} dx \\ &= \sqrt{x^2-4} - 2 \operatorname{arcosh}\left(\frac{x}{2}\right) + C \end{aligned}$$

$$\begin{aligned} \text{b } \int \frac{2x-1}{\sqrt{2-x^2}} dx &= \int \frac{2x}{\sqrt{2-x^2}} dx - \int \frac{1}{\sqrt{2-x^2}} dx \\ &= -2\sqrt{2-x^2} - \arcsin\left(\frac{x}{\sqrt{2}}\right) + C \end{aligned}$$

$$\begin{aligned} \text{c } \int \frac{2+3x}{1+3x^2} dx &= \int \frac{2}{1+3x^2} dx + \int \frac{3x}{1+3x^2} dx \\ &= \frac{2}{3} \int \frac{1}{\left(\frac{1}{3}+x^2\right)} dx + \frac{1}{2} \int \frac{6x}{1+3x^2} dx \\ &= \frac{2\sqrt{3}}{3} \arctan(\sqrt{3}x) + \frac{1}{2} \ln(1+3x^2) + C \end{aligned}$$

$a = \frac{1}{\sqrt{3}} \text{ in } \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 12

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

Use the method of partial fractions to find $\int \frac{x^2 + 4x + 10}{x^3 + 5x} dx, x > 0$.

Solution:

Setting up the model $\frac{x^2 + 4x + 10}{x(x^2 + 5)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 5}$

$$\Rightarrow x^2 + 4x + 10 = A(x^2 + 5) + (Bx + C)x$$

$$x = 0 \Rightarrow 10 = 5A \Rightarrow A = 2$$

$$\text{Coefficient of } x \Rightarrow 4 = C$$

$$\text{Coefficient of } x^2 \Rightarrow 1 = A + B \Rightarrow B = -1$$

$$\text{So } \int \frac{x^2 + 4x + 10}{x^3 + 5x} dx = \int \left(\frac{2}{x} + \frac{-x + 4}{x^2 + 5} \right) dx$$

$$= \int \left(\frac{2}{x} + \frac{4}{x^2 + 5} - \frac{1}{2} \frac{2x}{x^2 + 5} \right) dx$$

$$= 2 \ln x + \frac{4}{\sqrt{5}} \arctan \left(\frac{x}{\sqrt{5}} \right) - \frac{1}{2} \ln(x^2 + 5) + C$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 13

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

Show that $\int_0^1 \frac{2}{(x+1)(x^2+1)} dx = \frac{1}{4}(\pi + 2\ln 2)$.

Solution:

Setting up the model $\frac{2}{(x+1)(x^2+1)} = \frac{A}{x+1} + \frac{Bx+C}{x^2+1}$

$$\Rightarrow 2 \equiv A(x^2+1) + (Bx+C)(x+1)$$

$$x = -1 \Rightarrow 2 = 2A \Rightarrow A = 1$$

Coefficient of $x^2 \Rightarrow 0 = A + B \Rightarrow B = -1$

Coefficient of $x \Rightarrow 0 = B + C \Rightarrow C = 1$

So $\int_0^1 \frac{2}{(x+1)(x^2+1)} dx = \int_0^1 \frac{1}{(x+1)} dx + \int_0^1 \frac{1-x}{(x^2+1)} dx$

$$= \int_0^1 \frac{1}{(x+1)} dx + \int_0^1 \frac{1}{(x^2+1)} dx - \int_0^1 \frac{x}{(x^2+1)} dx$$

$$= [\ln(1+x)]_0^1 + [\arctan x]_0^1 - \left[\frac{1}{2} \ln(1+x^2) \right]_0^1$$

$$= \ln 2 + \arctan 1 - \frac{1}{2} \ln 2$$

$$= \frac{\pi}{4} + \frac{1}{2} \ln 2$$

$$= \frac{1}{4}(\pi + 2\ln 2)$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 14

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

By using the substitution $u = x^2$ evaluate $\int_2^3 \frac{2x}{\sqrt{x^4-1}} dx$.

Solution:

With $u = x^2$ and $du = 2x dx$,

$$\begin{aligned}\int_2^3 \frac{2x}{\sqrt{x^4-1}} dx &= \int_4^9 \frac{du}{\sqrt{u^2-1}} \\ &= [\operatorname{ar} \cosh u]_4^9 \\ &= \operatorname{ar} \cosh 9 - \operatorname{ar} \cosh 4 \\ &= 0.824 \quad (3 \text{ s.f.})\end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 15

Question:

By using the substitution $x = \frac{1}{2} \sin \theta$, show that $\int_0^{\frac{1}{4}} \frac{x^2}{\sqrt{1-4x^2}} dx = \frac{1}{192}(2\pi - 3\sqrt{3})$.

Solution:

With $x = \frac{1}{2} \sin \theta$, $dx = \frac{1}{2} \cos \theta d\theta$

$$1 - 4x^2 = 1 - \sin^2 \theta = \cos^2 \theta \quad \text{and so} \quad \frac{x^2}{\sqrt{1-4x^2}} = \frac{\sin^2 \theta}{4 \cos \theta}$$

$$\begin{aligned} \text{So } \int_0^{\frac{1}{4}} \frac{x^2}{\sqrt{1-4x^2}} dx &= \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta}{4 \cos \theta} \times \frac{1}{2} \cos \theta d\theta \\ &= \frac{1}{8} \int_0^{\frac{\pi}{6}} \sin^2 \theta d\theta \\ &= \frac{1}{16} \int_0^{\frac{\pi}{6}} (1 - \cos 2\theta) d\theta \\ &= \frac{1}{16} \left[\theta - \frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{6}} \\ &= \frac{1}{16} \left[\frac{\pi}{6} - \frac{\sqrt{3}}{4} \right] \\ &= \frac{1}{192} (2\pi - 3\sqrt{3}) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 16

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

- a Use the substitution $x = 2 \cosh u$ to show that

$$\int \sqrt{x^2 - 4} \, dx = \frac{1}{2} x \sqrt{x^2 - 4} - 2 \operatorname{arcosh} \left(\frac{x}{2} \right) + C.$$

- b Find the area enclosed between the hyperbola with equation $\frac{x^2}{4} - \frac{y^2}{9} = 1$ and the line $x = 4$.

Solution:

a Using $x = 2 \cosh u$, $dx = 2 \sinh u \, du$


$$\begin{aligned} \int \sqrt{x^2 - 4} \, dx &= \int 2\sqrt{\cosh^2 u - 1} \times 2 \sinh u \, du \\ &= 4 \int \sinh^2 u \, du \\ &= 2 \int (\cosh 2u - 1) \, du \\ &= 2 \left\{ \frac{\sinh 2u}{2} - u \right\} + C \\ &= 2 \sinh u \cosh u - 2u + C \\ &= 2 \left(\sqrt{\left(\frac{x}{2}\right)^2 - 1} \right) \left(\frac{x}{2}\right) - 2 \operatorname{arcosh} \left(\frac{x}{2}\right) + C \\ &= 2 \left(\frac{\sqrt{x^2 - 4}}{2} \right) \left(\frac{x}{2}\right) - 2 \operatorname{arcosh} \left(\frac{x}{2}\right) + C \\ &= \frac{1}{2} x \sqrt{x^2 - 4} - 2 \operatorname{arcosh} \left(\frac{x}{2}\right) + C \end{aligned}$$

$\begin{aligned} \cosh u &= \frac{x}{2} \text{ and} \\ \sinh u &= \sqrt{\cosh^2 u - 1} \end{aligned}$

b Area = $2 \int_2^4 y \, dx$

Rearranging $\frac{x^2}{4} - \frac{y^2}{9} = 1$ gives $9x^2 - 4y^2 = 36$

$$\begin{aligned} 4y^2 &= 9x^2 - 36 \\ &= 9(x^2 - 4) \end{aligned}$$



So $y = \frac{3}{2} \sqrt{x^2 - 4}$, taking the +ve value, representing the part of curve in first quadrant

$$\begin{aligned} \text{Area} &= 3 \int_2^4 \sqrt{x^2 - 4} \, dx = \left[\frac{3}{2} x \sqrt{x^2 - 4} - 6 \operatorname{arcosh} \left(\frac{x}{2}\right) \right]_2^4 \\ &= [6\sqrt{12} - 6 \operatorname{arcosh} 2] - [0 - 6 \operatorname{arcosh} 1] \\ &= 12.9 \text{ (3 s.f.)} \end{aligned}$$

Using result from a

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 17

Question:

Unless a substitution is given or asked for, use the standard results 7 to 14. Give numerical answers to 3 significant figures, unless otherwise stated.

- a Show that $\int \frac{1}{2\cosh x - \sinh x} dx$ can be written as $\int \frac{2e^x}{e^{2x} + 3} dx$.
- b Hence, by using the substitution $u = e^x$, find $\int \frac{1}{2\cosh x - \sinh x} dx$.

Solution:

$$\text{a } 2\cosh x - \sinh x = 2\left(\frac{e^x + e^{-x}}{2}\right) - \left(\frac{e^x - e^{-x}}{2}\right) = \frac{e^x + 3e^{-x}}{2}$$

$$\begin{aligned} \text{So } \int \frac{1}{2\cosh x - \sinh x} dx &= \int \frac{2}{e^x + 3e^{-x}} dx \\ &= \int \frac{2e^x}{e^{2x} + 3} dx \end{aligned}$$

Multiplying numerator
and denominator by e^x .

- b Using the substitution $u = e^x$, $du = e^x dx$ and

$$\begin{aligned} \int \frac{2e^x}{e^{2x} + 3} dx &= 2 \int \frac{du}{u^2 + 3} \\ &= \frac{2}{\sqrt{3}} \arctan\left(\frac{u}{\sqrt{3}}\right) + C \\ &= \frac{2}{\sqrt{3}} \arctan\left(\frac{e^x}{\sqrt{3}}\right) + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 18

Question:

Using the substitution $u = \frac{2}{3} \sinh x$, evaluate $\int_0^1 \frac{\cosh x}{\sqrt{4 \sinh^2 x + 9}} dx$.

Solution:

With $u = \frac{2}{3} \sinh x$, $du = \frac{2}{3} \cosh x dx$ or $\cosh x dx = \frac{3}{2} du$

$$4 \sinh^2 x + 9 = 4 \left(\frac{3u}{2} \right)^2 + 9 = 9u^2 + 9 = 9(u^2 + 1)$$

$$\text{so } \int_0^1 \frac{\cosh x}{\sqrt{4 \sinh^2 x + 9}} dx = \int_0^{\frac{2}{3} \sinh 1} \frac{1}{3\sqrt{u^2 + 1}} \times \frac{3}{2} du$$

$$= \frac{1}{2} \operatorname{arsinh}(u) \quad \text{between the given limits}$$

$$= \frac{1}{2} \operatorname{arsinh} \left(\frac{2}{3} \sinh 1 \right)$$

$$= 0.360 \text{ (3 s.f.)}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 19

Question:

- a Find $\int \frac{dx}{a^2 - x^2}$ $|x| < a$, by using
- partial fractions,
 - the substitution $x = a \tanh \theta$.
- b Deduce the logarithmic form of $\operatorname{artanh}\left(\frac{x}{a}\right)$.

Solution:

- a i Using partial fractions $\frac{1}{a^2 - x^2} = \frac{1}{2a} \left\{ \frac{1}{a-x} + \frac{1}{a+x} \right\}$

$$\begin{aligned} \text{So } \int \frac{dx}{a^2 - x^2} &= \frac{1}{2a} \int \left\{ \frac{1}{a-x} + \frac{1}{a+x} \right\} dx \\ &= \frac{1}{2a} [-\ln|a-x| + \ln|a+x|] + C \\ &= \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right| + C \end{aligned}$$

- ii Using the substitution $x = a \tanh \theta$, $dx = a \operatorname{sech}^2 \theta d\theta$

$$\begin{aligned} \int \frac{dx}{a^2 - x^2} &= \int \frac{a \operatorname{sech}^2 \theta}{a^2 \operatorname{sech}^2 \theta} d\theta \\ &= \frac{1}{a} \theta + D \\ &= \frac{1}{a} \operatorname{artanh}\left(\frac{x}{a}\right) + D \end{aligned}$$

- b Using the result in a $\operatorname{artanh}\left(\frac{x}{a}\right) = \frac{1}{2} \ln \left| \frac{a+x}{a-x} \right| + \text{constant}$

$$\text{At } x=0, 0 = 0 + \text{constant}, \Rightarrow \text{constant} = 0 \text{ and so } \operatorname{artanh}\left(\frac{x}{a}\right) = \frac{1}{2} \ln \left| \frac{a+x}{a-x} \right|$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise C, Question 20

Question:

Using the substitution $x = \sec \theta$, find

a $\int \frac{1}{x\sqrt{x^2-1}} dx$

b $\int \frac{\sqrt{x^2-1}}{x} dx$.

Solution:

With $x = \sec \theta$,

a
$$\int \frac{1}{x\sqrt{x^2-1}} dx = \int \frac{1}{\sec \theta \sqrt{\sec^2 \theta - 1}} \sec \theta \tan \theta d\theta$$

$$= \int 1 d\theta$$

$$= \theta + C$$

$$= \operatorname{arcsec} x + C$$

b
$$\int \frac{\sqrt{x^2-1}}{x} dx = \int \frac{\sqrt{\sec^2 \theta - 1}}{\sec \theta} \sec \theta \tan \theta d\theta$$

$$= \int \tan^2 \theta d\theta$$

$$= \int (\sec^2 \theta - 1) d\theta$$

$$= \tan \theta - \theta + C$$

$$= \sqrt{\sec^2 \theta - 1} - \theta + C$$

$$= \sqrt{x^2 - 1} - \operatorname{arcsec} x + C$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 1

Question:

Find the following.

a $\int \frac{1}{\sqrt{5-4x-x^2}} dx$

b $\int \frac{1}{\sqrt{x^2-4x-12}} dx$

c $\int \frac{1}{\sqrt{x^2+6x+10}} dx$

d $\int \frac{1}{\sqrt{x(x-2)}} dx$

e $\int \frac{1}{2x^2+4x+7} dx$

f $\int \frac{1}{\sqrt{-4x^2-12x}} dx$

g $\int \frac{1}{\sqrt{14-12x-2x^2}} dx$

h $\int \frac{1}{\sqrt{9x^2-8x+1}} dx$

Solution:

$$\text{a } 5 - 4x - x^2 = -(x^2 + 4x - 5) = -\{(x+2)^2 - 9\} = 9 - (x+2)^2$$

$$\text{So } \int \frac{1}{\sqrt{5-4x-x^2}} dx = \int \frac{1}{\sqrt{9-(x+2)^2}} dx$$

Let $u = (x+2)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{5-4x-x^2}} dx &= \int \frac{1}{\sqrt{9-u^2}} du \\ &= \arcsin\left(\frac{u}{3}\right) + C \\ &= \arcsin\left(\frac{x+2}{3}\right) + C \end{aligned}$$

$$\text{b } x^2 - 4x - 12 = \{(x-2)^2 - 16\}$$

$$\text{So } \int \frac{1}{\sqrt{x^2-4x-12}} dx = \int \frac{1}{\sqrt{(x-2)^2-16}} dx$$

Let $u = (x-2)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{x^2-4x-12}} dx &= \int \frac{1}{\sqrt{u^2-16}} du \\ &= \operatorname{arcosh}\left(\frac{u}{4}\right) + C \\ &= \operatorname{arcosh}\left(\frac{x-2}{4}\right) + C \end{aligned}$$

$$\text{c } x^2 + 6x + 10 = \{(x+3)^2 + 1\}$$

$$\text{So } \int \frac{1}{\sqrt{x^2+6x+10}} dx = \int \frac{1}{\sqrt{(x+3)^2+1}} dx$$

Let $u = (x+3)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{x^2+6x+10}} dx &= \int \frac{1}{\sqrt{u^2+1}} du \\ &= \operatorname{arsinh}(u) + C \\ &= \operatorname{arsinh}(x+3) + C \end{aligned}$$

$$\text{d } x(x-2) = x^2 - 2x = \{(x-1)^2 - 1\}$$

$$\text{So } \int \frac{1}{\sqrt{x(x-2)}} dx = \int \frac{1}{\sqrt{(x-1)^2 - 1}} dx$$

$$\text{Let } u = (x-1), \text{ so } du = dx.$$

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{x(x-2)}} dx &= \int \frac{1}{\sqrt{u^2 - 1}} du \\ &= \operatorname{arcosh}(u) + C \\ &= \operatorname{arcosh}(x-1) + C \end{aligned}$$

$$\text{e } 2x^2 + 4x + 7 = 2\left(x^2 + 2x + \frac{7}{2}\right) = 2\left\{(x+1)^2 + \frac{5}{2}\right\}$$

$$\text{Let } u = (x+1), \text{ so } du = dx.$$

$$\begin{aligned} \text{Then } \int \frac{1}{2x^2 + 4x + 7} dx &= \frac{1}{2} \int \frac{1}{u^2 + \left(\frac{\sqrt{5}}{\sqrt{2}}\right)^2} du \\ &= \frac{1}{2} \left\{ \frac{\sqrt{2}}{\sqrt{5}} \arctan\left(\frac{\sqrt{2}u}{\sqrt{5}}\right) \right\} + C \\ &= \frac{\sqrt{10}}{10} \arctan\left(\frac{\sqrt{2}(x+1)}{\sqrt{5}}\right) + C \end{aligned}$$

$$\text{f } -4x^2 - 12x = -4\left(x^2 + 3x\right) = -4\left\{\left(x + \frac{3}{2}\right)^2 - \frac{9}{4}\right\} = 4\left\{\frac{9}{4} - \left(x + \frac{3}{2}\right)^2\right\}$$

$$\text{So } \int \frac{1}{\sqrt{-4x^2 - 12x}} dx = \frac{1}{2} \int \frac{1}{\sqrt{\left(\frac{3}{2}\right)^2 - \left(x + \frac{3}{2}\right)^2}} dx$$

$$\text{Let } u = \left(x + \frac{3}{2}\right), \text{ so } du = dx.$$

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{-4x^2 - 12x}} dx &= \frac{1}{2} \int \frac{1}{\sqrt{\left(\frac{3}{2}\right)^2 - u^2}} du \\ &= \frac{1}{2} \arcsin\left(\frac{2u}{3}\right) + C \\ &= \frac{1}{2} \arcsin\left(\frac{2x+3}{3}\right) + C \end{aligned}$$

$$\begin{aligned} \text{g } 14 - 12x - 2x^2 &= -2(x^2 + 6x - 7) \\ &= -2((x+3)^2 - 16) \\ &= 2(16 - (x+3)^2) \end{aligned}$$

$$\text{So } \int \frac{1}{\sqrt{14 - 12x - 2x^2}} dx = \frac{1}{\sqrt{2}} \int \frac{1}{\sqrt{4^2 - (x+3)^2}} dx$$

Let $u = x+3$, so $du = dx$

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{14 - 12x - 2x^2}} dx &= \frac{1}{\sqrt{2}} \int \frac{1}{\sqrt{4^2 - u^2}} du \\ &= \frac{1}{\sqrt{2}} \arcsin\left(\frac{u}{4}\right) + C \\ &= \frac{1}{\sqrt{2}} \arcsin\left(\frac{x+3}{4}\right) + C \end{aligned}$$

$$\text{h } 9x^2 - 8x + 1 = 9\left(x^2 - \frac{8}{9}x + \frac{1}{9}\right) = 9\left[\left(x - \frac{4}{9}\right)^2 - \frac{7}{81}\right]$$

$$\text{So } \int \frac{1}{\sqrt{9x^2 - 8x + 1}} dx = \frac{1}{3} \int \frac{1}{\sqrt{\left(x - \frac{4}{9}\right)^2 - \left(\frac{\sqrt{7}}{9}\right)^2}} dx$$

Let $u = \left(x - \frac{4}{9}\right)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{9x^2 - 8x + 1}} dx &= \frac{1}{3} \int \frac{1}{\sqrt{u^2 - \left(\frac{\sqrt{7}}{9}\right)^2}} du \\ &= \frac{1}{3} \operatorname{arcosh}\left(\frac{9u}{\sqrt{7}}\right) + C \\ &= \frac{1}{3} \operatorname{arcosh}\left(\frac{9x - 4}{\sqrt{7}}\right) + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 2

Question:

Find

$$\text{a } \int \frac{1}{\sqrt{4x^2 - 12x + 10}} dx$$

$$\text{b } \int \frac{1}{\sqrt{4x^2 - 12x + 4}} dx.$$

Solution:

$$\text{a } 4x^2 - 12x + 10 = 4\left(x^2 - 3x + \frac{5}{2}\right) = 4\left\{\left(x - \frac{3}{2}\right)^2 + \frac{1}{4}\right\}$$

$$\text{So } \int \frac{1}{\sqrt{4x^2 - 12x + 10}} dx = \frac{1}{2} \int \frac{1}{\sqrt{\left(x - \frac{3}{2}\right)^2 + \left(\frac{1}{2}\right)^2}} dx$$

$$\text{Let } u = \left(x - \frac{3}{2}\right), \text{ so } du = dx.$$

$$\text{Then } \int \frac{1}{\sqrt{4x^2 - 12x + 10}} dx = \frac{1}{2} \int \frac{1}{\sqrt{u^2 + \left(\frac{1}{2}\right)^2}} du$$

$$= \frac{1}{2} \operatorname{arsinh}(2u) + C$$

$$= \frac{1}{2} \operatorname{arsinh}(2x - 3) + C$$

$$\text{b } 4x^2 - 12x + 4 = 4(x^2 - 3x + 1) = 4\left\{\left(x - \frac{3}{2}\right)^2 - \frac{5}{4}\right\}$$

$$\text{So } \int \frac{1}{\sqrt{4x^2 - 12x + 4}} dx = \frac{1}{2} \int \frac{1}{\sqrt{\left(x - \frac{3}{2}\right)^2 - \left(\frac{\sqrt{5}}{2}\right)^2}} dx$$

$$\text{Let } u = \left(x - \frac{3}{2}\right), \text{ so } du = dx.$$

$$\text{Then } \int \frac{1}{\sqrt{4x^2 - 12x + 4}} dx = \frac{1}{2} \int \frac{1}{\sqrt{u^2 - \left(\frac{\sqrt{5}}{2}\right)^2}} du$$

$$= \frac{1}{2} \operatorname{arcosh}\left(\frac{2u}{\sqrt{5}}\right) + C$$

$$= \frac{1}{2} \operatorname{arcosh}\left(\frac{2x - 3}{\sqrt{5}}\right) + C$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 3

Question:

Evaluate the following, giving answers to 3 significant figures.

a $\int_1^3 \frac{1}{\sqrt{x^2 + 2x + 5}} dx$

b $\int_1^3 \frac{1}{x^2 + x + 1} dx$

c $\int_0^1 \frac{1}{\sqrt{2 + 3x - 2x^2}} dx$

Solution:

a $x^2 + 2x + 5 = (x+1)^2 + 4$

So $\int_0^1 \frac{1}{\sqrt{x^2 + 2x + 5}} dx = \int_0^1 \frac{1}{\sqrt{(x+1)^2 + 4}} dx$

Let $u = (x+1)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int_0^1 \frac{1}{\sqrt{x^2 + 2x + 5}} dx &= \int_1^2 \frac{1}{\sqrt{u^2 + 2^2}} du \\ &= \left[\operatorname{arsinh}\left(\frac{u}{2}\right) \right]_1^2 \\ &= \left[\operatorname{arsinh} 1 - \operatorname{arsinh}\left(\frac{1}{2}\right) \right] \\ &= 0.400 \text{ (3 s.f.)} \end{aligned}$$

b $\int_1^3 \frac{1}{x^2 + x + 1} dx = \int_1^3 \frac{1}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} dx$

Let $u = \left(x + \frac{1}{2}\right)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int_1^3 \frac{1}{x^2 + x + 1} dx &= \int_{\frac{3}{2}}^{\frac{7}{2}} \frac{1}{u^2 + \left(\frac{\sqrt{3}}{2}\right)^2} du \\ &= \left[\frac{2}{\sqrt{3}} \arctan\left(\frac{2u}{\sqrt{3}}\right) \right]_{\frac{3}{2}}^{\frac{7}{2}} \\ &= \frac{2}{\sqrt{3}} \left[\arctan\left(\frac{7}{\sqrt{3}}\right) - \arctan(\sqrt{3}) \right] \\ &= 0.325 \text{ (3 s.f.)} \end{aligned}$$

c $2 + 3x - 2x^2 = -2\left(x^2 - \frac{3}{2}x - 1\right) = -2\left\{\left(x - \frac{3}{4}\right)^2 - \frac{25}{16}\right\} = 2\left\{\frac{25}{16} - \left(x - \frac{3}{4}\right)^2\right\}$

So $\int_0^1 \frac{1}{\sqrt{2 + 3x - 2x^2}} dx = \frac{1}{\sqrt{2}} \int_0^1 \frac{1}{\sqrt{\left(\frac{5}{4}\right)^2 - \left(x - \frac{3}{4}\right)^2}} dx$

Let $u = \left(x - \frac{3}{4}\right)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int_0^1 \frac{1}{\sqrt{2 + 3x - 2x^2}} dx &= \frac{1}{\sqrt{2}} \int_{-\frac{3}{4}}^{\frac{1}{4}} \frac{1}{\sqrt{\left(\frac{5}{4}\right)^2 - u^2}} du \\ &= \frac{1}{\sqrt{2}} \left[\arcsin\left(\frac{4u}{5}\right) \right]_{-\frac{3}{4}}^{\frac{1}{4}} \\ &= \frac{1}{\sqrt{2}} \left[\arcsin\left(\frac{1}{5}\right) - \arcsin\left(\frac{-3}{5}\right) \right] \\ &= 0.597 \text{ (3 s.f.)} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 4

Question:

Evaluate

a $\int_1^3 \frac{1}{\sqrt{x^2 - 2x + 2}} dx$, giving your answer as a single natural logarithm,

b $\int_1^2 \frac{1}{\sqrt{1 + 6x - 3x^2}} dx$, giving your answer in the form $k\pi$.

Solution:

a $x^2 - 2x + 2 = (x - 1)^2 + 1$

$$\text{So } \int_1^3 \frac{1}{\sqrt{x^2 - 2x + 2}} dx = \int_1^3 \frac{1}{\sqrt{(x-1)^2 + 1}} dx$$

$$= [\operatorname{arsinh}(x-1)]_1^3$$

$$= \operatorname{arsinh} 2$$

$$= \ln \{2 + \sqrt{5}\}$$

$$\operatorname{arsinh} x = \ln \{x + \sqrt{x^2 + 1}\}$$

b $1 + 6x - 3x^2 = -3\left(x^2 - 2x - \frac{1}{3}\right) = -3\left\{(x-1)^2 - \frac{4}{3}\right\} = 3\left[\frac{4}{3} - (x-1)^2\right]$

$$\text{So } \int_1^2 \frac{1}{\sqrt{1 + 6x - 3x^2}} dx = \frac{1}{\sqrt{3}} \int_1^2 \frac{1}{\sqrt{\left(\frac{2}{\sqrt{3}}\right)^2 - (x-1)^2}} dx$$

$$= \frac{1}{\sqrt{3}} \left[\arcsin \left(\frac{\sqrt{3}(x-1)}{2} \right) \right]_1^2$$

$$= \frac{1}{\sqrt{3}} \arcsin \left(\frac{\sqrt{3}}{2} \right)$$

$$= \frac{\pi}{3\sqrt{3}}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 5

Question:

Show that $\int_1^3 \frac{1}{\sqrt{3x^2 - 6x + 7}} dx = \frac{1}{\sqrt{3}} \ln(2 + \sqrt{3})$.

Solution:

$$3x^2 - 6x + 7 = 3\left(x^2 - 2x + \frac{7}{3}\right) = 3\left\{(x-1)^2 + \frac{4}{3}\right\}$$

$$\text{So } \int \frac{1}{\sqrt{3x^2 - 6x + 7}} dx = \frac{1}{\sqrt{3}} \int \frac{1}{\sqrt{(x-1)^2 + \left(\frac{2}{\sqrt{3}}\right)^2}} dx$$

Let $u = (x-1)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int_1^3 \frac{1}{\sqrt{3x^2 - 6x + 7}} &= \frac{1}{\sqrt{3}} \int_0^2 \frac{1}{\sqrt{u^2 + \left(\frac{2}{\sqrt{3}}\right)^2}} du \\ &= \frac{1}{\sqrt{3}} \left[\operatorname{arsinh} \left(\frac{\sqrt{3}u}{2} \right) \right]_0^2 \\ &= \frac{1}{\sqrt{3}} \operatorname{arsinh} \sqrt{3} \\ &= \frac{1}{\sqrt{3}} \ln \left\{ \sqrt{3} + \sqrt{3+1} \right\} \quad \operatorname{arsinh} x = \ln \left\{ x + \sqrt{x^2 + 1} \right\} \\ &= \frac{1}{\sqrt{3}} \ln \left\{ 2 + \sqrt{3} \right\} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 6

Question:

Using a suitable hyperbolic or trigonometric substitution find

a $\int \frac{1}{\sqrt{x^2 + 4x + 5}} dx$

b $\int \frac{1}{\sqrt{-x^2 + 4x + 5}} dx.$

Solution:

a $x^2 + 4x + 5 = (x + 2)^2 + 1$

So let $(x + 2) = \sinh u$, then $dx = \cosh u du$ and $(x + 2)^2 + 1 = \sinh^2 u + 1 = \cosh^2 u$

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{x^2 + 4x + 5}} dx &= \int \frac{1}{\cosh u} \cosh u du \\ &= \int 1 du \\ &= u + C \\ &= \operatorname{arsinh}(x + 2) + C \end{aligned}$$

b $-x^2 + 4x + 5 = -(x^2 - 4x - 5) = -\{(x - 2)^2 - 9\} = 9 - (x - 2)^2$

So let $(x - 2) = 3 \sin \theta$, then $dx = 3 \cos \theta d\theta$

and $9 - (x - 2)^2 = 9(1 - \sin^2 \theta) = 9 \cos^2 \theta$

$$\begin{aligned} \text{Then } \int \frac{1}{\sqrt{-x^2 + 4x + 5}} dx &= \int \frac{1}{3 \cos \theta} 3 \cos \theta d\theta \\ &= \int 1 d\theta \\ &= \theta + C \\ &= \arcsin\left(\frac{x - 2}{3}\right) + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 7

Question:

Using the substitution $x = \frac{1}{5}(\sqrt{3} \tan \theta - 1)$, obtain $\int_{-0.2}^0 \frac{1}{25x^2 + 10x + 4} dx$, giving your answer in terms of π .

Solution:

Using the substitution $x = \frac{1}{5}(\sqrt{3} \tan \theta - 1)$, $dx = \frac{\sqrt{3}}{5} \sec^2 \theta d\theta$ and $25x^2 + 10x + 4 = (3 \tan^2 \theta - 2\sqrt{3} \tan \theta + 1) + 2(\sqrt{3} \tan \theta - 1) + 4$

$$= 3 \tan^2 \theta + 3$$

$$= 3(\tan^2 \theta + 1) = 3 \sec^2 \theta$$

$$\begin{aligned} \text{Then } \int_{-0.2}^0 \frac{1}{25x^2 + 10x + 4} dx &= \frac{\sqrt{3}}{5} \int_0^{\frac{\pi}{6}} \frac{1}{3 \sec^2 \theta} \sec^2 \theta d\theta \\ &= \frac{\sqrt{3}}{15} \int_0^{\frac{\pi}{6}} 1 d\theta \\ &= \frac{\pi \sqrt{3}}{90} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 8

Question:

Evaluate $\int_3^4 \frac{1}{\sqrt{(x-2)(x+4)}} dx$, giving your answer in the form $\ln(a+b\sqrt{c})$, where a , b and c are integers to be found.

Solution:

$$(x-2)(x+4) = x^2 + 2x - 8 = (x+1)^2 - 9$$

$$\text{So } \int_3^4 \frac{1}{\sqrt{(x-2)(x+4)}} dx = \int_3^4 \frac{1}{\sqrt{(x+1)^2 - 3^2}} dx$$

Let $u = (x+1)$, so $du = dx$.

$$\begin{aligned} \text{Then } \int_3^4 \frac{1}{\sqrt{(x-2)(x+4)}} dx &= \int_4^5 \frac{1}{\sqrt{u^2 - 3^2}} du \\ &= \left[\operatorname{arcosh} \left(\frac{u}{3} \right) \right]_4^5 \\ &= \operatorname{arcosh} \left(\frac{5}{3} \right) - \operatorname{arcosh} \left(\frac{4}{3} \right) \\ &= \ln \left\{ \left(\frac{5}{3} \right) + \sqrt{\frac{25}{9} - 1} \right\} - \ln \left\{ \left(\frac{4}{3} \right) + \sqrt{\frac{16}{9} - 1} \right\} \quad \boxed{\operatorname{arcosh} x = \ln \{x + \sqrt{x^2 - 1}\}} \\ &= \ln 3 - \ln \left\{ \frac{4 + \sqrt{7}}{3} \right\} \\ &= \ln \left(\frac{9}{4 + \sqrt{7}} \right) \quad \boxed{\ln a - \ln b = \ln \left(\frac{a}{b} \right)} \\ &= \ln \left(\frac{9(4 - \sqrt{7})}{9} \right) \quad \boxed{\text{Rationalising the denominator}} \\ &= \ln(4 - \sqrt{7}) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 9

Question:

Using the substitution $x = 1 + \sinh \theta$, show that

$$\int \frac{x}{(x^2 - 2x + 2)^{\frac{3}{2}}} dx = \frac{x-1}{\sqrt{x^2 - 2x + 2}} + C.$$

Solution:

Using the substitution $x = 1 + \sinh \theta$, $dx = \cosh \theta d\theta$ and

$$x^2 - 2x + 2 = (\sinh^2 \theta + 2\sinh \theta + 1) - 2(\sinh \theta + 1) + 2 = \sinh^2 \theta + 1 = \cosh^2 \theta$$

$$\text{So } \int \frac{1}{(x^2 - 2x + 2)^{\frac{3}{2}}} dx = \int \frac{1}{\cosh^3 \theta} \cdot \cosh \theta d\theta$$

$$= \int \operatorname{sech}^2 \theta d\theta$$

$$= \tanh \theta + C$$

$$= \frac{x-1}{\sqrt{x^2 - 2x + 2}} + C$$

$$\begin{aligned} \sinh \theta &= x-1 \\ \cosh \theta &= \sqrt{1 + \sinh^2 \theta} = \sqrt{2 - 2x + x^2} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise D, Question 10

Question:

Use the substitution $x = 2 \sin \theta - 1$ to find $\int \frac{x}{\sqrt{3-2x-x^2}} dx$.

Solution:

Using the substitution $x = 2 \sin \theta - 1$, $dx = 2 \cos \theta d\theta$

$$\begin{aligned} \text{and } 3-2x-x^2 &= 3-2(2 \sin \theta - 1) - (4 \sin^2 \theta - 4 \sin \theta + 1) \\ &= 4 - 4 \sin^2 \theta \\ &= 4 \cos^2 \theta \end{aligned}$$

$$\text{So } \int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{2 \sin \theta - 1}{2 \cos \theta} \cdot 2 \cos \theta d\theta$$

$$= \int (2 \sin \theta - 1) d\theta$$

$$= -2 \cos \theta - \theta + C$$

$$= -2 \sqrt{1 - \left(\frac{x+1}{2}\right)^2} - \theta + C$$

$\begin{aligned} \cos \theta &= \sqrt{1 - \sin^2 \theta} \\ \text{and } \sin \theta &= \frac{x+1}{2} \end{aligned}$

$$= -\sqrt{3-2x-x^2} - \arcsin\left(\frac{x+1}{2}\right) + C$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise E, Question 1

Question:

- a Show that $\int \operatorname{arsinh} x \, dx = x \operatorname{arsinh} x - \sqrt{1+x^2} + C$.
- b Evaluate $\int_0^1 \operatorname{arsinh} x \, dx$, giving your answer to 3 significant figures.
- c Using the substitution $u = 2x+1$ and the result in a, or otherwise, find $\int \operatorname{arsinh}(2x+1) \, dx$.

Solution:

a $I = \int 1 \cdot \operatorname{arsinh} x \, dx$

Let $u = \operatorname{arsinh} x \quad \frac{dv}{dx} = 1$

So $\frac{dx}{\sqrt{x^2+1}} \quad v = x$

So $I = x \operatorname{arsinh} x - \int \frac{x}{\sqrt{x^2+1}} \, dx$

Using integration by parts

$= x \operatorname{arsinh} x - \sqrt{x^2+1} + C$

Using

$$\int f^n(x) f'(x) \, dx = \frac{1}{n+1} f^{n+1}(x) + C, n \neq -1$$

b $\int_0^1 \operatorname{arsinh} x \, dx = \left[x \operatorname{arsinh} x - \sqrt{x^2+1} \right]_0^1$

$= \left[\operatorname{arsinh} 1 - \sqrt{2} \right] - [-1]$

$= 0.467 \text{ (3 s.f.)}$

c Let $u = 2x+1$, so $du = 2 \, dx$

Then $\int \operatorname{arsinh}(2x+1) \, dx = \frac{1}{2} \int \operatorname{arsinh} u \, du$

$= \frac{1}{2} \operatorname{arsinh} u - \sqrt{1+u^2} + C$ using a

$= \frac{1}{2} (2x+1) \operatorname{arsinh}(2x+1) - \sqrt{4x^2+4x+2} + C$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise E, Question 2

Question:

Show that $\int \arctan 3x \, dx = x \arctan 3x - \frac{1}{6} \ln(1+9x^2) + C$.

Solution:

$$\text{Let } u = \arctan 3x \quad \frac{dv}{dx} = 1$$

$$\text{So } \frac{du}{dx} = \frac{3}{1+(3x)^2} \quad v = x$$

$$\begin{aligned} \text{Then } \int \arctan 3x \, dx &= x \arctan 3x - \int \frac{3x}{1+9x^2} \, dx \\ &= x \arctan 3x - \frac{1}{6} \int \frac{18x}{1+9x^2} \, dx \\ &= x \arctan 3x - \frac{1}{6} \ln(1+9x^2) + C \end{aligned}$$

Using $\frac{du}{dx} = \frac{du}{dt} \times \frac{dt}{dx}$, where $u = \arctan t$
and $t = 3x$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise E, Question 3

Question:

- a Show that $\int \operatorname{arcosh} x \, dx = x \operatorname{arcosh} x - \sqrt{x^2 - 1} + C$.
- b Hence show that $\int_1^2 \operatorname{arcosh} x = \ln(7 + 4\sqrt{3}) - \sqrt{3}$.

Solution:

a Let $u = \operatorname{arcosh} x$ $\frac{dv}{dx} = 1$

So $\frac{dx}{dx} = \frac{1}{\sqrt{x^2 - 1}}$ $v = x$

$$\begin{aligned} \text{So } \int \operatorname{arcosh} x \, dx &= x \operatorname{arcosh} x - \int \frac{x}{\sqrt{x^2 - 1}} \, dx \\ &= x \operatorname{arcosh} x - \sqrt{x^2 - 1} + C \end{aligned}$$

- b Using limits

$$\int_1^2 \operatorname{arcosh} x = [2 \operatorname{arcosh} 2 - \sqrt{3}] - [\operatorname{arcosh} 1] = [2 \operatorname{arcosh} 2 - \sqrt{3}] \quad \boxed{\text{as } \operatorname{arcosh} 1 = 0}$$

As $\operatorname{arcosh} x = \ln \{x + \sqrt{x^2 - 1}\}$

$$\begin{aligned} \int_1^2 \operatorname{arcosh} x &= [2 \ln \{2 + \sqrt{3}\} - \sqrt{3}] \\ &= [\ln \{2 + \sqrt{3}\}^2 - \sqrt{3}] \\ &= \ln(7 + 4\sqrt{3}) - \sqrt{3} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise E, Question 4

Question:

a Show that $\int \arctan x \, dx = x \arctan x - \frac{1}{2} \ln(1+x^2) + C$.

b Hence show that $\int_{-1}^{\sqrt{3}} \arctan x \, dx = \frac{(4\sqrt{3}-3)\pi}{12} - \frac{1}{2} \ln 2$.

The curve C has equation $y = 2 \arctan x$. The region R is enclosed by C , the y -axis, the line $y = \pi$ and the line $x = 3$.

c Find the area of R , giving your answer to 3 significant figures.

Solution:

a $I = \int 1 \times \arctan x \, dx$

Let $u = \arctan x$ $\frac{dv}{dx} = 1$

So $\frac{du}{dx} = \frac{1}{1+x^2}$ $v = x$

So $I = x \arctan x - \int \frac{x}{1+x^2} \, dx$
 $= x \arctan x - \frac{1}{2} \ln(1+x^2) + C$

Using integration by parts

Using $\int \frac{f'(x)}{f(x)} \, dx = \ln f(x) + C$

b $\int_{-1}^{\sqrt{3}} \arctan x \, dx = \left[x \arctan x - \frac{1}{2} \ln(1+x^2) \right]_{-1}^{\sqrt{3}}$
 $= \left[\sqrt{3} \arctan \sqrt{3} - \frac{1}{2} \ln 4 \right] - \left[-\arctan(-1) - \frac{1}{2} \ln 2 \right]$
 $= \frac{\sqrt{3}\pi}{3} - \ln 2 + \left(-\frac{\pi}{4} \right) + \frac{1}{2} \ln 2$
 $= \frac{(4\sqrt{3}-3)\pi}{12} - \frac{1}{2} \ln 2$

c Area of $R = \text{area of rectangle} - \int_0^3 2 \arctan x \, dx$

$= 3\pi - 2 \left[x \arctan x - \frac{1}{2} \ln(1+x^2) \right]_0^3$

Using α

$= 3\pi - 6 \arctan 3 + \ln 10$

$= 4.23$ (3 s.f.)

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise E, Question 5

Question:

Evaluate

a $\int_0^{\frac{\sqrt{2}}{2}} \arcsin x \, dx$

b $\int_0^1 x \arctan x \, dx$ giving your answers in terms of π .

Solution:

a Let $u = \arcsin x \quad \frac{dv}{dx} = 1$

So $\frac{dx}{dx} = \frac{1}{\sqrt{1-x^2}} \quad v = x$

$$\begin{aligned} \text{Then } \int_0^{\frac{\sqrt{2}}{2}} \arcsin x \, dx &= \left[x \arcsin x \right]_0^{\frac{\sqrt{2}}{2}} - \int_0^{\frac{\sqrt{2}}{2}} \frac{x}{\sqrt{1-x^2}} \, dx \\ &= \left[x \arcsin x + \sqrt{1-x^2} \right]_0^{\frac{\sqrt{2}}{2}} \\ &= \left[\frac{\sqrt{2}}{2} \frac{\pi}{4} + \sqrt{\frac{1}{2}} \right] - [1] \\ &= \frac{\sqrt{2}}{8} \pi - 1 + \frac{\sqrt{2}}{2} = 0.262 \text{ (3 s.f.)} \end{aligned}$$

b Let $u = \arctan x \quad \frac{dv}{dx} = x$

So $\frac{dx}{dx} = \frac{1}{1+x^2} \quad v = \frac{x^2}{2}$

$$\begin{aligned} \text{Then } \int_0^1 x \arctan x \, dx &= \left[\frac{x^2}{2} \arctan x \right]_0^1 - \frac{1}{2} \int_0^1 \frac{x^2}{1+x^2} \, dx \\ &= \left[\frac{1}{2} \arctan 1 \right]_0^1 - \frac{1}{2} \int_0^1 \frac{1+x^2-1}{1+x^2} \, dx \\ &= \left[\frac{\pi}{8} \right] - \frac{1}{2} \int_0^1 \left(1 - \frac{1}{1+x^2} \right) \, dx \\ &= \left[\frac{\pi}{8} \right] - \frac{1}{2} \left[x - \arctan x \right]_0^1 \\ &= \left[\frac{\pi}{8} \right] - \frac{1}{2} \left[1 - \frac{\pi}{4} \right] \\ &= \frac{\pi-2}{4} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise E, Question 6

Question:

Using the result that if $y = \operatorname{arcsec} x$, then $\frac{dy}{dx} = \frac{1}{x\sqrt{x^2-1}}$, show that

$$\int \operatorname{arcsec} x dx = x \operatorname{arcsec} x - \ln(x + \sqrt{x^2 - 1}) + C.$$

Solution:

$$\text{Let } u = \operatorname{arcsec} x \quad \frac{dv}{dx} = 1$$

$$\text{So } \frac{du}{dx} = \frac{1}{x\sqrt{x^2-1}} \quad v = x$$

$$\begin{aligned} \text{and } \int \operatorname{arcsec} x dx &= x \operatorname{arcsec} x - \int \frac{x}{x\sqrt{x^2-1}} dx \\ &= x \operatorname{arcsec} x - \operatorname{arcosh} x + C \\ &= x \operatorname{arcsec} x - \ln \{x + \sqrt{x^2 - 1}\} + C \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise E, Question 7

Question:

- a Show that $\int \operatorname{arsinh}(2x+1) dx = x \operatorname{arsinh}(2x+1) - \int \frac{2x}{\sqrt{(2x+1)^2+1}} dx$.
- b Find $\int \frac{2x}{\sqrt{(2x+1)^2+1}} dx$, using the substitution $2x+1 = \sinh u$, and hence find $\int \operatorname{arcsin}(2x+1) dx$.

Solution:

a Let $u = \operatorname{arsinh}(2x+1)$ $\frac{dv}{dx} = 1$

So $\frac{dx}{dx} = \frac{2}{\sqrt{(2x+1)^2+1}}$ $v = x$

Then $\int \operatorname{arsinh}(2x+1) dx = x \operatorname{arsinh}(2x+1) - \int \frac{2x}{\sqrt{(2x+1)^2+1}} dx$

b Let $2x+1 = \sinh u$ then $2 dx = \cosh u du$

So $\int \frac{2x}{\sqrt{(2x+1)^2+1}} dx = \frac{1}{2} \int \frac{(\sinh u - 1)}{\cosh u} \cosh u du$

$$= \frac{1}{2} \left[\int \sinh u du - u \right]$$

$$= \frac{1}{2} [\cosh u - u] + C$$

$$= \frac{1}{2} \left(\sqrt{1+(2x+1)^2} - \operatorname{arsinh}(2x+1) \right) + C$$

$\int \operatorname{arsinh}(2x+1) dx = x \operatorname{arsinh}(2x+1) + \frac{1}{2} \operatorname{arsinh}(2x+1) - \frac{1}{2} \sqrt{1+(2x+1)^2} + C$

Using a and b.

$$= \frac{1}{2} (2x+1) \operatorname{arsinh}(2x+1) - \frac{1}{2} \sqrt{1+(2x+1)^2} + C$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 1

Question:

Given that $I_n = \int x^n e^{\frac{x}{2}} dx$,

a show that $I_n = 2x^n e^{\frac{x}{2}} - 2nI_{n-1}$, $n \geq 1$.

b Hence find $\int x^3 e^{\frac{x}{2}} dx$.

Solution:

a Integrating by parts with $u = x^n$ and $\frac{dv}{dx} = e^{\frac{x}{2}}$

so $\frac{du}{dx} = nx^{n-1}$, $v = 2e^{\frac{x}{2}}$

$$\begin{aligned} \text{So } I_n &= 2x^n e^{\frac{x}{2}} - \int 2nx^{n-1} e^{\frac{x}{2}} dx \\ &= 2x^n e^{\frac{x}{2}} - 2n \int x^{n-1} e^{\frac{x}{2}} dx \\ &= 2x^n e^{\frac{x}{2}} - 2nI_{n-1} \quad * \end{aligned}$$

b $I_3 = 2x^3 e^{\frac{x}{2}} - 6I_2$

$$= 2x^3 e^{\frac{x}{2}} - 6 \left(2x^2 e^{\frac{x}{2}} - 4I_1 \right)$$

$$= 2x^3 e^{\frac{x}{2}} - 12x^2 e^{\frac{x}{2}} + 24 \left(2xe^{\frac{x}{2}} - 2I_0 \right), \text{ where } I_0 = \int e^{\frac{x}{2}} dx = 2e^{\frac{x}{2}} + C$$

$$= 2x^3 e^{\frac{x}{2}} - 12x^2 e^{\frac{x}{2}} + 48xe^{\frac{x}{2}} - 48I_0$$

So $\int x^3 e^{\frac{x}{2}} dx = 2x^3 e^{\frac{x}{2}} - 12x^2 e^{\frac{x}{2}} + 48xe^{\frac{x}{2}} - 96e^{\frac{x}{2}} + C$

Substituting $n = 3, 2$ and 1 respectively in *

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 2

Question:

Given that $I_n = \int_1^e x(\ln x)^n dx, n \in \mathbb{N}$,

a show that $I_n = \frac{e^2}{2} - \frac{n}{2} I_{n-1}, n \in \mathbb{N}$.

b Hence show that $\int_1^e x(\ln x)^4 dx = \frac{e^2 - 3}{4}$.

Solution:

a Let $u = (\ln x)^n$ and $\frac{dv}{dx} = x$, so $\frac{du}{dx} = n \frac{(\ln x)^{n-1}}{x}, v = \frac{x^2}{2}$

Integration by parts:

$$\begin{aligned} \int_1^e x(\ln x)^n dx &= \left[\frac{x^2 (\ln x)^n}{2} \right]_1^e - \int_1^e \frac{nx^2 (\ln x)^{n-1}}{2x} dx \\ &= \left[\frac{e^2}{2} - 0 \right] - \frac{n}{2} \int_1^e x(\ln x)^{n-1} dx \end{aligned}$$

$$\text{So } I_n = \frac{e^2}{2} - \frac{n}{2} I_{n-1} \quad *$$

b $\int_1^e x(\ln x)^4 dx = I_4$

Substituting $n = 4, 3, 2$ and 1 respectively in the reduction formula *

$$\begin{aligned} I_4 &= \frac{e^2}{2} - \frac{4}{2} I_3 \\ &= \frac{e^2}{2} - 2 \left(\frac{e^2}{2} - \frac{3}{2} I_2 \right) \\ &= \frac{e^2}{2} - e^2 + 3 \left(\frac{e^2}{2} - \frac{2}{2} I_1 \right) \\ &= \frac{e^2}{2} - e^2 + \frac{3e^2}{2} - 3 \left(\frac{e^2}{2} - \frac{1}{2} I_0 \right), \text{ where } I_0 = \int_1^e x dx = \left[\frac{x^2}{2} \right]_1^e = \frac{e^2}{2} - \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \text{So } \int_1^e x(\ln x)^4 dx &= \frac{e^2}{2} - e^2 + \frac{3e^2}{2} - \frac{3e^2}{2} + \frac{3}{2} \left(\frac{e^2}{2} - \frac{1}{2} \right) \\ &= \frac{e^2}{4} - \frac{3}{4} = \frac{e^2 - 3}{4} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 3

Question:

In Example 21, you saw that, if $I_n = \int_0^1 x^n \sqrt{1-x} dx$, then $I_n = \frac{2n}{2n+3} I_{n-1}, n \geq 1$.

Use this reduction formula to evaluate $\int_0^1 (x+1)(x+2)\sqrt{1-x} dx$

Solution:

$$\begin{aligned} \int_0^1 [(x+1)(x+2)\sqrt{1-x}] dx &= \int_0^1 [(x^2+3x+2)\sqrt{1-x}] dx \\ &= \int_0^1 [x^2\sqrt{1-x}] dx + \int_0^1 [3x\sqrt{1-x}] dx + \int_0^1 [2\sqrt{1-x}] dx \\ &= I_2 + 3I_1 + 2I_0 \end{aligned}$$

$$\text{Now } I_0 = \int_0^1 \sqrt{1-x} dx = \left[-\frac{2}{3}(1-x)^{\frac{3}{2}} \right]_0^1 = 0 - \left(-\frac{2}{3} \right) = \frac{2}{3}$$

$$I_1 = \frac{2}{5} I_0 = \left(\frac{2}{5} \right) \left(\frac{2}{3} \right) = \frac{4}{15}$$

← Using the given formula with $n = 1$

$$I_2 = \frac{4}{7} I_1 = \left(\frac{4}{7} \right) \left(\frac{4}{15} \right) = \frac{16}{105}$$

← Using the given formula with $n = 2$

$$\begin{aligned} \text{So } \int_0^1 [(x+1)(x+2)\sqrt{1-x}] dx &= \frac{16}{105} + 3 \left(\frac{4}{15} \right) + 2 \left(\frac{2}{3} \right) \\ &= \frac{16 + 12(7) + 4(35)}{105} \\ &= \frac{240}{105} = \frac{16}{7} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 4

Question:

Given that $I_n = \int x^n e^{-x} dx$, where n is a positive integer,

a show that $I_n = -x^n e^{-x} + nI_{n-1}$, $n \geq 1$.

b Find $\int x^3 e^{-x} dx$.

c Evaluate $\int_0^1 x^4 e^{-x} dx$, giving your answer in terms of e .

Solution:

a Using integration by parts with $u = x^n$ and $\frac{dv}{dx} = e^{-x}$

so $\frac{du}{dx} = nx^{n-1}$ and $v = -e^{-x}$

$$\int x^n e^{-x} dx = -x^n e^{-x} - \int -nx^{n-1} e^{-x} dx, \text{ so } I_n = -x^n e^{-x} + nI_{n-1}$$

b Repeatedly using the reduction formula to find I_3

$$\begin{aligned} I_3 &= -x^3 e^{-x} + 3I_2 \\ &= -x^3 e^{-x} + 3(-x^2 e^{-x} + 2I_1) \\ &= -x^3 e^{-x} - 3x^2 e^{-x} + 6I_1 \\ &= -x^3 e^{-x} - 3x^2 e^{-x} + 6(-x e^{-x} + I_0) \end{aligned}$$

$$\text{But } I_0 = \int e^{-x} dx = -e^{-x} + C$$

$$\text{So } I_3 = -x^3 e^{-x} - 3x^2 e^{-x} - 6x e^{-x} - 6e^{-x} + K$$

c $I_4 = -x^4 e^{-x} + 4I_3$

$$= -x^4 e^{-x} + 4(-x^3 e^{-x} - 3x^2 e^{-x} - 6x e^{-x} - 6e^{-x} + C) \quad \boxed{\text{Using the result from b}}$$

$$\begin{aligned} \text{So } \int_0^1 x^4 e^{-x} dx &= [-x^4 e^{-x} - 4x^3 e^{-x} - 12x^2 e^{-x} - 24x e^{-x} - 24e^{-x}]_0^1 \\ &= [-65e^{-1}] - [-24] \\ &= 24 - 65e^{-1} \quad \text{or} \quad \frac{24e - 65}{e} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 5

Question:

$$I_n = \int \tanh^n x \, dx,$$

a By writing $\tanh^n x = \tanh^{n-2} x \tanh^2 x$, show that for $n \geq 2$,

$$I_n = I_{n-2} - \frac{1}{n-1} \tanh^{n-1} x.$$

b Find $\int \tanh^5 x \, dx$.

c Show that $\int_0^{\ln 2} \tanh^4 x \, dx = \ln 2 - \frac{84}{125}$.

Solution:

$$\begin{aligned} \text{a } I_n &= \int \tanh^n x \, dx = \int \tanh^{n-2} x \tanh^2 x \, dx \\ &= \int \tanh^{n-2} x (1 - \operatorname{sech}^2 x) \, dx \\ &= \int \tanh^{n-2} x \, dx - \int \tanh^{n-2} \operatorname{sech}^2 x \, dx \end{aligned}$$

Using $1 - \tanh^2 x = \operatorname{sech}^2 x$

$$\text{So } I_n = I_{n-2} - \frac{1}{n-1} \tanh^{n-1} x, \quad n \neq 1$$

$$\begin{aligned} \text{b } \int \tanh^5 x \, dx &= I_5 = I_3 - \frac{1}{4} \tanh^4 x \\ &= \left(I_1 - \frac{1}{2} \tanh^2 x \right) - \frac{1}{4} \tanh^4 x \\ &= \int \tanh x \, dx - \frac{1}{2} \tanh^2 x - \frac{1}{4} \tanh^4 x \\ &= \ln \cosh x - \frac{1}{2} \tanh^2 x - \frac{1}{4} \tanh^4 x + C \end{aligned}$$

$$\text{c } \text{As } \int \tanh^n x \, dx = \int \tanh^{n-2} x \, dx - \frac{1}{n-1} \tanh^{n-1} x, \text{ it follows that}$$

$$\int_0^{\ln 2} \tanh^n x \, dx = \int_0^{\ln 2} \tanh^{n-2} x \, dx - \left[\frac{1}{n-1} \tanh^{n-1} x \right]_0^{\ln 2} \quad *$$

$$\text{Now } \tanh(\ln 2) = \frac{e^{\ln 2} - e^{-\ln 2}}{e^{\ln 2} + e^{-\ln 2}} = \frac{2 - \frac{1}{2}}{2 + \frac{1}{2}} = \frac{3}{5}$$

Reminder: $e^{-\ln a} = e^{\ln a^{-1}} = a^{-1}$

$$\begin{aligned} \text{So } \int_0^{\ln 2} \tanh^4 x \, dx &= \int_0^{\ln 2} \tanh^2 x \, dx - \frac{1}{3} \times \left(\frac{3}{5} \right)^3 \\ &= \left[\int_0^{\ln 2} \tanh^0 x \, dx - 1 \times \left(\frac{3}{5} \right) \right] - \frac{1}{3} \times \frac{27}{125} \\ &= \ln 2 - \frac{3}{5} - \frac{9}{125} \\ &= \ln 2 - \frac{84}{125} \end{aligned}$$

Using * with $n = 4$ and $\tanh(\ln 2) = \frac{3}{5}$

Using * with $n = 2$ and $\tanh(\ln 2) = \frac{3}{5}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 6

Question:

Given that $\int \tan^x x \, dx = \frac{1}{x-1} \tan^{x-1} x - \int \tan^{x-2} x \, dx$ (derived in Example 23)

a find $\int \tan^4 x \, dx$.

b Evaluate $\int_0^{\frac{\pi}{4}} \tan^5 x \, dx$.

c Show that $\int_0^{\frac{\pi}{3}} \tan^6 x \, dx = \frac{9\sqrt{3}}{5} - \frac{\pi}{3}$.

Solution:

$$\begin{aligned}
 \text{a } \int \tan^4 x \, dx &= \frac{1}{3} \tan^3 x - \int \tan^2 x \, dx \\
 &= \frac{1}{3} \tan^3 x - \left(\tan x - \int \tan^0 x \, dx \right) \\
 &= \frac{1}{3} \tan^3 x - \tan x + \int 1 \, dx \\
 &= \frac{1}{3} \tan^3 x - \tan x + x + C
 \end{aligned}$$

$$\text{b } \int_0^{\frac{\pi}{4}} \tan^n x \, dx = \left[\frac{1}{n-1} \tan^{n-1} x \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \tan^{n-2} x \, dx = \frac{1}{n-1} - \int_0^{\frac{\pi}{4}} \tan^{n-2} x \, dx$$

$$\text{Let } I_n = \int_0^{\frac{\pi}{4}} \tan^n x \, dx, \text{ then } I_n = \frac{1}{n-1} - I_{n-2}$$

$$\begin{aligned}
 I_5 &= \frac{1}{4} - I_3 = \frac{1}{4} - \left(\frac{1}{2} - I_1 \right) = \frac{1}{4} - \frac{1}{2} + \int_0^{\frac{\pi}{4}} \tan x \, dx = \frac{1}{4} - \frac{1}{2} + [\ln \sec x]_0^{\frac{\pi}{4}} \\
 &= -\frac{1}{4} + (\ln \sqrt{2} - \ln 1)
 \end{aligned}$$

$$\text{So } \int_0^{\frac{\pi}{4}} \tan^5 x \, dx = \ln \sqrt{2} - \frac{1}{4}$$

$$\text{c } \text{Defining } J_n = \int_0^{\frac{\pi}{3}} \tan^n x \, dx,$$

$$J_n = \left[\frac{1}{n-1} \tan^{n-1} x \right]_0^{\frac{\pi}{3}} - J_{n-2} = \frac{(\sqrt{3})^{n-1}}{n-1} - J_{n-2}$$

$$\text{So } J_6 = \frac{(\sqrt{3})^5}{5} - J_4 = \frac{(\sqrt{3})^5}{5} - \left(\frac{(\sqrt{3})^3}{3} - J_2 \right) = \frac{(\sqrt{3})^5}{5} - \frac{(\sqrt{3})^3}{3} + \left(\frac{\sqrt{3}}{1} - J_0 \right)$$

$$\text{As } J_0 = \int_0^{\frac{\pi}{3}} 1 \, dx = \frac{\pi}{3}, \int_0^{\frac{\pi}{3}} \tan^6 x \, dx = \frac{9\sqrt{3}}{5} - \frac{3\sqrt{3}}{3} + \sqrt{3} - \frac{\pi}{3} = \frac{9\sqrt{3}}{5} - \frac{\pi}{3}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 7

Question:

Given that $I_n = \int_1^a (\ln x)^n dx$, where $a > 1$ is a constant,

a show that, for $n \geq 1$, $I_n = a(\ln a)^n - nI_{n-1}$.

b Find the exact value of $\int_1^2 (\ln x)^3 dx$.

c Show that $\int_1^e (\ln x)^6 dx = 5(53e - 144)$.

Solution:

$$\text{a } I_n = \int_1^a (\ln x)^n dx = \int_1^a 1(\ln x)^n dx$$

$$\text{Let } u = (\ln x)^n \text{ and } \frac{dv}{dx} = 1, \text{ so } \frac{du}{dx} = n \frac{(\ln x)^{n-1}}{x}, v = x$$

Integration by parts:

$$\begin{aligned} \int_1^a (\ln x)^n dx &= \left[x(\ln x)^n \right]_1^a - \int_1^a \frac{n(\ln x)^{n-1}}{x} x dx \\ &= \left[a(\ln a)^n - 0 \right] - n \int_1^a (\ln x)^{n-1} dx \end{aligned}$$

$$\text{So } I_n = a(\ln a)^n - nI_{n-1}$$

$$\text{b Putting } a = 2, I_n = \int_1^2 (\ln x)^n dx = 2(\ln 2)^n - nI_{n-1}$$

$$\begin{aligned} I_3 &= \int_1^2 (\ln x)^3 dx = 2(\ln 2)^3 - 3I_2 \\ &= 2(\ln 2)^3 - 3\{2(\ln 2)^2 - 2I_1\} \\ &= 2(\ln 2)^3 - 6(\ln 2)^2 + 6\{2(\ln 2) - I_0\} \\ &= 2(\ln 2)^3 - 6(\ln 2)^2 + 12(\ln 2) - 6I_0 \end{aligned}$$

$$\text{As } I_0 = \int_1^2 1 dx = [x]_1^2 = 1,$$

$$\int_1^2 (\ln x)^3 dx = 2(\ln 2)^3 - 6(\ln 2)^2 + 12(\ln 2) - 6$$

$$\text{c Putting } a = e, I_n = \int_1^e (\ln x)^n dx = e(\ln e)^n - nI_{n-1} = e - nI_{n-1}$$

$$\begin{aligned} I_6 &= \int_1^e (\ln x)^6 dx = e - 6I_5 \\ &= e - 6(e - 5I_4) \\ &= e - 6e + 30(e - 4I_3) \\ &= e - 6e + 30e - 120(e - 3I_2) \\ &= e - 6e + 30e - 120e + 360(e - 2I_1) \\ &= e - 6e + 30e - 120e + 360e - 720(e - I_0) \end{aligned}$$

$$\text{As } I_0 = \int_1^e 1 dx = [x]_1^e = e - 1,$$

$$\begin{aligned} \int_1^e (\ln x)^6 dx &= e - 6e + 30e - 120e + 360e - 720e + 720(e - 1) \\ &= 265e - 720 \\ &= 5(53e - 144) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 8

Question:

Using the results given in Example 22, evaluate

a $\int_0^{\frac{\pi}{2}} \sin^7 x \, dx$

b $\int_0^{\frac{\pi}{2}} \sin^2 x \cos^4 x \, dx$

c $\int_0^1 x^5 \sqrt{1-x^2} \, dx$, using the substitution $x = \sin \theta$

d $\int_0^{\frac{\pi}{6}} \sin^8 3t \, dt$, using a suitable substitution.

Solution:

a $I_7 = \frac{6}{7} \times \frac{4}{5} \times \frac{2}{3} \times 1 = \frac{16}{35}$

b $\int_0^{\frac{\pi}{2}} \sin^2 x \cos^4 x \, dx = \int_0^{\frac{\pi}{2}} \sin^2 x (1 - \sin^2 x)^2 \, dx = \int_0^{\frac{\pi}{2}} (\sin^2 x - 2\sin^4 x + \sin^6 x) \, dx$
 $= I_2 - 2I_4 + I_6$

$$I_2 = \frac{1}{2} \times \frac{\pi}{2} = \frac{\pi}{4}; \quad I_4 = \frac{3}{4} I_2 = \frac{3\pi}{16}; \quad I_6 = \frac{5}{6} I_4 = \frac{5\pi}{32}$$

So $\int_0^{\frac{\pi}{2}} \sin^2 x \cos^4 x \, dx = \frac{\pi}{4} - \frac{3\pi}{8} + \frac{5\pi}{32} = \frac{\pi}{32}$

c Using $x = \sin \theta$, $\int_0^1 x^5 \sqrt{1-x^2} \, dx = \int_0^{\frac{\pi}{2}} \sin^5 \theta \cos \theta (\cos \theta \, d\theta)$
 $= \int_0^{\frac{\pi}{2}} \sin^5 x (1 - \sin^2 x) \, dx = I_5 - I_7$

$$I_5 = \frac{4}{5} \times \frac{2}{3} \times 1 = \frac{8}{15} \quad \text{and} \quad I_7 = \frac{16}{35} \quad \text{from a}$$

So $\int_0^1 x^5 \sqrt{1-x^2} \, dx = \frac{8}{15} - \frac{16}{35} = \frac{56-48}{105} = \frac{8}{105}$

d Using $x = 3t$, $\int_0^{\frac{\pi}{6}} \sin^8 3t \, dt = \int_0^{\frac{\pi}{2}} \sin^8 x \left(\frac{1}{3} \, dx \right) = \frac{1}{3} I_8$
 $= \frac{1}{3} \times \frac{7}{8} \times \frac{5}{6} \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2} = \frac{35\pi}{768}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 9

Question:

Given that $I_n = \int \frac{\sin^{2n} x}{\cos x} dx$,

a write down a similar expression for I_{n+1} and hence show that $I_n - I_{n+1} = \frac{\sin^{2n+1} x}{2n+1}$.

b Find $\int \frac{\sin^4 x}{\cos x} dx$ and hence show that $\int_0^{\frac{\pi}{4}} \frac{\sin^4 x}{\cos x} dx = \ln(1+\sqrt{2}) - \frac{7\sqrt{2}}{12}$.

Solution:

$$\mathbf{a} \quad I_{n+1} = \int \frac{\sin^{2n+2} x}{\cos x} dx$$

$$\begin{aligned} \text{So } I_n - I_{n+1} &= \int \frac{\sin^{2n} x - \sin^{2n+2} x}{\cos x} dx \\ &= \int \frac{\sin^{2n} x (1 - \sin^2 x)}{\cos x} dx \\ &= \int \sin^{2n} x \cos x dx \end{aligned}$$

as $1 - \sin^2 x = \cos^2 x$

$$\text{So } I_n - I_{n+1} = \frac{\sin^{2n+1} x}{2n+1}$$

$$\text{or } I_{n+1} = I_n - \frac{\sin^{2n+1} x}{2n+1} \quad \#$$

[+C not necessary at this stage]

$$\mathbf{b} \quad \mathbf{i} \quad \int \frac{\sin^4 x}{\cos x} dx = I_2$$

$$\text{Substituting } n=1 \text{ in } \# \text{ gives } I_2 = I_1 - \frac{\sin^3 x}{3}$$

$$= \left(I_0 - \frac{\sin x}{1} \right) - \frac{\sin^3 x}{3} \text{ using } n=0 \text{ in } \#$$

$$I_0 = \int \frac{1}{\cos x} dx = \int \sec x dx = \ln |(\sec x + \tan x)| + C$$

$$\text{So } \int \frac{\sin^4 x}{\cos x} dx = \ln |(\sec x + \tan x)| - \sin x - \frac{\sin^3 x}{3} + C$$

Applying the given limits gives

$$\int_0^{\frac{\pi}{4}} \frac{\sin^4 x}{\cos x} dx = \left[\ln |(\sec x + \tan x)| - \sin x - \frac{\sin^3 x}{3} \right]_0^{\frac{\pi}{4}}$$

$$= \ln(1 + \sqrt{2}) - \frac{\sqrt{2}}{2} - \frac{\left(\frac{\sqrt{2}}{2}\right)^3}{3}$$

$$= \ln(1 + \sqrt{2}) - \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{12}$$

$$= \ln(1 + \sqrt{2}) - \frac{7\sqrt{2}}{12}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 10

Question:

- a Given that $I_n = \int_0^1 x(1-x^3)^n dx$, show that $I_n = \frac{3n}{3n+2} I_{n-1}$, $n \geq 1$.
- b Use your reduction formula to evaluate I_4 .

Hint: After integrating by parts, write x^4 as $x\{1-(1-x^3)\}$

Solution:

- a Let $u = (1-x^3)^n$ and $\frac{dv}{dx} = x$, so $\frac{du}{dx} = n(1-x^3)^{n-1}(-3x^2)$, $v = \frac{x^2}{2}$

Integration by parts gives

$$\begin{aligned} \int_0^1 x(1-x^3)^n dx &= \left[\frac{x^2}{2}(1-x^3)^n \right]_0^1 - \int_0^1 -3nx^2(1-x^3)^{n-1} \frac{x^2}{2} dx \\ &= [0-0] + \frac{3n}{2} \int_0^1 x^4(1-x^3)^{n-1} dx \quad \text{providing } n \geq 0 \end{aligned}$$

Writing $x^4 = x \cdot x^3 = x\{1-(1-x^3)\}$ and $I_n = \int_0^1 x(1-x^3)^n dx$

$$\begin{aligned} \text{we have } I_n &= \frac{3n}{2} \int_0^1 x\{1-(1-x^3)\}(1-x^3)^{n-1} dx \\ &= \frac{3n}{2} \int_0^1 x(1-x^3)^{n-1} dx - \frac{3n}{2} \int_0^1 x(1-x^3)^n dx \\ &= \frac{3n}{2} I_{n-1} - \frac{3n}{2} I_n \end{aligned}$$

$$\Rightarrow (3n+2)I_n = 3nI_{n-1}, \text{ so } I_n = \frac{3n}{3n+2} I_{n-1}, n \geq 1$$

- b $I_4 = \frac{12}{14} I_3 = \frac{12}{14} \times \frac{9}{11} I_2 = \frac{12}{14} \times \frac{9}{11} \times \frac{6}{8} I_1 = \frac{12}{14} \times \frac{9}{11} \times \frac{6}{8} \times \frac{3}{5} I_0 = \frac{12}{14} \times \frac{9}{11} \times \frac{6}{8} \times \frac{3}{5} \int_0^1 x dx$
 $= \frac{12}{14} \times \frac{9}{11} \times \frac{6}{8} \times \frac{3}{5} \times \frac{1}{2} = \frac{243}{1540}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 11

Question:

Given that $I_n = \int_0^a (a^2 - x^2)^n dx$, where a is a positive constant,

a show that, for $n > 0$, $I_n = \frac{2na^2}{2n+1} I_{n-1}$.

b Use the reduction formula to evaluate

i $\int_0^1 (1-x^2)^4 dx$

ii $\int_0^3 (9-x^2)^3 dx$

iii $\int_0^2 \sqrt{4-x^2} dx$.

c Check your answer to part **b iii** by using another method.

Solution:

- a Integrating by parts with $u = (a^2 - x^2)^n$ and $\frac{dv}{dx} = 1$

$$\frac{du}{dx} = -2nx(a^2 - x^2)^{n-1} \quad v = x$$

$$\begin{aligned} \text{So } \int_0^a (a^2 - x^2)^n dx &= \left[x(a^2 - x^2)^n \right]_0^a - \int_0^a x \{ -2nx(a^2 - x^2)^{n-1} \} dx \\ &= [0 - 0] + 2n \int_0^a x^2 (a^2 - x^2)^{n-1} dx = 2n \int_0^a x^2 (a^2 - x^2)^{n-1} dx \quad (\text{if } n > 0) \end{aligned}$$

Writing x^2 as $\{a^2 - (a^2 - x^2)\}$ and defining $I_n = \int_0^a (a^2 - x^2)^n dx$,

we have

$$\begin{aligned} I_n &= 2n \int_0^a \{ a^2 (a^2 - x^2)^{n-1} - (a^2 - x^2)^n \} dx \\ &= 2na^2 I_{n-1} - 2n I_n \end{aligned}$$

$$\text{So } (2n+1)I_n = 2na^2 I_{n-1}$$

- b i With $a = 1$, $I_n = \int_0^1 (1 - x^2)^n dx$ and $I_n = \frac{2n}{2n+1} I_{n-1}$

$$\text{So } I_4 = \frac{8}{9} I_3 = \frac{8}{9} \times \frac{6}{7} I_2 = \frac{8}{9} \times \frac{6}{7} \times \frac{4}{5} I_1 = \frac{8}{9} \times \frac{6}{7} \times \frac{4}{5} \times \frac{2}{3} \times 1 = \frac{128}{315}$$

$$I_0 = \int_0^a dx = a$$

- ii With $a = 3$, $I_n = \int_0^3 (9 - x^2)^n dx$ and $I_n = \frac{18n}{2n+1} I_{n-1}$

$$\text{So } I_3 = \frac{54}{7} I_2 = \frac{54}{7} \times \frac{36}{5} I_1 = \frac{54}{7} \times \frac{36}{5} \times \frac{18}{3} I_0 = \frac{54}{7} \times \frac{36}{5} \times \frac{18}{3} \times 3 = \frac{34\,992}{35}$$

- iii With $a = 2$, $I_n = \int_0^2 (4 - x^2)^n dx$ and $I_n = \frac{8n}{2n+1} I_{n-1}$

$$\text{So } I_{\frac{1}{2}} = \frac{4}{2} I_{0\frac{1}{2}} = 2 \int_0^2 \frac{dx}{\sqrt{4-x^2}} = 2 \left[\arcsin \left(\frac{x}{2} \right) \right]_0^2 = 2 \arcsin 1 = 2 \times \frac{\pi}{2} = \pi$$

- c Using the substitution $x = 2 \sin \theta$,

$$\begin{aligned} \int_0^2 (4 - x^2)^{\frac{1}{2}} dx &= \int_0^{\frac{\pi}{2}} (2 \cos \theta)(2 \cos \theta d\theta) \\ &= 2 \int_0^{\frac{\pi}{2}} (1 + \cos 2\theta) d\theta \\ &= [2\theta + \sin 2\theta]_0^{\frac{\pi}{2}} = \pi \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 12

Question:

Given that $I_n = \int_0^4 x^n \sqrt{4-x} \, dx$,

a establish the reduction formula $I_n = \frac{8n}{2n+3} I_{n-1}, n \geq 1$.

b Evaluate $\int_0^4 x^3 \sqrt{4-x} \, dx$, giving your answer correct to 3 significant figures.

Solution:

a Integrating by parts with $u = x^n$ and $\frac{dv}{dx} = \sqrt{4-x}$

$$\frac{du}{dx} = nx^{n-1}, \quad v = -\frac{2}{3}(4-x)^{\frac{3}{2}}$$

$$\begin{aligned} \text{So } \int_0^4 x^n \sqrt{4-x} \, dx &= \left[-\frac{2}{3} x^n (4-x)^{\frac{3}{2}} \right]_0^4 - \int_0^4 -\frac{2}{3} nx^{n-1} (4-x)^{\frac{3}{2}} \, dx \\ &= [0-0] + \frac{2}{3} n \int_0^4 x^{n-1} (4-x)^{\frac{3}{2}} \, dx \quad (n > 0) \\ &= \frac{2}{3} n \int_0^4 x^{n-1} \{(4-x)\sqrt{4-x}\} \, dx \\ &= \frac{2}{3} n \int_0^4 x^{n-1} 4\sqrt{4-x} \, dx + \frac{2}{3} n \int_0^4 x^{n-1} \{-x\sqrt{4-x}\} \, dx \\ &= \frac{8}{3} n \int_0^4 x^{n-1} \sqrt{4-x} \, dx - \frac{2}{3} n \int_0^4 x^n \sqrt{4-x} \, dx \end{aligned}$$

You need to write $(4-x)^{\frac{3}{2}}$
as $(4-x)\sqrt{4-x}$

$$\text{So } I_n = \frac{8}{3} n I_{n-1} - \frac{2}{3} n I_n$$

$$\Rightarrow (2n+3)I_n = 8nI_{n-1} \leq I_n = \frac{8n}{2n+3} I_{n-1}, n \geq 1$$

b $\int_0^4 x^3 \sqrt{4-x} \, dx = I_3 = \frac{24}{9} I_2 = \frac{24}{9} \times \frac{16}{7} I_1 = \frac{24}{9} \times \frac{16}{7} \times \frac{8}{5} I_0 = \frac{1024}{105} I_0$

$$\text{As } I_0 = \int_0^4 \sqrt{4-x} \, dx = \left[-\frac{2}{3}(4-x)^{\frac{3}{2}} \right]_0^4 = \left[0 - \left\{ -\frac{2}{3}(4)^{\frac{3}{2}} \right\} \right] = \frac{16}{3},$$

$$\int_0^4 x^3 \sqrt{4-x} \, dx = \frac{1024}{105} \times \frac{16}{3} = 52.0 \text{ (3 s.f.)}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 13

Question:

Given that $I_n = \int \cos^n x \, dx$,

a establish, for $n \geq 2$, the reduction formula $nI_n = \cos^{n-1} x \sin x + (n-1)I_{n-2}$.

Defining $J_n = \int_0^{2\pi} \cos^n x \, dx$,

b write down a reduction formula relating J_n and J_{n-2} , for $n \geq 2$.

c Hence evaluate

i J_4

ii J_8 .

d Show that if n is odd, J_n is always equal to zero.

Solution:

a $I_n = \int \cos^n x \, dx = \int \cos^{n-1} x \cos x \, dx$

Integrating by parts with $u = \cos^{n-1} x$ and $\frac{dv}{dx} = \cos x$

$$\frac{du}{dx} = (n-1)\cos^{n-2} x(-\sin x), \quad v = \sin x$$

$$\begin{aligned} \text{So } I_n &= \int \cos^n x \, dx = \cos^{n-1} x \sin x - \int -(n-1)\cos^{n-2} x \sin^2 x \, dx \\ &= \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x (1 - \cos^2 x) \, dx \\ &= \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x \, dx - (n-1) \int \cos^n x \, dx \end{aligned}$$

Giving $I_n = \cos^{n-1} x \sin x + (n-1)I_{n-2} - (n-1)I_n$

So $nI_n = \cos^{n-1} x \sin x + (n-1)I_{n-2}$

b It follows that $n \int_0^{2\pi} \cos^n x \, dx = [\cos^{n-1} x \sin x]_0^{2\pi} + (n-1) \int_0^{2\pi} \cos^{n-2} x \, dx$

So $nJ_n = (n-1)J_{n-2}$, as $[\cos^{n-1} x \sin x]_0^{2\pi} = 0$

c i $J_4 = \int_0^{2\pi} \cos^4 x \, dx = \frac{3}{4}J_2 = \frac{3}{4} \times \frac{1}{2}J_0 = \frac{3}{8} \int_0^{2\pi} 1 \, dx = \frac{3}{8} \times 2\pi = \frac{3\pi}{4}$

ii $J_8 = \int_0^{2\pi} \cos^8 x \, dx = \frac{7}{8}J_6 = \frac{7}{8} \times \frac{5}{6}J_4 = \frac{35}{48}J_4 = \frac{35}{48} \times \frac{3\pi}{4} = \frac{35\pi}{64}$ using c i

d If n is odd, J_n always reduces to a multiple of J_1 .

but $J_1 = \int_0^{2\pi} \cos x \, dx = [\sin x]_0^{2\pi} = 0$.

(You could also consider the graphical representation.)

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 14

Question:

Given $I_n = \int_0^1 x^n \sqrt{1-x^2} dx, n \geq 0,$

a show that $(n+2)I_n = (n-1)I_{n-2}, n \geq 2.$

b Hence evaluate $\int_0^1 x^7 \sqrt{1-x^2} dx.$

Hint: Write $x^n \sqrt{1-x^2}$ as $x^{n-1} \{x\sqrt{1-x^2}\}$ before integrating by parts.

Solution:

a Integrating by parts with $u = x^{n-1}$ and $\frac{dv}{dx} = x\sqrt{1-x^2}$

Using the hint.

$$\frac{du}{dx} = (n-1)x^{n-2}, \quad v = -\frac{1}{3}(1-x^2)^{\frac{3}{2}}$$

$$\text{So } I_n = \int_0^1 x^{n-1} \{x\sqrt{1-x^2}\} dx = \left[-\frac{1}{3}x^{n-1}(1-x^2)^{\frac{3}{2}} \right]_0^1 + \frac{(n-1)}{3} \int_0^1 x^{n-2}(1-x^2)^{\frac{3}{2}} dx$$

$$= \frac{(n-1)}{3} \int_0^1 x^{n-2}(1-x^2)^{\frac{3}{2}} dx \quad \text{as } \left[-\frac{1}{3}x^{n-1}(1-x^2)^{\frac{3}{2}} \right]_0^1 = 0$$

$$= \frac{(n-1)}{3} \int_0^1 x^{n-2}(1-x^2)\sqrt{1-x^2} dx$$

$$= \frac{(n-1)}{3} \int_0^1 \{x^{n-2}\sqrt{1-x^2} - x^n\sqrt{1-x^2}\} dx$$

$$\text{So } I_n = \frac{(n-1)}{3} I_{n-2} - \frac{(n-1)}{3} I_n$$

$$\Rightarrow \{3+(n-1)\}I_n = (n-1)I_{n-2}$$

$$\Rightarrow (n+2)I_n = (n-1)I_{n-2} \quad *$$

b Using * $I_7 = \frac{6}{9}I_5 = \frac{6}{9} \times \frac{4}{7}I_3 = \frac{6}{9} \times \frac{4}{7} \times \frac{2}{5}I_1 = \frac{48}{315} \int_0^1 x\sqrt{1-x^2} dx$

$$= \frac{48}{315} \left[-\frac{1}{3}(1-x^2)^{\frac{3}{2}} \right]_0^1$$

$$= \frac{48}{315} \left[\frac{1}{3} \right] = \frac{16}{315}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 15

Question:

Given $I_n = \int x^n \cosh x \, dx$

a show that for $n \geq 2$, $I_n = x^n \sinh x - nx^{n-1} \cosh x + n(n-1)I_{n-2}$

b Find $I_4 = \int x^4 \cosh x \, dx$.

c Evaluate $\int_0^1 x^3 \cosh x$, giving your answer in terms of e.

Solution:

a Integrating by parts with $u = x^n$ and $\frac{dv}{dx} = \cosh x$

$$\frac{du}{dx} = nx^{n-1}, \quad v = \sinh x$$

$$\text{So } \int x^n \cosh x dx = x^n \sinh x - \int nx^{n-1} \sinh x dx$$

Integrating by parts again with $u = x^{n-1}$ and $\frac{dv}{dx} = \sinh x$

$$\frac{du}{dx} = (n-1)x^{n-2}, \quad v = \cosh x$$

$$\begin{aligned} \text{So } I_n &= x^n \sinh x - n \left\{ x^{n-1} \cosh x - \int (n-1)x^{n-2} \cosh x dx \right\} \\ &= x^n \sinh x - nx^{n-1} \cosh x + n(n-1)I_{n-2}, \quad n \geq 2 \quad * \end{aligned}$$

b $I_4 = x^4 \sinh x - 4x^3 \cosh x + 12I_2$, ← Substituting $n = 4$ in *

$$= x^4 \sinh x - 4x^3 \cosh x + 12 \left\{ x^2 \sinh x - 2x \cosh x + 2I_0 \right\}$$

$$= x^4 \sinh x - 4x^3 \cosh x + 12 \left\{ x^2 \sinh x - 2x \cosh x \right\} + 24 \int \cosh x dx$$

$$= x^4 \sinh x - 4x^3 \cosh x + 12 \left\{ x^2 \sinh x - 2x \cosh x \right\} + 24 \sinh x + C$$

$$= (x^4 + 12x^2 + 24) \sinh x - (4x^3 + 24x) \cosh x + C$$

c $\int_0^1 x^3 \cosh x dx = \left[x^3 \sinh x - 3x^2 \cosh x \right]_0^1 + 6 \int_0^1 x \cosh x dx$ Using a

$$= \{ \sinh 1 - 3 \cosh 1 \} + 6 \left\{ \left[x \sinh x \right]_0^1 - \int_0^1 1 \sinh x dx \right\}$$
 Integrating by parts

$$= \{ \sinh 1 - 3 \cosh 1 \} + 6 \{ \sinh 1 - [\cosh 1 - 1] \}$$

$$= 7 \sinh 1 - 9 \cosh 1 + 6$$

$$= 7 \left(\frac{e^1 - e^{-1}}{2} \right) - 9 \left(\frac{e^1 + e^{-1}}{2} \right) + 6$$

$$= 6 - e - 8e^{-1} \text{ or } \frac{6e - e^2 - 8}{e}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 16

Question:

Given that $I_n = \int \frac{\sin nx}{\sin x} dx, n > 0$,

a write down a similar expression for I_{n-2} , and hence show that

$$I_n - I_{n-2} = \frac{2 \sin(n-1)x}{n-1}.$$

b Find

i $\int \frac{\sin 4x}{\sin x} dx$

ii the exact value of $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin 5x}{\sin x} dx$.

Solution:

$$\text{a } I_{n-2} = \int \frac{\sin(n-2)x}{\sin x} dx$$

$$\text{So } I_n - I_{n-2} = \int \frac{\sin nx - \sin(n-2)x}{\sin x} dx$$

$$= \int \frac{2 \cos \left\{ \frac{n+(n-2)}{2} \right\} x \sin \left\{ \frac{n-(n-2)}{2} \right\} x}{\sin x} dx$$

Using Edexcel formula booklet

$$= \int \frac{2 \cos(n-1)x \sin x}{\sin x} dx$$

$$= \int 2 \cos(n-1)x dx$$

$$= \frac{2 \sin(n-1)x}{n-1}, n \geq 2$$

It is not necessary to have +C.

$$\text{b i } \int \frac{\sin 4x}{\sin x} dx = I_4$$

$$\text{Using a with } n=4: I_4 = I_2 + \frac{2 \sin 3x}{3}$$

$$= \int 2 \cos x dx + \frac{2 \sin 3x}{3}$$

$$= 2 \sin x + \frac{2 \sin 3x}{3} + C$$

$I_2 = \int \frac{\sin 2x}{\sin x} dx = \int \frac{2 \sin x \cos x}{\sin x} dx$

$$\text{ii Using a with } n=5: I_5 = I_3 + \frac{2 \sin 4x}{4}$$

$$= \left\{ I_1 + \frac{2 \sin 2x}{2} \right\} + \frac{2 \sin 4x}{4}$$

$$= \int 1 dx + \sin 2x + \frac{\sin 4x}{2}$$

$$= x + \sin 2x + \frac{\sin 4x}{2}$$

$$\text{It follows that } \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin 5x}{\sin x} dx = \left[x + \sin 2x + \frac{\sin 4x}{2} \right]_{\frac{\pi}{6}}^{\frac{\pi}{3}}$$

$$= \left[\frac{\pi}{3} + \frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{4} \right] - \left[\frac{\pi}{6} + \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{4} \right]$$

$$= \frac{\pi}{6} - \frac{\sqrt{3}}{2}$$

$$= \frac{\pi - 3\sqrt{3}}{6}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise F, Question 17

Question:

Given that $I_n = \int \sinh^n x \, dx, n \in \mathbb{N}$,

a derive the reduction formula $nI_n = \sinh^{n-1} x \cosh x - (n-1)I_{n-2}, n \geq 2$.

b Hence

i evaluate $\int_0^{\ln 3} \sinh^5 x \, dx$,

ii show that $\int_0^{\operatorname{arsinh} 1} \sinh^4 x \, dx = \frac{1}{8}(3\ln(1+\sqrt{2})-\sqrt{2})$.

Solution:

$$\text{a } I_n = \int \sinh^n x \, dx = \int \sinh^{n-1} x \sinh x \, dx$$

Integrating by parts with $u = \sinh^{n-1} x$ and $\frac{dv}{dx} = \sinh x$

$$\frac{du}{dx} = (n-1) \sinh^{n-2} x \cosh x, \quad v = \cosh x$$

$$\begin{aligned} \text{So } I_n &= \int \sinh^n x \, dx = \sinh^{n-1} x \cosh x - \int (n-1) \sinh^{n-2} x \cosh^2 x \, dx \\ &= \sinh^{n-1} x \cosh x - (n-1) \int \sinh^{n-2} x (1 + \sinh^2 x) \, dx \\ &= \sinh^{n-1} x \cosh x - (n-1) \int \sinh^{n-2} x \, dx - (n-1) \int \sinh^n x \, dx \end{aligned}$$

Giving $I_n = \sinh^{n-1} x \cosh x - (n-1)I_{n-2} - (n-1)I_n$

So $nI_n = \sinh^{n-1} x \cosh x - (n-1)I_{n-2}$, $n \geq 2$ *

$$\text{b i } I_5 = \frac{1}{5} \sinh^4 x \cosh x - \frac{4}{5} I_3 \quad \leftarrow \text{using * with } n=5$$

$$\begin{aligned} &= \frac{1}{5} \sinh^4 x \cosh x - \frac{4}{5} \left\{ \frac{1}{3} \sinh^2 x \cosh x - \frac{2}{3} I_1 \right\} \\ &= \frac{1}{5} \sinh^4 x \cosh x - \frac{4}{15} \sinh^2 x \cosh x + \frac{8}{15} \int \sinh x \, dx \\ &= \frac{1}{5} \sinh^4 x \cosh x - \frac{4}{15} \sinh^2 x \cosh x + \frac{8}{15} \cosh x + C \end{aligned}$$

$$\text{When } x = \ln 3, \sinh x = \frac{e^{\ln 3} - e^{-\ln 3}}{2} = \frac{3 - \frac{1}{3}}{2} = \frac{4}{3}, \cosh x = \frac{e^{\ln 3} + e^{-\ln 3}}{2} = \frac{3 + \frac{1}{3}}{2} = \frac{5}{3}$$

When $x = 0$, $\sinh x = 0$, $\cosh x = 1$

Applying the limits 0 and $\ln 3$ to the result in b

$$\begin{aligned} \int_0^{\ln 3} \sinh^5 x \, dx &= \left[\frac{1}{5} \left(\frac{4}{3} \right)^4 \left(\frac{5}{3} \right) - \frac{4}{15} \left(\frac{4}{3} \right)^2 \left(\frac{5}{3} \right) + \frac{8}{15} \left(\frac{5}{3} \right) \right] - \left[0 + 0 + \frac{8}{15} \right] \\ &= \frac{752}{1215} = 0.619 \text{ (3 s.f.)} \end{aligned}$$

$$\text{ii } \int \sinh^4 x \, dx = I_4 = \frac{1}{4} \sinh^3 x \cosh x - \frac{3}{4} I_2 \quad \leftarrow \text{Using * with } n=4$$

$$\begin{aligned} &= \frac{1}{4} \sinh^3 x \cosh x - \frac{3}{4} \left\{ \frac{1}{2} \sinh x \cosh x - \frac{1}{2} I_0 \right\} \\ &= \frac{1}{4} \sinh^3 x \cosh x - \frac{3}{8} \sinh x \cosh x + \frac{3}{8} \int 1 \, dx \\ &= \frac{1}{4} \sinh^3 x \cosh x - \frac{3}{8} \sinh x \cosh x + \frac{3}{8} x + C \end{aligned}$$

When $x = \operatorname{arsinh} 1$ $\sinh x = 1$, $\cosh x = \sqrt{1 + \sinh^2 x} = \sqrt{2}$

When $x = 0$ $\sinh x = 0$ $\cosh x = 1$

Applying the limits 0 and $\operatorname{arsinh} 1$ gives

$$\begin{aligned}\int_0^{\operatorname{arsinh} 1} \sinh^4 x \, dx &= \frac{1}{4}(1)^3(\sqrt{2}) - \frac{3}{8}(1)(\sqrt{2}) + \frac{3}{8} \operatorname{arsinh} 1 \\ &= \frac{\sqrt{2}}{4} - \frac{3\sqrt{2}}{8} + \frac{3}{8} \ln(1 + \sqrt{1^2 + 1}) \\ &= -\frac{\sqrt{2}}{8} + \frac{3}{8} \ln(1 + \sqrt{2}) \\ &= \frac{1}{8} \{3 \ln(1 + \sqrt{2}) - \sqrt{2}\}\end{aligned}$$

© Pearson Education Ltd 2009

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 1

Question:

Find the length of the arc of the curve with equation $y = \frac{1}{3}x^{\frac{3}{2}}$, from the origin to the point with x -coordinate 12.

Solution:

$$y = \frac{1}{3}x^{\frac{3}{2}}, \text{ so } \frac{dy}{dx} = \frac{1}{2}x^{\frac{1}{2}}$$

$$\begin{aligned} \text{Arc length} &= \int_0^{12} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^{12} \sqrt{1 + \frac{x}{4}} dx \\ &= \frac{1}{2} \int_0^{12} \sqrt{4+x} dx \\ &= \frac{1}{2} \left[\frac{2}{3} (4+x)^{\frac{3}{2}} \right]_0^{12} \\ &= \frac{1}{3} \left[16^{\frac{3}{2}} - 4^{\frac{3}{2}} \right] \\ &= \frac{1}{3} [64 - 8] \\ &= \frac{56}{3} \text{ or } 18\frac{2}{3} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 2

Question:

The curve C has equation $y = \ln \cos x$. Find the length of the arc of C between the points with x -coordinates 0 and $\frac{\pi}{3}$.

Solution:

$$y = \ln \cos x, \text{ so } \frac{dy}{dx} = \frac{-\sin x}{\cos x} = -\tan x$$

$$\begin{aligned} \text{Arc length} &= \int_0^{\frac{\pi}{3}} \sqrt{1 + \tan^2 x} \, dx = \int_0^{\frac{\pi}{3}} \sec x \, dx \\ &= \left[\ln(\sec x + \tan x) \right]_0^{\frac{\pi}{3}} \\ &= \ln(2 + \sqrt{3}) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 3

Question:

Find the length of the arc on the catenary, with equation $y = 2 \cosh\left(\frac{x}{2}\right)$, between the points with x -coordinates 0 and $\ln 4$.

Solution:

$$y = 2 \cosh\left(\frac{x}{2}\right), \text{ so } \frac{dy}{dx} = \sinh\left(\frac{x}{2}\right)$$

$$\text{arc length} = \int_0^{\ln 4} \sqrt{1 + \sinh^2\left(\frac{x}{2}\right)} dx$$

$$= \int_0^{\ln 4} \cosh\left(\frac{x}{2}\right) dx$$

$$= \left[2 \sinh\left(\frac{x}{2}\right) \right]_0^{\ln 4}$$

$$= 2 \frac{e^{\frac{\ln 4}{2}} - e^{-\frac{\ln 4}{2}}}{2}$$

$$= e^{\ln 2} - e^{-\ln 2}$$

$$= 2 - \frac{1}{2} = \frac{3}{2}$$

As $\ln 4 = \ln 2^2 = 2 \ln 2$

As $e^{\ln k} = k$, $e^{-\ln k} = e^{\ln k^{-1}} = k^{-1}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 4

Question:

Find the length of the arc of the curve with equation $y^2 = \frac{4}{9}x^3$, from the origin to the point $(3, 2\sqrt{3})$.

Solution:

$$y^2 = \frac{4}{9}x^3, \text{ so } 2y \frac{dy}{dx} = \frac{4}{3}x^2 \Rightarrow \frac{dy}{dx} = \frac{2x^2}{3y} = \pm \frac{x^2}{\frac{3}{2}y} = \pm \sqrt{x} \leftarrow \begin{array}{|l} \text{The arc in question is above} \\ \text{the } x\text{-axis.} \end{array}$$

$$\begin{aligned} \text{arc length} &= \int_0^3 \sqrt{1+x} \, dx \\ &= \left[\frac{2}{3}(1+x)^{\frac{3}{2}} \right]_0^3 \\ &= \frac{2}{3}[8-1] = 4\frac{2}{3} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 5

Question:

The curve C has equation $y = \frac{1}{2} \sinh^2 2x$. Find the length of the arc on C from the origin to the point whose x -coordinate is 1, giving your answer to 3 significant figures.

Solution:

$$y = \frac{1}{2} \sinh^2 2x, \text{ so } \frac{dy}{dx} = 2 \sinh 2x \cosh 2x = \sinh 4x$$

$$\begin{aligned} \text{So arc length} &= \int_0^1 \sqrt{1 + \sinh^2 4x} dx \\ &= \int_0^1 \cosh 4x dx \\ &= \frac{1}{4} [\sinh 4x]_0^1 \\ &= \frac{1}{4} \sinh 4 = 6.82 \quad (3 \text{ s.f.}) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 6

Question:

The curve C has equation $y = \frac{1}{4}(2x^2 - \ln x)$, $x > 0$. The points A and B on C have x -coordinates 1 and 2 respectively. Show that the length of the arc from A to B is $\frac{1}{4}(6 + \ln 2)$.

Solution:

$$y = \frac{1}{4}(2x^2 - \ln x), \text{ so } \frac{dy}{dx} = x - \frac{1}{4x}$$

$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + x^2 - \frac{1}{2} + \frac{1}{16x^2} = x^2 + \frac{1}{2} + \frac{1}{16x^2} = \left(x + \frac{1}{4x}\right)^2$$

$$\begin{aligned} \text{So arc length} &= \int_1^2 \left(x + \frac{1}{4x}\right) dx \\ &= \left[\frac{x^2}{2} + \frac{1}{4}\ln x\right]_1^2 \\ &= \left[2 + \frac{1}{4}\ln 2\right] - \left[\frac{1}{2}\right] \\ &= \frac{1}{4}(6 + \ln 2) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 7

Question:

Find the length of the arc on the curve $y = 2\text{arcosh}\left(\frac{x}{2}\right)$, from the point at which the curve crosses the x -axis to the point with x -coordinate $\frac{5}{2}$. Compare your answer with that in Example 25 and explain the relationship.

Solution:

$$y = 2\text{arcosh}\left(\frac{x}{2}\right), \text{ so } \frac{dy}{dx} = 2 \times \frac{1}{2} \frac{1}{\sqrt{\left(\frac{x}{2}\right)^2 - 1}} = \frac{2}{\sqrt{x^2 - 4}}$$

$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{4}{x^2 - 4} = \frac{x^2}{x^2 - 4}$$

The curve crosses the x -axis at $x = 2$,

$$\begin{aligned} \text{So arc length} &= \int_2^{\frac{5}{2}} x(x^2 - 4)^{-\frac{1}{2}} dx \\ &= \left[\sqrt{x^2 - 4} \right]_2^{\frac{5}{2}} \\ &= 1.5 \end{aligned}$$

Eliminating t from the two equations in Example 25, you find that the Cartesian equation is $\frac{x}{2} = \cosh\left(\frac{y}{2}\right)$. For $t \geq 1$, the curve is $y = 2\text{arcosh}\left(\frac{x}{2}\right)$. The limits in both questions correspond, and so they are essentially the same question.

[For $0 < t < 1$, the reflection of $y = 2\text{arcosh}\left(\frac{x}{2}\right)$ in the x -axis is generated.]

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 8


Question:

The line $y=4$ intersects the parabola with equation $y=x^2$ at the points A and B . Find the length of the arc of the parabola from A to B .

Solution:

The line $y=4$ intersects the parabola with equation $y=x^2$ where $x=-2$ and $x=+2$.

$$\begin{aligned} \text{Using symmetry arc length} &= 2 \int_0^2 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\ &= 2 \int_0^2 \sqrt{1 + 4x^2} dx \end{aligned}$$



Using the substitution $2x = \sinh u$, so that $2dx = \cosh u du$,

$$\begin{aligned} \text{arc length} &= \int_0^{\text{arsinh } 4} \sqrt{1 + \sinh^2 u} \cosh u du \\ &= \int_0^{\text{arsinh } 4} \cosh^2 u du \\ &= \int_0^{\text{arsinh } 4} \frac{(1 + \cosh 2u)}{2} du \\ &= \frac{1}{2} \left[u + \frac{1}{2} \sinh 2u \right]_0^{\text{arsinh } 4} \\ &= \frac{1}{2} \left[u + \sinh u \cosh u \right]_0^{\text{arsinh } 4} \\ &= \frac{1}{2} \text{arsinh } 4 + \frac{1}{2} (4\sqrt{1+16}) \leftarrow \text{Using } \cosh u = \sqrt{1 + \sinh^2 u} \text{ and } \sinh u = 4 \\ &= \frac{1}{2} \text{arsinh } 4 + 2\sqrt{17} \\ &= \frac{1}{2} \ln (4 + \sqrt{17}) + 2\sqrt{17} \leftarrow \text{Using } \text{arsinh } x = \ln \{x + \sqrt{(1+x^2)}\} \\ &= 9.29 \text{ (3 s.f.)} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 9

Question:

The circle C has parametric equations $x = r \cos \theta$, $y = r \sin \theta$. Use the formula for arc length on page 79 for to show that the length of the circumference is $2\pi r$.

Solution:

$$\text{As } x = r \cos \theta, y = r \sin \theta, \frac{dx}{d\theta} = -r \sin \theta, \frac{dy}{d\theta} = r \cos \theta$$

$$\text{So } \left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 = r^2 (\cos^2 \theta + \sin^2 \theta) = r^2$$

$$\begin{aligned} \text{The circumference of the circle} &= 4 \int_0^{\frac{\pi}{2}} r \, d\theta \\ &= 4r \left[\theta\right]_0^{\frac{\pi}{2}} \\ &= 2\pi r \end{aligned}$$

Using symmetry.

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics


Integration

Exercise G, Question 10

Question:

The diagram shows the astroid, with parametric equations
 $x = 2a \cos^3 t, y = 2a \sin^3 t, 0 \leq t < 2\pi$.

Find the length of the arc of the curve AB , and hence find the total length of the curve.



Solution:

$$x = 2a \cos^3 t, y = 2a \sin^3 t, \text{ so } \frac{dx}{dt} = -6a \cos^2 t \sin t, \frac{dy}{dt} = 6a \sin^2 t \cos t,$$

$$\begin{aligned} \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 &= 36a^2 (\cos^4 t \sin^2 t + \sin^4 t \cos^2 t) = 36a^2 \sin^2 t \cos^2 t (\cos^2 t + \sin^2 t) \\ &= 36a^2 \sin^2 t \cos^2 t \end{aligned}$$

$$\text{At } A, t = 0, \text{ at } B, t = \frac{\pi}{2},$$

$$\begin{aligned} \text{so arc length } AB &= \int_0^{\frac{\pi}{2}} 6a \sin t \cos t \, dt \\ &= 3a \int_0^{\frac{\pi}{2}} \sin 2t \, dt \\ &= \frac{3}{2}a [-\cos 2t]_0^{\frac{\pi}{2}} \\ &= \frac{3}{2}a [1 - (-1)] \\ &= 3a \end{aligned}$$

$$\text{Total length of curve} = 4 \times 3a = 12a \text{ (symmetry)}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 11

Question:

Calculate the length of the arc on the curve with parametric equations $x = \tanh u$, $y = \operatorname{sech} u$, between the points with parameters $u = 0$ and $u = 1$.

Solution:

$$x = \tanh u, y = \operatorname{sech} u, \text{ so } \frac{dx}{du} = \operatorname{sech}^2 u, \frac{dy}{du} = -\operatorname{sech} u \tanh u,$$

$$\left(\frac{dx}{du}\right)^2 + \left(\frac{dy}{du}\right)^2 = \operatorname{sech}^4 u + \operatorname{sech}^2 u \tanh^2 u = \operatorname{sech}^2 u (\operatorname{sech}^2 u + \tanh^2 u) = \operatorname{sech}^2 u$$

So arc length = $\int_0^1 \operatorname{sech} u \, du$ ← See Example 7.

$$= \int_0^1 \frac{2}{e^u + e^{-u}} \, du$$

$$= \int_0^1 \frac{2e^u}{(e^u)^2 + 1} \, du$$

$$= 2 \left[\arctan(e^u) \right]_0^1$$

$$= 2 \arctan(e) - \frac{\pi}{2} \text{ or } 0.866 \text{ (3 s.f.)}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 12

Question:

The cycloid has parametric equations $x = a(\theta + \sin \theta)$, $y = a(1 - \cos \theta)$. Find the length of the arc from $\theta = 0$ to $\theta = \pi$.

Solution:

$$\text{As } x = a(\theta + \sin \theta), y = a(1 - \cos \theta), \frac{dx}{d\theta} = a(1 + \cos \theta), \frac{dy}{d\theta} = a \sin \theta$$

$$\begin{aligned} \left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 &= a^2(1 + 2\cos\theta + \cos^2\theta + \sin^2\theta) \\ &= a^2(2 + 2\cos\theta) \end{aligned}$$

$$= 4a^2 \cos^2\left(\frac{\theta}{2}\right)$$

Using $\cos 2A = 2\cos^2 A - 1$ with $A = \left(\frac{\theta}{2}\right)$

$$\text{So arc length} = 2a \int_0^\pi \cos\left(\frac{\theta}{2}\right) d\theta$$

$$= 4a \left[\sin\left(\frac{\theta}{2}\right) \right]_0^\pi$$

$$= 4a$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 13

Question:

Show that the length of the arc, between the points with parameters $t = 0$ and $t = \frac{\pi}{3}$ on the curve defined by the equations $x = t + \sin t, y = 1 - \cos t$, is 2.

Solution:

$$x = t + \sin t, y = 1 - \cos t$$

$$\frac{dx}{dt} = 1 + \cos t, \frac{dy}{dt} = \sin t$$

$$\begin{aligned} \text{So } \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 &= \{(1 + 2\cos t + \cos^2 t) + (\sin^2 t)\} \\ &= 2(1 + \cos t) = 4\cos^2\left(\frac{t}{2}\right) \end{aligned}$$

$$\text{Using } s = \int_{t_A}^{t_B} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$\begin{aligned} \text{arc length} &= \int_0^{\frac{\pi}{3}} \sqrt{4\cos^2\left(\frac{t}{2}\right)} dt \\ &= 2 \int_0^{\frac{\pi}{3}} \cos\left(\frac{t}{2}\right) dt \\ &= 4 \left[\sin\left(\frac{t}{2}\right) \right]_0^{\frac{\pi}{3}} \\ &= 2 \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 14

Question:

Find the length of the arc of the curve given by the equations $x = e^t \cos t, y = e^t \sin t$,
between the points with parameters $t = 0$ and $t = \frac{\pi}{4}$.

Solution:

$$x = e^t \cos t, y = e^t \sin t$$

$$\frac{dx}{dt} = e^t (\cos t - \sin t), \frac{dy}{dt} = e^t (\sin t + \cos t)$$

$$\begin{aligned} \text{So } \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 &= (e^t)^2 \{(\cos^2 t - 2 \sin t \cos t + \sin^2 t) + (\sin^2 t + 2 \sin t \cos t + \cos^2 t)\} \\ &= 2(e^t)^2 (\sin^2 t + \cos^2 t) \\ &= 2(e^t)^2 \end{aligned}$$

$$\begin{aligned} \text{arc length} &= \int_0^{\frac{\pi}{4}} \sqrt{2(e^t)^2} dt \\ &= \sqrt{2} \int_0^{\frac{\pi}{4}} e^t dt \\ &= \sqrt{2} [e^t]_0^{\frac{\pi}{4}} \\ &= \sqrt{2} \left[e^{\frac{\pi}{4}} - 1 \right] \text{ or } 1.69 \quad (3 \text{ s.f.}) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise G, Question 15

Question:

- a Denoting the length of one complete wave of the sine curve with equation

$$y = \sqrt{3} \sin x \text{ by } L, \text{ show that } L = 4 \int_0^{\frac{\pi}{2}} \sqrt{1 + 3 \cos^2 x} \, dx.$$

- b The ellipse has parametric equations $x = \cos t, y = 2 \sin t$. Show that the length of its circumference is equal to that of the wave in a.


Solution:

a $y = \sqrt{3} \sin x$, so $\frac{dy}{dx} = \sqrt{3} \cos x$

$$\begin{aligned} \text{Using the symmetry of the sine curve } s &= 4 \int_0^{\frac{\pi}{2}} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx \\ &= 4 \int_0^{\frac{\pi}{2}} \sqrt{1 + 3 \cos^2 x} \, dx \end{aligned}$$

b $x = \cos t, y = 2 \sin t$

$$\begin{aligned} \frac{dx}{dt} &= -\sin t, \frac{dy}{dt} = 2 \cos t \\ \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 &= \sin^2 t + 4 \cos^2 t \\ &= 1 - \cos^2 t + 4 \cos^2 t \\ &= 1 + 3 \cos^2 t \end{aligned}$$



From the diagram, at A, $t = 0$,

at B, $t = \frac{\pi}{2}$,

so using the symmetry of the ellipse, the length of the circumference is

$$4 \int_0^{\frac{\pi}{2}} \sqrt{1 + 3 \cos^2 t} \, dt, \text{ equal to that of the sine curve in a}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 1

Question:

- a The section of the line $y = \frac{3}{4}x$ between points with x -coordinates 4 and 8 is rotated completely about the x -axis. Use integration to find the area of the surface generated.
- b The same section of line is rotated completely about the y -axis. Show that the area of the surface generated is 60π .

Solution:

$$\text{a } y = \frac{3}{4}x \Rightarrow \frac{dy}{dx} = \frac{3}{4} \Rightarrow 1 + \left(\frac{dy}{dx}\right)^2 = \frac{25}{16}$$

$$\text{Surface area} = \int_4^8 2\pi \left(\frac{3}{4}x\right) \left(\frac{5}{4}\right) dx$$

$$= \frac{15}{8}\pi \int_4^8 x dx$$

$$= \frac{15}{8}\pi \left[\frac{x^2}{2}\right]_4^8 = 45\pi$$

Using $\int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$

- b Rotating about the y -axis:

$$\text{From the work in a } 1 + \left(\frac{dx}{dy}\right)^2 = 1 + \frac{16}{9} = \frac{25}{9}$$

As integration is w.r.t. y , the integrand must be in terms of y
The limits for y are 3 (when $x = 4$) and 6 (when $x = 8$),

$$\text{so area of surface is } \int_3^6 2\pi \left(\frac{4}{3}y\right) \left(\frac{5}{3}\right) dy,$$

$$= \frac{40}{9}\pi \left[\frac{y^2}{2}\right]_3^6$$

$$= \frac{40 \times 27}{9 \times 2}\pi = 60\pi$$

Although it is quicker to use

$$\int_a^b 2\pi x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx,$$

$$\text{here } \int_{y_1}^{y_2} 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

is used to give an example of its use.

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 2

Question:

The arc of the curve $y = x^3$, between the origin and the point (1, 1), is rotated through 4 right-angles about the x -axis. Find the area of the surface generated.

Solution:

$$y = x^3 \text{ so } \frac{dy}{dx} = 3x^2$$

$$\text{Using } \int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx,$$

$$\begin{aligned} \text{the area of the surface is } & \int_0^1 2\pi x^3 \sqrt{1 + 9x^4} dx \\ &= \frac{2\pi}{36} \int_0^1 36x^3 \sqrt{1 + 9x^4} dx \\ &= \frac{2\pi}{36} \left[\frac{2}{3} (1 + 9x^4)^{\frac{3}{2}} \right]_0^1 \\ &= \frac{\pi}{27} [10\sqrt{10} - 1] \quad (3.56, 3 \text{ s.f.}) \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 3

Question:

The arc of the curve $y = \frac{1}{2}x^2$, between the origin and the point (2, 2), is rotated through 4 right-angles about the y -axis. Find the area of the surface generated.

Solution:

$$y = \frac{1}{2}x^2, \text{ so } \frac{dy}{dx} = x$$

$$\text{Using } \int_{x_1}^{x_2} 2\pi x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx,$$

$$\begin{aligned} \text{the area of the surface is } & \int_0^2 2\pi x \sqrt{1 + x^2} dx \\ &= \pi \int_0^2 2x \sqrt{1 + x^2} dx \\ &= \pi \left[\frac{2}{3} (1 + x^2)^{\frac{3}{2}} \right]_0^2 \\ &= \frac{2\pi}{3} [5\sqrt{5} - 1] \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 4

Question:

The points A and B , in the first quadrant, on the curve $y^2 = 16x$ have x -coordinates 5 and 12 respectively. Find, in terms π , the area of the surface generated when the arc AB is rotated completely about the x -axis.

Solution:

$$y^2 = 16x \text{ so } 2y \frac{dy}{dx} = 16 \Rightarrow \frac{dy}{dx} = \frac{8}{y}$$

$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{64}{y^2} = 1 + \frac{4}{x} = \frac{x+4}{x}$$

$$\text{Using } \int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx,$$

$$\begin{aligned} \text{the area of the surface is } & \int_5^{12} 2\pi 4\sqrt{x} \sqrt{\frac{4+x}{x}} dx \\ & = 8\pi \int_5^{12} \sqrt{4+x} dx \\ & = 8\pi \left[\frac{2}{3} (4+x)^{\frac{3}{2}} \right]_5^{12} \\ & = \frac{16\pi}{3} [37] \\ & = \frac{592\pi}{3} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 5

Question:

The curve C has equation $y = \cosh x$. The arc s on C , has end points $(0, 1)$ and $(1, \cosh 1)$.

- Find the area of the surface generated when s is rotated completely about the x -axis.
- Show that the area of the surface generated when s is rotated completely about the

y -axis is $2\pi \left(\frac{e-1}{e} \right)$.

Solution:

$$y = \cosh x, \text{ so } \frac{dy}{dx} = \sinh x$$

$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \sinh^2 x = \cosh^2 x$$

a Using $\int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$,

the area of the surface is $\int_0^1 2\pi \cosh^2 x dx$

$$= \pi \int_0^1 (\cosh 2x + 1) dx$$

$$= \pi \left[\frac{\sinh 2x}{2} + x \right]_0^1$$

$$= \pi [\sinh x \cosh x + x]_0^1$$

$$= \pi [\sinh 1 \cosh 1 + 1]$$

$$= 8.84 \text{ (3 s.f.)}$$

b Using $\int_{x_1}^{x_2} 2\pi x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$,

the area of the surface is $\int_0^1 2\pi x \cosh x dx$

$$= 2\pi \left\{ [x \sinh x]_0^1 - \int_0^1 \sinh x dx \right\}$$

$$= 2\pi \left\{ \sinh 1 - [\cosh x]_0^1 \right\}$$

$$= 2\pi \left\{ \sinh 1 - \cosh 1 + 1 \right\}$$

$$= 2\pi \left\{ \frac{1}{2} \left(e - \frac{1}{e} - e - \frac{1}{e} \right) + 1 \right\}$$

$$= 2\pi \left(1 - \frac{1}{e} \right)$$

$$= 2\pi \left(\frac{e-1}{e} \right)$$

Using integration by parts

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 6

Question:

The curve C has equation $y = \frac{1}{2x} + \frac{x^3}{6}$.

a Show that $\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \frac{1}{2}\left(x^2 + \frac{1}{x^2}\right)$.

The arc of the curve between points with x -coordinates 1 and 3 is rotated completely about the x -axis.

b Find the area of the surface generated.

Solution:

a $y = \frac{1}{2x} + \frac{x^3}{6}$, so $\frac{dy}{dx} = -\frac{1}{2x^2} + \frac{x^2}{2} = \frac{1}{2}\left(x^2 - \frac{1}{x^2}\right)$

$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{1}{4}\left(x^4 - 2 + \frac{1}{x^4}\right) = \frac{1}{4}\left(x^4 + 2 + \frac{1}{x^4}\right) = \frac{1}{4}\left(x^2 + \frac{1}{x^2}\right)^2$$

$$\text{So } \sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \frac{1}{2}\left(x^2 + \frac{1}{x^2}\right)$$

b Using $\int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$,

the area of the surface is $\pi \int_1^3 \left(\frac{1}{2x} + \frac{x^3}{6}\right) \left(x^2 + \frac{1}{x^2}\right) dx$

$$= \pi \int_1^3 \left(\frac{2x}{3} + \frac{x^5}{6} + \frac{1}{2x^3}\right) dx$$

$$= \pi \left[\frac{x^2}{3} + \frac{x^6}{36} - \frac{1}{4x^2}\right]_1^3$$

$$= 23\frac{1}{9}\pi = 72.6 \text{ (3 s.f.)}$$

Solutionbank FP3


Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 7

Question:

The diagram shows part of the curve with equation $x^{\frac{2}{3}} + y^{\frac{2}{3}} = 4$. Find the area of the surface generated when this arc is rotated completely about the y -axis.



Solution:

$$x^{\frac{2}{3}} + y^{\frac{2}{3}} = 4, \text{ so } \frac{2}{3}x^{-\frac{1}{3}} + \frac{2}{3}y^{-\frac{1}{3}} \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{x^{-\frac{1}{3}}}{y^{-\frac{1}{3}}} = -\frac{y^{\frac{1}{3}}}{x^{\frac{2}{3}}}$$

$$\text{So } 1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{y^{\frac{2}{3}}}{x^{\frac{4}{3}}} = \frac{x^{\frac{2}{3}} + y^{\frac{2}{3}}}{x^{\frac{2}{3}}} = \frac{4}{x^{\frac{2}{3}}}$$

$$\text{Using } \int_{x_1}^{x_2} 2\pi x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx,$$

$$\begin{aligned} \text{the area of the surface is } & 2\pi \int_0^8 x \left(\frac{2}{x^{\frac{2}{3}}}\right) dx \\ & = 2\pi \int_0^8 2x^{\frac{2}{3}} dx \\ & = 2\pi \left[\frac{6}{5} x^{\frac{5}{3}} \right]_0^8 \\ & = \frac{12\pi}{5} [32] \\ & = \frac{384\pi}{5} = 241 \text{ (3 s.f.)} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 8

Question:

- a The arc of the circle with equation $x^2 + y^2 = R^2$, between the points $(-R, 0)$ and $(R, 0)$, is rotated through 2π radians about the x -axis. Use integration to find the surface area of the sphere S formed.
- b The axis of a cylinder C of radius R is the x -axis. Show that the areas of the surface of S and C , contained between planes with equations $x = a$ and $x = b$, where $a < b < R$, are equal.

Solution:

a $x^2 + y^2 = R^2$, so $2x + 2y \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{x}{y}$

So $1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{x^2}{y^2} = \frac{x^2 + y^2}{y^2} = \frac{R^2}{y^2}$

Using $\int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$,

the area of the surface of the sphere is $2\pi \int_{-R}^R y \left(\frac{R}{y}\right) dx$

$$= 4\pi \int_0^R R dx$$

Using the symmetry

$$= 4\pi R [x]_0^R$$

$$= 4\pi R^2$$


b The required area is $2\pi \int_a^b y \left(\frac{R}{y}\right) dx$

see diagram

$$= 2\pi \int_a^b R dx$$

$$= 2\pi R(b-a)$$

This is the same area as that of a cylinder of radius R and height $(b-a)$.



Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 9

Question:

The finite arc of the parabola with parametric equations $x = at^2$, $y = 2at$, where a is a positive constant, cut off by the line $x = 4a$, is rotated through 180° about the x -axis.

Show that the area of the surface generated is $\frac{8}{3}\pi a^2(5\sqrt{5}-1)$.

Solution:

$$x = at^2, y = 2at, \text{ so } \frac{dx}{dt} = 2at, \frac{dy}{dt} = 2a$$

$$\text{So } \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 4a^2t^2 + 4a^2 = 4a^2(1+t^2)$$

$x = 4a$ when $t = \pm 2$ (See diagram.)

A rotation of π radians gives a surface which would be found by rotating the section $y \geq 0$, i.e.

$t = 0$ to $t = 2$ through 2π radians.


$$\text{Using } \int_{t_1}^{t_2} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt,$$

$$\text{the area of the surface is } 2\pi \int_0^2 4a^2t\sqrt{1+t^2} dt$$

$$= 8\pi a^2 \left[\frac{1}{3}(1+t^2)^{\frac{3}{2}} \right]_0^2$$

$$= \frac{8}{3}\pi a^2 \left[5^{\frac{3}{2}} - 1 \right]$$

$$= \frac{8}{3}\pi a^2 (5\sqrt{5} - 1)$$



Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 10

Question:

The arc, in the first quadrant, of the curve with parametric equations
 $x = \operatorname{sech} t, y = \tanh t$, between the points where $t = 0$ and $t = \ln 2$, is rotated
 completely about the x -axis. Show that the area of the surface generated is $\frac{2\pi}{5}$.

Solution:

$$x = \operatorname{sech} t, y = \tanh t, \text{ so } \frac{dx}{dt} = -\operatorname{sech} t \tanh t, \frac{dy}{dt} = \operatorname{sech}^2 t$$

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \operatorname{sech}^2 t \tanh^2 t + \operatorname{sech}^4 t = \operatorname{sech}^2 t (\tanh^2 t + \operatorname{sech}^2 t) = \operatorname{sech}^2 t$$

$$\text{Using } \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt,$$

$$\text{the area of the surface is } 2\pi \int_0^{\ln 2} \tanh t \operatorname{sech} t dt$$

$$= 2\pi \left[-\operatorname{sech} t \right]_0^{\ln 2}$$

$$= 2\pi \left[-\frac{2}{e^t + e^{-t}} \right]_0^{\ln 2}$$

$$= \frac{2\pi}{5} \left[\frac{-2}{2.5} + 1 \right]$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 11

Question:

The arc of the curve given by $x = 3t^2, y = 2t^3$, from $t = 0$ and $t = 2$, is completely rotated about the y -axis.

- a Show that the area of the surface generated can be expressed as $36\pi \int_0^2 t^3 \sqrt{1+t^2} dt$.
- b Using integration by parts, find the exact value of this area.

Solution:

a $x = 3t^2, y = 2t^3$, so $\frac{dx}{dt} = 6t, \frac{dy}{dt} = 6t^2$

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 36t^2(t^2 + 1)$$

Using $\int_{t_1}^{t_2} 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$,

the area of the surface is $2\pi \int_0^2 3t^2 \times 6t \sqrt{1+t^2} dt$

$$= 36\pi \int_0^2 t^3 \sqrt{1+t^2} dt$$

b Let $u = t^2, \frac{du}{dt} = 2t \sqrt{1+t^2}$

So $\frac{du}{dt} = 2t, v = \frac{1}{3}(1+t^2)^{\frac{3}{2}}$

$$\begin{aligned} 36\pi \int_0^2 t^3 (t\sqrt{1+t^2}) dt &= 36\pi \left\{ \left[\frac{1}{3} t^2 (1+t^2)^{\frac{3}{2}} \right]_0^2 - \int_0^2 \frac{2}{3} t (1+t^2)^{\frac{3}{2}} dt \right\} \\ &= 12\pi \left[t^2 (1+t^2)^{\frac{3}{2}} - \frac{2}{5} (1+t^2)^{\frac{5}{2}} \right]_0^2 \\ &= 12\pi \left[4(5\sqrt{5}) - \frac{2}{5}(25\sqrt{5}) + \frac{2}{5} \right] \\ &= 12\pi \left[10\sqrt{5} + \frac{2}{5} \right] \\ &= \frac{24\pi}{5} [25\sqrt{5} + 1] \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 12

Question:

The arc of the curve with parametric equations $x = t^2, y = t - \frac{1}{3}t^3$, between the points where $t = 0$ and $t = 1$, is rotated through 360° about the x -axis. Calculate the area of the surface generated.

Solution:

$$x = t^2, y = t - \frac{1}{3}t^3, \text{ so } \frac{dx}{dt} = 2t, \frac{dy}{dt} = 1 - t^2$$

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 4t^2 + 1 - 2t^2 + t^4 = (1 + t^2)^2$$

$$\text{Using } \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt,$$

$$\text{the area of the surface is } 2\pi \int_0^1 \left(t - \frac{1}{3}t^3\right)(1 + t^2) dt$$

$$= 2\pi \int_0^1 \left(t + \frac{2}{3}t^3 - \frac{1}{3}t^5\right) dt$$

$$= 2\pi \left[\frac{t^2}{2} + \frac{t^4}{6} - \frac{t^6}{18} \right]_0^1$$

$$= \frac{11\pi}{9}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 13

Question:

The astroid C has parametric equations $x = a \cos^3 t, y = a \sin^3 t$, where a is a positive constant. The arc of C , between $t = \frac{\pi}{6}$ and $t = \frac{\pi}{2}$ is rotated through 2π radians about the x -axis. Find the area of the surface of revolution formed.

Solution:

$$x = a \cos^3 t, y = a \sin^3 t, \text{ so } \frac{dx}{dt} = -3a \cos^2 t \sin t, \frac{dy}{dt} = 3a \sin^2 t \cos t$$

$$\begin{aligned} \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 &= 9a^2 (\cos^4 t \sin^2 t + \sin^4 t \cos^2 t) \\ &= 9a^2 \sin^2 t \cos^2 t (\cos^2 t + \sin^2 t) \\ &= 9a^2 \sin^2 t \cos^2 t \end{aligned}$$

$$\text{Using } \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt,$$

$$\text{the area of the surface is } 2\pi \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} a \sin^3 t (3a \sin t \cos t) dt$$

$$= 6\pi a^2 \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \sin^4 t \cos t dt$$

$$= 6\pi a^2 \left[\frac{1}{5} \sin^5 t \right]_{\frac{\pi}{6}}^{\frac{\pi}{2}}$$

$$= \frac{6\pi a^2}{5} \left[1 - \frac{1}{32} \right]$$

$$= \frac{93\pi a^2}{80}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise H, Question 14

Question:

The part of the curve $y = e^x$, between $(0, 1)$ and $(\ln 2, 2)$, is rotated completely about the x -axis. Show that the area of the surface generated is $\pi(\operatorname{arsinh} 2 - \operatorname{arsinh} 1 + 2\sqrt{5} - \sqrt{2})$.

Solution:

$$y = e^x, \frac{dy}{dx} = e^x$$

$$\text{Using } \int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx,$$

$$\text{the area of the surface is } 2\pi \int_0^{\ln 2} e^x \sqrt{1 + e^{2x}} dx$$

Make the substitution $e^x = \sinh u$, so $e^x dx = \cosh u du$

Limits: when $x = \ln 2, u = \operatorname{arsinh} e^{\ln 2} = \operatorname{arsinh} 2$

when $x = 0, u = \operatorname{arsinh} e^0 = \operatorname{arsinh} 1$

$$\text{Then the area of the surface is } 2\pi \int_{\operatorname{arsinh} 1}^{\operatorname{arsinh} 2} \cosh^2 u du$$

$$= \pi \int_{\operatorname{arsinh} 1}^{\operatorname{arsinh} 2} (1 + \cosh 2u) du$$

$$= \pi \left[u + \frac{\sinh 2u}{2} \right]_{\operatorname{arsinh} 1}^{\operatorname{arsinh} 2}$$

$$= \pi \left[u + \sinh u \cosh u \right]_{\operatorname{arsinh} 1}^{\operatorname{arsinh} 2}$$

$$= \pi \left[\operatorname{arsinh} 2 + 2\sqrt{5} - \left(\operatorname{arsinh} 1 + (1)(\sqrt{2}) \right) \right]_{\operatorname{arsinh} 1}$$

$$= \pi (\operatorname{arsinh} 2 - \operatorname{arsinh} 1 + 2\sqrt{5} - \sqrt{2})$$

$\cosh u = \sqrt{1 + \sinh^2 u}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 1

Question:

Show that the volume of the solid generated when the finite region enclosed by the curve with equation $y = \tanh x$, the line $x = 1$ and the x -axis is rotated through 2π radians about the x -axis is $\frac{2\pi}{1+e^2}$. [E]

Solution:

$$\begin{aligned}\text{Volume} &= \pi \int_0^1 y^2 dx = \pi \int_0^1 \tanh^2 x dx \\ &= \pi \int_0^1 (1 - \operatorname{sech}^2 x) dx \\ &= \pi [x - \tanh x]_0^1 \\ &= \pi (1 - \tanh 1) \\ &= \pi \left(1 - \frac{e^2 - 1}{e^2 + 1} \right) \\ &= \frac{2\pi}{1 + e^2}\end{aligned}$$

$\tanh 1 = \frac{e - e^{-1}}{e + e^{-1}} = \frac{e^2 - 1}{e^2 + 1}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 2

Question:

$$4x^2 + 4x + 17 \equiv (ax + b)^2 + c, a > 0.$$

a Find the values of a , b and c .

b Find the exact value of $\int_{-0.5}^{1.5} \frac{1}{4x^2 + 4x + 17} dx$ [E]

Solution:

$$4x^2 + 4x + 17 \equiv (ax + b)^2 + c, \quad a > 0$$

a $4x^2 + 4x + 17 \equiv (2x + b)^2 + c \quad a = 2$

$$\equiv 4x^2 + 4bx + b^2 + c$$

Comparing coefficient of x : $b = 1$

Comparing constant term: $17 = 1 + c \Rightarrow c = 16$

b Using a, $\int \frac{1}{4x^2 + 4x + 17} dx = \int \frac{1}{(2x + 1)^2 + 16} dx$

Let $2x + 1 = 4 \tan \theta$, then $2 dx = 4 \sec^2 \theta d\theta$

and $\int \frac{1}{(2x + 1)^2 + 16} dx = \int \frac{2 \sec^2 \theta}{16 \tan^2 \theta + 16} d\theta$

$$= \int \frac{2 \sec^2 \theta}{16 \sec^2 \theta} d\theta$$

$$= \frac{1}{8} \theta + C$$

$$= \frac{1}{8} \arctan \left(\frac{2x + 1}{4} \right) + C$$

So $\int_{-0.5}^{1.5} \frac{1}{4x^2 + 4x + 17} dx = \frac{1}{8} [\arctan 1 - \arctan 0]$

$$= \frac{\pi}{32}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 3

Question:

Find the following.

a $\int \sinh 4x \cosh 6x \, dx$

b $\int \frac{\operatorname{sech} x \tanh x}{1 + 2\operatorname{sech} x} \, dx$

c $\int e^x \sinh x \, dx$

Solution:

a Using the definitions of $\sinh 4x$ and $\cosh 6x$

$$\begin{aligned} \int \sinh 4x \cosh 6x \, dx &= \int \left(\frac{e^{4x} - e^{-4x}}{2} \right) \left(\frac{e^{6x} + e^{-6x}}{2} \right) dx \\ &= \frac{1}{4} \int (e^{10x} + e^{-2x} - e^{2x} - e^{-10x}) dx \end{aligned}$$

You could use hyperbolic identities to split up into a difference of two sinhs.

$$= \frac{1}{4} \left\{ \frac{e^{10x}}{10} + \frac{e^{-2x}}{-2} - \frac{e^{2x}}{2} - \frac{e^{-10x}}{-10} \right\} + C$$

$$= \frac{1}{4} \left\{ \frac{e^{10x}}{10} + \frac{e^{-10x}}{10} - \frac{e^{2x}}{2} - \frac{e^{-2x}}{2} \right\} + C$$

$$= \frac{1}{20} \cosh 10x - \frac{1}{4} \cosh 2x + C$$

$$\text{as } \cosh ax = \frac{e^{ax} + e^{-ax}}{2}$$

b $\int \frac{\operatorname{sech} x \tanh x}{1 + 2\operatorname{sech} x} \, dx = -\frac{1}{2} \int \frac{-2\operatorname{sech} x \tanh x}{1 + 2\operatorname{sech} x} \, dx = -\frac{1}{2} \ln(1 + 2\operatorname{sech} x) + C$

c You cannot use by parts for $\int e^x \sinh x \, dx$

Using the definition of $\sinh x$

$$\begin{aligned} \int e^x \sinh x \, dx &= \int e^x \left(\frac{e^x - e^{-x}}{2} \right) dx \\ &= \frac{1}{2} \int (e^{2x} - 1) dx \\ &= \frac{1}{2} \left(\frac{1}{2} e^{2x} - x \right) + C \\ &= \frac{1}{4} e^{2x} - \frac{1}{2} x + C \end{aligned}$$


Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 4

Question:



The diagram shows the cross-section R of an artificial ski slope. The slope is modelled by the curve with equation

$$y = \frac{10}{\sqrt{4x^2 + 9}}, 0 \leq x \leq 5.$$

Given that 1 unit on each axis represents 10 metres, use integration to calculate the area R . Show your method clearly and give your answer to 2 significant figures. [E]

Solution:

$$\begin{aligned} \text{Area under curve} &= \int_0^5 y \, dx = \int_0^5 \frac{10}{\sqrt{4x^2 + 9}} \, dx \\ &= 5 \int_0^5 \frac{1}{\sqrt{x^2 + \frac{9}{4}}} \, dx \\ &= 5 \left[\operatorname{arsinh} \left(\frac{2x}{3} \right) \right]_0^5 \\ &= 5 \operatorname{arsinh} \left(\frac{10}{3} \right) \text{ (sq. units)} \end{aligned}$$

Using $\int \frac{1}{\sqrt{x^2 + a^2}} \, dx = \operatorname{arsinh} \left(\frac{x}{a} \right)$

$$\text{'Real' area} = 5 \operatorname{arsinh} \left(\frac{10}{3} \right) \times 100 \, \text{m}^2 = 960 \text{ (2 s.f.)}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 5

Question:

a Find $\int \frac{1+2x}{1+4x^2} dx$.

b Find the exact value of

$$\int_0^{0.5} \frac{1+2x}{1+4x^2} dx.$$

Solution:

$$\begin{aligned} \text{a } \int \frac{1+2x}{1+4x^2} dx &= \int \frac{1}{1+4x^2} dx + \int \frac{2x}{1+4x^2} dx \\ &= \int \frac{1}{4\left(\frac{1}{4} + x^2\right)} dx + \frac{1}{4} \int \frac{8x}{1+4x^2} dx \\ &= \frac{1}{2} \arctan 2x + \frac{1}{4} \ln(1+4x^2) + C \end{aligned}$$

$$\text{b } \int_0^{0.5} \frac{1+2x}{1+4x^2} dx = \frac{1}{2} \arctan 1 + \frac{1}{4} \ln 2$$

Using the result from a

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 6

Question:

A rope is hung from points two points on the same horizontal level. The curve formed by the rope is modelled by the equation

$$y = 4 \cosh\left(\frac{x}{4}\right), -20 \leq x \leq 20,$$

Find the length of the rope, giving your answer to 3 significant figures.

Solution:

$$y = 4 \cosh\left(\frac{x}{4}\right), \text{ so } \frac{dy}{dx} = \frac{4}{4} \sinh\left(\frac{x}{4}\right) = \sinh\left(\frac{x}{4}\right)$$

$$\text{arc length} = \int_{-20}^{20} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$= 2 \int_0^{20} \sqrt{1 + \sinh^2\left(\frac{x}{4}\right)} dx$$

Using the symmetry of the catenary

$$= 2 \int_0^{20} \cosh\left(\frac{x}{4}\right) dx$$

$$= 2 \left[4 \sinh\left(\frac{x}{4}\right) \right]_0^{20}$$

$$= 8 \sinh 5$$

$$= 594 \text{ (3 s.f.)}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 7

Question:

Show that $\int_0^{\frac{1}{2}} \operatorname{artanh} x \, dx = \frac{1}{4} \ln\left(\frac{a}{b}\right)$, where a and b are positive integers to be found.

Solution:

$$\text{Let } u = \operatorname{artanh} x \quad \frac{dv}{dx} = 1$$

$$\text{So } \frac{du}{dx} = \frac{1}{1-x^2} \quad v = x$$

$$\text{Then } \int_0^{\frac{1}{2}} \operatorname{artanh} x \, dx = [x \operatorname{artanh} x]_0^{\frac{1}{2}} - \int_0^{\frac{1}{2}} \frac{x}{1-x^2} \, dx$$

$$= [x \operatorname{artanh} x]_0^{\frac{1}{2}} + \frac{1}{2} \int_0^{\frac{1}{2}} \frac{-2x}{1-x^2} \, dx$$

$$= \left[x \operatorname{artanh} x + \frac{1}{2} \ln(1-x^2) \right]_0^{\frac{1}{2}}$$

$$= \frac{1}{2} \operatorname{artanh}\left(\frac{1}{2}\right) + \frac{1}{2} \ln\left(\frac{3}{4}\right)$$

$$= \frac{1}{2} \left\{ \frac{1}{2} \ln\left(\frac{3}{\frac{1}{2}}\right) \right\} + \frac{1}{2} \ln\left(\frac{3}{4}\right)$$

$$= \frac{1}{4} \ln 3 + \frac{1}{2} \ln\left(\frac{3}{4}\right)$$

$$= \frac{1}{4} \left\{ \ln 3 + 2 \ln\left(\frac{3}{4}\right) \right\}$$

$$= \frac{1}{4} \left\{ \ln 3 + \ln\left(\frac{9}{16}\right) \right\}$$

$$= \frac{1}{4} \ln\left(\frac{27}{16}\right) \text{ so } a = 27 \text{ and } b = 16$$

$$\text{Using } \operatorname{artanh} x = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right)$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 8

Question:

Given that $I_n = \int_0^{\frac{\pi}{2}} x^n \cos x \, dx$,

a find the values of

i I_0 and

ii I_1 .

b show, by using integration by parts twice, that $I_n = \left(\frac{\pi}{2}\right)^n - n(n-1)I_{n-2}$, $n \geq 2$.

c Hence show that $\int_0^{\frac{\pi}{2}} x^3 \cos x \, dx = \frac{1}{8}(\pi^3 - 24\pi + 48)$.

d Evaluate $\int_0^{\frac{\pi}{2}} x^4 \cos x \, dx$, leaving your answer in terms of π .

Solution:

$$\text{a i } I_0 = \int_0^{\frac{\pi}{2}} \cos x \, dx = [\sin x]_0^{\frac{\pi}{2}} = 1$$

$$\begin{aligned} \text{ii } I_1 &= \int_0^{\frac{\pi}{2}} x \cos x \, dx = [x \sin x]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \sin x \, dx && \text{Using integration by parts} \\ &= \frac{\pi}{2} + [\cos x]_0^{\frac{\pi}{2}} \\ &= \frac{\pi}{2} + [0 - 1] = \frac{\pi}{2} - 1 \end{aligned}$$

b Integrating by parts with $u = x^n$ and $\frac{dv}{dx} = \cos x$

$$\frac{du}{dx} = nx^{n-1}, \quad v = \sin x$$

$$\begin{aligned} \text{So } I_n &= \int_0^{\frac{\pi}{2}} x^n \cos x \, dx = [x^n \sin x]_0^{\frac{\pi}{2}} - n \int_0^{\frac{\pi}{2}} x^{n-1} \sin x \, dx \\ &= \left(\frac{\pi}{2}\right)^n - n \int_0^{\frac{\pi}{2}} x^{n-1} \sin x \, dx \quad * \end{aligned}$$

Integrating by parts on $\int_0^{\frac{\pi}{2}} x^{n-1} \sin x \, dx$ with $u = x^{n-1}$ and $\frac{dv}{dx} = \sin x$

$$\frac{du}{dx} = (n-1)x^{n-2}, \quad v = -\cos x$$

$$\begin{aligned} \text{gives } \int_0^{\frac{\pi}{2}} x^{n-1} \sin x \, dx &= [-x^{n-1} \cos x]_0^{\frac{\pi}{2}} + (n-1) \int_0^{\frac{\pi}{2}} x^{n-2} \cos x \, dx \\ &= (n-1)I_{n-2} \quad \text{as } [-x^{n-1} \cos x]_0^{\frac{\pi}{2}} = 0 \end{aligned}$$

Substituting in *

$$I_n = \left(\frac{\pi}{2}\right)^n - n(n-1)I_{n-2}$$

$$\begin{aligned} \text{c } \int_0^{\frac{\pi}{2}} x^3 \cos x \, dx &= I_3 = \left(\frac{\pi}{2}\right)^3 - 3(2)I_1 \\ &= \left(\frac{\pi}{2}\right)^3 - 6\left(\frac{\pi}{2} - 1\right) && \text{Using a ii} \\ &= \frac{\pi^3}{8} - 3\pi + 6 \\ &= \frac{1}{8}(\pi^3 - 24\pi + 48) \end{aligned}$$

$$\begin{aligned} \text{d } \int_0^{\frac{\pi}{2}} x^4 \cos x \, dx &= I_4 = \left(\frac{\pi}{2}\right)^4 - 4(3)I_2 \\ &= \left(\frac{\pi}{2}\right)^4 - 12 \left\{ \left(\frac{\pi}{2}\right)^2 - 2(1)I_0 \right\} \\ &= \frac{\pi^4}{16} - 3\pi^2 + 24 && \text{as } I_0 = 1 \text{ from a i} \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 9

Question:

a Find $\int \frac{dx}{\sqrt{x^2 - 2x + 10}}$.

b Find $\int \frac{dx}{x^2 - 2x + 10}$.

c By using the substitution $x = \sin \theta$, show that $\int_0^{\frac{1}{2}} \frac{x^4}{\sqrt{(1-x^2)}} = \frac{(4\pi - 7\sqrt{3})}{64}$ [E]

Solution:

a $x^2 - 2x + 10 = (x-1)^2 + 9$

$$\text{So } \int \frac{dx}{\sqrt{x^2 - 2x + 10}} = \int \frac{dx}{\sqrt{(x-1)^2 + 9}}$$

Let $x-1 = 3\sinh u$, then $dx = 3\cosh u du$

$$\begin{aligned} \text{so } \int \frac{dx}{\sqrt{x^2 - 2x + 10}} &= \int \frac{3\cosh u}{3\cosh u} du \\ &= u + C \\ &= \operatorname{arsinh}\left(\frac{x-1}{3}\right) + C \end{aligned}$$

b $\int \frac{dx}{x^2 - 2x + 10} = \int \frac{dx}{(x-1)^2 + 9}$

Let $x-1 = 3\tan \theta$, then $dx = 3\sec^2 \theta d\theta$

$$\begin{aligned} \text{so } \int \frac{dx}{x^2 - 2x + 10} &= \int \frac{3\sec^2 \theta}{9\tan^2 \theta + 9} d\theta \\ &= \int \frac{3\sec^2 \theta}{9\sec^2 \theta} d\theta \\ &= \frac{1}{3}\theta + C \\ &= \frac{1}{3}\arctan\left(\frac{x-1}{3}\right) + C \end{aligned}$$

c Using the substitution $x = \sin \theta$, so $dx = \cos \theta d\theta$

$$\begin{aligned} \int_0^{\frac{1}{2}} \frac{x^4 dx}{\sqrt{1-x^2}} &= \int_0^{\frac{\pi}{6}} \frac{\sin^4 \theta \cos \theta d\theta}{\cos \theta} \\ &= \int_0^{\frac{\pi}{6}} \sin^4 \theta d\theta \\ &= \frac{1}{4} \int_0^{\frac{\pi}{6}} (1 - 2\cos 2\theta + \cos^2 2\theta) d\theta \\ &= \frac{1}{4} \int_0^{\frac{\pi}{6}} \left(1 - 2\cos 2\theta + \frac{1 + \cos 4\theta}{2}\right) d\theta \\ &= \frac{1}{4} \left[\frac{3\theta}{2} - \sin 2\theta + \frac{\sin 4\theta}{8} \right]_0^{\frac{\pi}{6}} \\ &= \frac{1}{4} \left(\frac{\pi}{4} - \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{16} \right) \\ &= \frac{(4\pi - 7\sqrt{3})}{64} \end{aligned}$$

$$\sin^4 \theta = (\sin^2 \theta)^2 = \frac{1}{4}(1 - \cos 2\theta)^2$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 10

Question:

Given that $I_n = \int_0^1 x^n (1-x)^{\frac{1}{3}} dx, n \geq 0,$

a show that $I_n = \frac{3n}{3n+4} I_{n-1}, n \geq 1$

b Hence find the exact value of $\int_0^1 (x+1)(1-x)^{\frac{1}{3}} dx.$

[E]

Solution:

a Using integration by parts on I_n , with $u = x^n$ and $\frac{dv}{dx} = (1-x)^{\frac{1}{3}}$

so $\frac{du}{dx} = nx^{n-1}$ and $v = -\frac{3}{4}(1-x)^{\frac{4}{3}}$

$$I_n = -\frac{3}{4} \left[x^n (1-x)^{\frac{4}{3}} \right]_0^1 + \frac{3n}{4} \int_0^1 x^{n-1} (1-x)^{\frac{4}{3}} dx$$

$$= \frac{3n}{4} \int_0^1 x^{n-1} (1-x)^{\frac{4}{3}} dx$$

$$= \frac{3n}{4} \int_0^1 x^{n-1} (1-x)(1-x)^{\frac{1}{3}} dx$$

$$= 6n \int_0^1 x^{n-1} (1-x)^{\frac{1}{3}} dx - \frac{3n}{4} \int_0^1 x^n (1-x)^{\frac{1}{3}} dx$$

$$\Rightarrow 4I_n = 6nI_{n-1} - \frac{3n}{4} I_n \Rightarrow I_n = \frac{24n}{3n+4} I_{n-1}$$

b $\int_0^1 (1+x)(1-x)^{\frac{4}{3}} dx = \int_0^1 (1+x^2)(1-x)^{\frac{1}{3}} dx = I_0 - I_2$

$$I_0 = \int_0^1 (1+x)^{\frac{1}{3}} dx = \left[-\frac{3}{4}(1-x)^{\frac{4}{3}} \right]_0^1 = \frac{3}{4}$$

Using a $I_2 = \frac{3}{5} I_1 = \frac{3}{5} \left(\frac{3}{7} I_0 \right) = \left(\frac{27}{140} \right)$

$$\text{So } \int_0^1 (1+x)(1-x)^{\frac{4}{3}} dx = \frac{3}{4} - \frac{27}{140} = \frac{78}{140} = \frac{39}{70}$$


Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 11

Question:



The curve C has parametric equations

$$x = t - \ln t,$$

$$y = 4\sqrt{t}, 1 \leq t \leq 4.$$

a Show that the length of C is $3 + \ln 4$.

The curve is rotated through 2π radians about the x -axis.

b Find the exact area of the curved surface generated.

[E]

Solution:

$$x = t - \ln t, \text{ so } \frac{dx}{dt} = 1 - \frac{1}{t}$$

$$y = 4\sqrt{t}, \text{ so } \frac{dy}{dt} = \frac{2}{\sqrt{t}}$$

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 1 - \frac{2}{t} + \frac{1}{t^2} + \frac{4}{t} = 1 + \frac{2}{t} + \frac{1}{t^2} = \left(1 + \frac{1}{t}\right)^2$$

$$\text{a Arc length} = \int_1^4 \sqrt{\left(1 + \frac{1}{t}\right)^2} dt = \int_1^4 \left(1 + \frac{1}{t}\right) dt = [t + \ln t]_1^4 = (4 + \ln 4) - 1 = 3 + \ln 4$$

$$\text{b Using } \int_{t_1}^{t_2} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt,$$

$$\text{the area of the surface is } 2\pi \int_1^4 4\sqrt{t} \left(1 + \frac{1}{t}\right) dt$$

$$= 8\pi \int_1^4 \left(\sqrt{t} + \frac{1}{\sqrt{t}}\right) dt$$

$$= 8\pi \left[\frac{2}{3} t^{\frac{3}{2}} + 2t^{\frac{1}{2}} \right]_1^4$$

$$= 8\pi \left[\left(\frac{16}{3} + 4 \right) - \left(\frac{2}{3} + 2 \right) \right]$$

$$= \frac{160\pi}{3}$$


Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 12

Question:



Above is a sketch of part of the curve with equation

$$y = x^2 \operatorname{arsinh} x.$$

The region R , shown shaded, is bounded by the curve, the x -axis and the line $x = 3$.

Show that the area of R is

$$9 \ln(3 + \sqrt{10}) - \frac{1}{9}(2 + 7\sqrt{10}). \quad \text{[E]}$$

Solution:

$$\text{Area} = \int_0^3 y \, dx = \int_0^3 x^2 \operatorname{arsinh} x \, dx$$

Using integration by parts on I_x , with $u = \operatorname{arsinh} x$ and $\frac{dv}{dx} = x^2$

$$\text{so } \frac{du}{dx} = \frac{1}{\sqrt{1+x^2}} \text{ and } v = \frac{x^3}{3}$$

$$\int x^2 \operatorname{arsinh} x \, dx = \frac{1}{3} x^3 \operatorname{arsinh} x - \frac{1}{3} \int \frac{x^3}{\sqrt{1+x^2}} \, dx$$

Let $x = \sinh u$ so $dx = \cosh u \, du$

$$\begin{aligned} \int_0^3 x^2 \operatorname{arsinh} x \, dx &= 9 \operatorname{arsinh} 3 - \frac{1}{3} \int_0^{\operatorname{arsinh} 3} \frac{\sinh^3 u}{\cosh u} \cosh u \, du \\ &= 9 \operatorname{arsinh} 3 - \frac{1}{3} \int_0^{\operatorname{arsinh} 3} \sinh^3 u \, du \\ &= 9 \operatorname{arsinh} 3 - \frac{1}{3} \int_0^{\operatorname{arsinh} 3} \sinh u (\cosh^2 u - 1) \, du \\ &= 9 \operatorname{arsinh} 3 - \frac{1}{3} \left[\frac{1}{3} \cosh^3 u - \cosh u \right]_0^{\operatorname{arsinh} 3} \end{aligned}$$

When $x = 3$, $\sinh u = 3$ so $\cosh u = \sqrt{1 + \sinh^2 u} = \sqrt{10}$

$$\begin{aligned} \text{So } \int_0^3 x^2 \operatorname{arsinh} x \, dx &= 9 \ln \{3 + \sqrt{10}\} - \frac{1}{3} \left[\frac{10}{3} \sqrt{10} - \sqrt{10} - \left(\frac{1}{3} - 1 \right) \right] \\ &= 9 \ln \{3 + \sqrt{10}\} - \frac{1}{9} [7\sqrt{10} + 2] \end{aligned}$$

You could use integration by parts with

$$u = x^2 \text{ and } \frac{dv}{dx} = \frac{x}{\sqrt{1+x^2}}$$

$$\operatorname{arsinh} x = \ln \{x + \sqrt{1+x^2}\}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 13

Question:

a Use the substitution $u = x^2$ to find $\int_0^1 \frac{x}{1+x^4} dx$

b Find

i $\int \frac{1}{\sqrt{4x-x^2}} dx$

ii $\int \frac{4-2x}{\sqrt{4x-x^2}} dx$.

Hence, or otherwise, evaluate

iii $\int_3^4 \frac{5-2x}{\sqrt{4x-x^2}} dx$.

Solution:

a Using $x^2 = u$ '2x dx' becomes 'du'

$$\begin{aligned} \text{So } \int_0^1 \frac{x}{1+x^4} dx &= \frac{1}{2} \int_0^1 \frac{du}{1+u^2} \\ &= \frac{1}{2} [\arctan u]_0^1 \\ &= \frac{\pi}{8} \end{aligned}$$

b i $4x - x^2 = -(x^2 - 4x) = -[(x-2)^2 - 4]$
 $= 4 - (x-2)^2$

$$\begin{aligned} \int \frac{1}{\sqrt{4x-x^2}} dx &= \int \frac{1}{\sqrt{4-(x-2)^2}} dx \\ &= \arcsin\left(\frac{x-2}{2}\right) + C \end{aligned}$$

Using $\int \frac{1}{\sqrt{a^2-x^2}} = \arcsin\left(\frac{x}{a}\right) + C$

ii $\int \frac{4-2x}{\sqrt{4x-x^2}} dx$
 $= 2(4x-x^2)^{\frac{1}{2}} + C$

Notice that $\frac{d}{dx}(4x-x^2) = 4-2x$

iii $\int_3^4 \frac{5-2x}{\sqrt{4x-x^2}} dx = \int_3^4 \left\{ \frac{1}{\sqrt{4x-x^2}} + \frac{4-2x}{\sqrt{4x-x^2}} \right\} dx$
 $= \int_3^4 \frac{1}{\sqrt{4x-x^2}} dx + \int_3^4 \frac{4-2x}{\sqrt{4x-x^2}} dx$
 $= \left[\arcsin\left(\frac{x-2}{2}\right) + 2(4x-x^2)^{\frac{1}{2}} \right]_3^4$
 $= \left(\frac{\pi}{2}\right) - \left(\frac{\pi}{6} + 2\sqrt{3}\right) = \frac{\pi}{3} - 2\sqrt{3}$

Using i and ii

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics


Integration

Exercise I, Question 14

Question:

The curve C shown in the diagram has equation $y^2 = 4x, 0 \leq x \leq 1$.

The part of the curve in the first quadrant is rotated through 2π radians about the x -axis.



- a Show that the surface area of the solid generated is given by $4\pi \int_0^1 \sqrt{1+x} dx$.
- b Find the exact value of this surface area.
- c Show also that the length of the curve C , between the points $(1, -2)$ and $(1, 2)$, is given by $2 \int_0^1 \sqrt{\frac{x+1}{x}} dx$.
- d Use the substitution $x = \sinh^2 \theta$ to show that the exact value of this length is $2[\sqrt{2} + \ln(1 + \sqrt{2})]$. **[E]**

Solution:

$y = 2\sqrt{x}$ represents the section of curve for $x \geq 0, y \geq 0$, so $\frac{dy}{dx} = \frac{1}{\sqrt{x}}$

a Using $2\pi \int_{x_1}^{x_2} y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$

$$\begin{aligned} \text{area of surface} &= 2\pi \int_0^1 2\sqrt{x} \sqrt{1 + \frac{1}{x}} dx \\ &= 4\pi \int_0^1 \sqrt{x} \sqrt{\frac{x+1}{x}} dx \\ &= 4\pi \int_0^1 \sqrt{1+x} dx \end{aligned}$$

b $4\pi \int_0^1 \sqrt{1+x} dx = 4\pi \left[\frac{2}{3}(1+x)^{\frac{3}{2}} \right]_0^1$
 $= \frac{8\pi}{3}(2\sqrt{2}-1)$

c Using the symmetry of the parabola, arc length is $2 \times$ the length of arc from origin to $(1, 2)$

$$\begin{aligned} \text{so arc length} &= 2 \int_0^1 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\ &= 2 \int_0^1 \sqrt{\frac{x+1}{x}} dx \end{aligned}$$

d Using $x = \sinh^2 \theta, dx = 2 \sinh \theta \cosh \theta d\theta$

$$\begin{aligned} 2 \int \sqrt{\frac{x+1}{x}} dx &= 2 \int \sqrt{\frac{\sinh^2 \theta + 1}{\sinh^2 \theta}} 2 \sinh \theta \cosh \theta d\theta \\ &= 4 \int \cosh^2 \theta d\theta \\ &= 2 \int (1 + \cosh 2\theta) d\theta \\ &= 2 \left(\theta + \frac{\sinh 2\theta}{2} \right) + C \\ &= 2(\theta + \sinh \theta \cosh \theta) + C \\ &= 2 \{ \operatorname{arsinh} \sqrt{x} + \sqrt{x} \sqrt{1+x} \} + C \end{aligned}$$

$$\begin{aligned} \text{So arc length} &= 2 \int_0^1 \sqrt{\frac{x+1}{x}} dx = 2(\operatorname{arsinh} 1 + \sqrt{2}) \\ &= 2[\sqrt{2} + \ln(1 + \sqrt{2})] \end{aligned}$$

$\operatorname{arsinh} x = \ln \{ x + \sqrt{1+x^2} \}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 15

Question:

a Show that $\int x \operatorname{arcosh} x \, dx = \frac{1}{4}(2x^2 - 1) \operatorname{arcosh} x - \frac{1}{4}x\sqrt{x^2 - 1} + C$

b Hence, using the substitution $x = u^2$, find $\int \operatorname{arcosh}(\sqrt{x}) \, dx$.

Solution:

a Using integration by parts with $u = \operatorname{arcosh} x$ and $\frac{dv}{dx} = x$,

$$\frac{du}{dx} = \frac{1}{\sqrt{x^2 - 1}} \quad \text{and} \quad v = \frac{x^2}{2}$$

$$\text{So } \int x \operatorname{arcosh} x \, dx = \frac{x^2}{2} \operatorname{arcosh} x - \int \frac{x^2}{2\sqrt{x^2 - 1}} \, dx \quad *$$

Substitute $x = \cosh u$ in $\int \frac{x^2}{\sqrt{x^2 - 1}} \, dx$ gives

$$\begin{aligned} \int \frac{x^2}{\sqrt{x^2 - 1}} \, dx &= \int \frac{\cosh^2 u}{\sinh u} \sinh u \, du \\ &= \int \cosh^2 u \, du \\ &= \frac{1}{2} \int (1 + \cosh 2u) \, du \\ &= \frac{1}{2} [u + \sinh u \cosh u] + C \\ &= \frac{1}{2} [\operatorname{arcosh} x + x\sqrt{x^2 - 1}] + C \end{aligned}$$

You could use integration by parts with $u = x$ and

$$\frac{dv}{dx} = \frac{x}{\sqrt{x^2 - 1}}$$

$$\begin{aligned} \text{So } \int x \operatorname{arcosh} x \, dx &= \frac{x^2}{2} \operatorname{arcosh} x - \frac{1}{4} [\operatorname{arcosh} x + x\sqrt{x^2 - 1}] + C \quad \text{from } * \\ &= \frac{1}{4} (2x^2 - 1) \operatorname{arcosh} x - \frac{1}{4} x\sqrt{x^2 - 1} + C \end{aligned}$$

b Let $x = u^2$, so $dx = 2u \, du$,

$$\begin{aligned} \text{then } \int \operatorname{arcosh}(\sqrt{x}) \, dx &= 2 \int u \operatorname{arcosh} u \, du \\ &= \frac{1}{2} (2u^2 - 1) \operatorname{arcosh} u - \frac{1}{2} u\sqrt{u^2 - 1} + C \\ &= \frac{1}{2} (2x - 1) \operatorname{arcosh} \sqrt{x} - \frac{1}{2} \sqrt{x}\sqrt{x - 1} + C \end{aligned}$$

Using a

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 16

Question:

Given that $I_n = \int \frac{\sin(2n+1)x}{\sin x} dx$,

a show that $I_n - I_{n-1} = \frac{\sin 2nx}{n}$.

b Hence find I_5 .

c Show that, for all positive integers n , $\int_0^{\frac{\pi}{2}} \frac{\sin(2n+1)x}{\sin x} dx$ always has the same value, which should be found.

Solution:

$$\begin{aligned} \text{a } I_n - I_{n-1} &= \int \frac{[\sin(2n+1)x - \sin(2n-1)x]}{\sin x} dx \\ &= \int \frac{2 \cos 2nx \sin x}{\sin x} dx \\ &= \int 2 \cos 2nx dx \\ &= \frac{\sin 2nx}{n} \end{aligned}$$

$$\sin A - \sin B = 2 \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right)$$

$$\text{b } I_5 - I_4 = \frac{\sin 10x}{5}, I_4 - I_3 = \frac{\sin 8x}{4}, I_3 - I_2 = \frac{\sin 6x}{3}, I_2 - I_1 = \frac{\sin 4x}{2}$$

$$I_1 - I_0 = \sin 2x$$

$$\text{Adding: } I_5 = \frac{\sin 10x}{5} + \frac{\sin 8x}{4} + \frac{\sin 6x}{3} + \frac{\sin 4x}{2} + \sin 2x + I_0$$

$$\begin{aligned} \text{where } I_0 &= \int 1 dx = x + C \\ &= \frac{\sin 10x}{5} + \frac{\sin 8x}{4} + \frac{\sin 6x}{3} + \frac{\sin 4x}{2} + \sin 2x + x + C \end{aligned}$$

$$\text{c } \int_0^{\frac{\pi}{2}} \frac{\sin(2n+1)x}{\sin x} dx - \int_0^{\frac{\pi}{2}} \frac{\sin(2n-1)x}{\sin x} dx = \left[\frac{\sin 2nx}{n} \right]_0^{\frac{\pi}{2}} = \frac{\sin(n\pi)}{n}$$

$$\text{So, if } n \text{ is any a positive integer } \int_0^{\frac{\pi}{2}} \frac{\sin(2n+1)x}{\sin x} dx - \int_0^{\frac{\pi}{2}} \frac{\sin(2n-1)x}{\sin x} dx = 0$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \frac{\sin(2n+1)x}{\sin x} dx = \int_0^{\frac{\pi}{2}} \frac{\sin(2n-1)x}{\sin x} dx = \dots = \int_0^{\frac{\pi}{2}} \frac{\sin x}{\sin x} dx = \frac{\pi}{2}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 17


Question:

The diagram shows part of the graph of the curve with equation $y^2 = \frac{1}{3}x(x-1)^2$.

- a Show that the length of the loop is $\frac{4\sqrt{3}}{3}$.

The arc OA (in blue) is rotated completely about the x -axis.

- b Find the area of the surface generated.



Solution:

- a The point A on the curve has coordinates $(1, 0)$.

Using symmetry, the length of the loop is $2 \int_0^1 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$.

$$\text{As } y^2 = \frac{1}{3}x(x-1)^2 = \frac{1}{3}(x^3 - 2x^2 + x)$$

$$2y \frac{dy}{dx} = \frac{1}{3}(3x^2 - 4x + 1) = \frac{1}{3}(3x-1)(x-1)$$

$$\text{So } \frac{dy}{dx} = \frac{\frac{1}{3}(3x-1)(x-1)}{\pm 2\sqrt{\frac{x}{3}}(x-1)} = \pm \frac{1}{2\sqrt{3}} \frac{(3x-1)}{\sqrt{x}}$$

$$\text{and } 1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{9x^2 - 6x + 1}{12x} = \frac{9x^2 + 6x + 1}{12x} = \frac{(3x+1)^2}{12x}$$

$$\begin{aligned} \text{Therefore, arc length} &= 2 \int_0^1 \frac{3x+1}{2\sqrt{3}\sqrt{x}} dx \\ &= \frac{1}{\sqrt{3}} \int_0^1 \left(3\sqrt{x} + \frac{1}{\sqrt{x}}\right) dx \\ &= \frac{1}{\sqrt{3}} \left[2x^{\frac{3}{2}} + 2\sqrt{x}\right]_0^1 \\ &= \frac{4}{\sqrt{3}} = \frac{4\sqrt{3}}{3} \end{aligned}$$

- b Using $2\pi \int_x^{x_2} y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$ for area of surface generated about the x -axis

$$\begin{aligned} \text{Area of surface} &= 2\pi \int_0^1 \frac{1}{\sqrt{3}} \sqrt{x}(1-x) \frac{(3x+1)}{\sqrt{12x}} dx \\ &= \frac{\pi}{3} \int_0^1 (1-x)(3x+1) dx \\ &= \frac{\pi}{3} \int_0^1 (1+2x-3x^2) dx \\ &= \frac{\pi}{3} \left[x + x^2 - x^3\right]_0^1 \\ &= \frac{\pi}{3} \end{aligned}$$

Note: y is +ve for OA , so you need to take $y = -\frac{\sqrt{x}(x-1)}{\sqrt{3}} = \frac{\sqrt{x}(1-x)}{\sqrt{3}}$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 18

Question:

- a Find $\int \frac{1}{\sinh x + 2 \cosh x} dx$.
- b Show that $\int_1^4 \frac{3x-1}{\sqrt{x^2-2x+10}} dx = 9(\sqrt{2}-1) + 2\operatorname{arsinh} 1$. [E]

Solution:

- a Using the exponential forms

$$\begin{aligned} \int \frac{1}{\sinh x + 2 \cosh x} dx &= \int \frac{1}{\left(\frac{e^x - e^{-x}}{2}\right) + 2\left(\frac{e^x + e^{-x}}{2}\right)} dx \\ &= \int \frac{2}{3e^x + e^{-x}} dx \\ &= \int \frac{2e^x}{3e^{2x} + 1} dx \end{aligned}$$

Using the substitution $u = e^x$, then $\frac{du}{dx} = e^x$ so ' $e^x dx$ ' can be replaced by ' du '.

$$\begin{aligned} \text{So } \int \frac{1}{\sinh x + 2 \cosh x} dx &= \int \frac{2}{3u^2 + 1} du \\ &= \frac{2}{3} \int \frac{1}{u^2 + \frac{1}{3}} du \\ &= \frac{2}{3} (\sqrt{3}) \arctan(\sqrt{3}u) + C \\ &= \frac{2}{\sqrt{3}} \arctan(\sqrt{3}e^x) + C \end{aligned}$$

- b $x^2 - 2x + 10 = (x-1)^2 + 9$

So let $x-1 = 3 \sinh u$, then $dx = 3 \cosh u du$

$$\begin{aligned} \text{and } \int \frac{3x-1}{\sqrt{x^2-2x+10}} dx &= \int \frac{9 \sinh u + 2}{\sqrt{9 \sinh^2 u + 9}} 3 \cosh u du \\ &= \int \frac{9 \sinh u + 2}{3 \cosh u} 3 \cosh u du \\ &= 9 \cosh u + 2u + C \\ &= 9 \sqrt{1 + \left(\frac{x-1}{3}\right)^2} + 2 \operatorname{arsinh}\left(\frac{x-1}{3}\right) + C \end{aligned}$$

$$\begin{aligned} \text{So } \int_1^4 \frac{3x-1}{\sqrt{x^2-2x+10}} dx &= [9\sqrt{2} + 2\operatorname{arsinh} 1] - [9] \\ &= 9(\sqrt{2}-1) + 2\operatorname{arsinh} 1 \end{aligned}$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 19

Question:

Given that $I_n = \int \sec^n x \, dx$;

a by writing $\sec^n x = \sec^{n-2} x \sec^2 x$, show that, for $n \geq 2$,

$$(n-1)I_n = \sec^{n-2} x \tan x + (n-2)I_{n-2}.$$

b Find I_5 .

c Hence show that $\int_0^{\frac{\pi}{4}} \sec^5 x \, dx = \frac{1}{8}(7\sqrt{2} + 3\ln(1+\sqrt{2}))$

Solution:

a $\int \sec^n x \, dx = \int \sec^{n-2} x \sec^2 x \, dx$

Let $u = \sec^{n-2} x$ and $\frac{dv}{dx} = \sec^2 x$

$$\frac{du}{dx} = (n-2)\sec^{n-3} x (\sec x \tan x) = (n-2)\sec^{n-2} x \tan x \text{ and } v = \tan x$$

Integrating by parts

$$\begin{aligned} \int \sec^n x \, dx &= I_n = \sec^{n-2} x \tan x - (n-2) \int \sec^{n-2} x \tan^2 x \, dx \\ &= \sec^{n-2} x \tan x - (n-2) \int \sec^{n-2} x (\sec^2 x - 1) \, dx \\ &= \sec^{n-2} x \tan x - (n-2) \int \sec^n x \, dx + (n-2) \int \sec^{n-2} x \, dx \\ I_n &= \sec^{n-2} x \tan x - (n-2)I_n + (n-2)I_{n-2} \end{aligned}$$

So $(n-1)I_n = \sec^{n-2} x \tan x + (n-2)I_{n-2}$, $n \geq 2$, *

b $\int \sec^5 x \, dx = I_5 = \frac{1}{4} \sec^3 x \tan x + \frac{3}{4} I_3$ ← Substituting $n=5$ in *

$$= \frac{1}{4} \sec^3 x \tan x + \frac{3}{4} \left(\frac{1}{2} \sec x \tan x + \frac{1}{2} I_1 \right)$$
 ← Substituting $n=3$ in *

But $I_1 = \int \sec x \, dx = \ln |\sec x + \tan x| + C$ ← On Edexcel formula sheet

So $\int \sec^5 x \, dx = I_5 = \frac{1}{4} \sec^3 x \tan x + \frac{3}{8} \sec x \tan x + \frac{3}{8} \ln |\sec x + \tan x| + C$

c $\int_0^{\frac{\pi}{4}} \sec^5 x \, dx = \frac{1}{4} (\sqrt{2})^3 + \frac{3}{8} (\sqrt{2}) + \frac{3}{8} \ln (\sqrt{2} + 1)$
 $= \frac{1}{8} [7\sqrt{2} + 3\ln(\sqrt{2} + 1)]$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 20

Question:

- a Show by using a suitable substitution for x , that

$$\int \sqrt{a^2 - x^2} \, dx = \frac{a^2}{2} \arcsin\left(\frac{x}{a}\right) + \frac{x}{2} \sqrt{a^2 - x^2} + C$$

- b Hence show that the area of the region enclosed by the ellipse with equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ is } \pi ab.$$

Solution:

- a Let $x = a \sin \theta$, then $\frac{dx}{d\theta} = a \cos \theta$

$$\begin{aligned} \text{So } \int \sqrt{a^2 - x^2} \, dx &= \int a^2 \cos^2 \theta \, d\theta \\ &= \frac{a^2}{2} \int (1 + \cos 2\theta) \, d\theta \\ &= \frac{a^2}{2} \left(\theta + \frac{\sin 2\theta}{2} \right) + C \\ &= \frac{a^2}{2} (\theta + \sin \theta \cos \theta) + C \\ &= \frac{a^2}{2} \left(\arcsin\left(\frac{x}{a}\right) + \frac{x}{a} \sqrt{1 - \left(\frac{x}{a}\right)^2} \right) + C \\ &= \frac{a^2}{2} \arcsin\left(\frac{x}{a}\right) + \frac{x}{2} \sqrt{a^2 - x^2} + C \end{aligned}$$

- b Area enclosed by the ellipse = $4 \times$ area enclosed by arc in first quadrant and the positive coordinate axes (symmetry)

$$= 4 \int_0^a y \, dx$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \Rightarrow y = \pm \frac{b}{a} \sqrt{a^2 - x^2}$$

+ve square root required

$$\text{So area} = 4 \frac{b}{a} \left[\frac{a^2}{2} \arcsin\left(\frac{x}{a}\right) + \frac{x}{2} \sqrt{a^2 - x^2} \right]_0^a$$

from a

$$= 2ab \arcsin 1$$

$$= \pi ab$$

Solutionbank FP3

Edexcel AS and A Level Modular Mathematics

Integration

Exercise I, Question 21

Question:

- a Show by using a suitable substitution for x , that

$$\int \sqrt{a^2 - x^2} \, dx = \frac{a^2}{2} \arcsin\left(\frac{x}{a}\right) + \frac{x}{2} \sqrt{a^2 - x^2} + C$$

- b Hence show that the area of the region enclosed by the ellipse with equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ is } \pi ab.$$

Solution:

- a Let $x = a \sin \theta$, then $\frac{dx}{d\theta} = a \cos \theta$

$$\begin{aligned} \text{So } \int \sqrt{a^2 - x^2} \, dx &= \int a^2 \cos^2 \theta \, d\theta \\ &= \frac{a^2}{2} \int (1 + \cos 2\theta) \, d\theta \\ &= \frac{a^2}{2} \left(\theta + \frac{\sin 2\theta}{2} \right) + C \\ &= \frac{a^2}{2} (\theta + \sin \theta \cos \theta) + C \\ &= \frac{a^2}{2} \left(\arcsin\left(\frac{x}{a}\right) + \frac{x}{a} \sqrt{1 - \left(\frac{x}{a}\right)^2} \right) + C \\ &= \frac{a^2}{2} \arcsin\left(\frac{x}{a}\right) + \frac{x}{2} \sqrt{a^2 - x^2} + C \end{aligned}$$

- b Area enclosed by the ellipse = $4 \times$ area enclosed by arc in first quadrant (symmetry)

$$= 4 \int_0^a y \, dx$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \Rightarrow y = \pm \frac{b}{a} \sqrt{a^2 - x^2}$$

{ +ve square root required }

$$\text{So area} = 4 \frac{b}{a} \left[\frac{a^2}{2} \arcsin\left(\frac{x}{a}\right) + \frac{x}{2} \sqrt{a^2 - x^2} \right]_0^a$$

from a

$$= 2ab \arcsin 1$$

$$= \pi ab$$