



## MECHANICS (C) UNIT 2 TEST PAPER 1

Take  $g = 9.8 \text{ ms}^{-2}$  and give all answers correct to 3 significant figures where necessary.

- A car of mass  $1200 \text{ kg}$  decelerates from  $30 \text{ ms}^{-1}$  to  $20 \text{ ms}^{-1}$  in 6 seconds at a constant rate.
  - Find the magnitude, in  $\text{N}$ , of the decelerating force. [2]
  - Find the loss, in  $\text{J}$ , in the car's kinetic energy. [2]
- Eddie, whose mass is  $71 \text{ kg}$ , rides a bicycle of mass  $25 \text{ kg}$  up a hill inclined at an angle  $\alpha$  to the horizontal, where  $\sin \alpha = \frac{1}{12}$ . When Eddie is working at a rate of  $600 \text{ W}$ , he is moving at a constant speed of  $6 \text{ ms}^{-1}$ .  
Find the magnitude of the non-gravitational resistance to his motion. [6]

- A bird of mass  $0.5 \text{ kg}$ , flying around a vertical feeding post at a constant speed of  $4 \text{ ms}^{-1}$ , inclines its wings so as to move in a horizontal circle of radius  $2 \text{ m}$ . The lifting force  $L$  newtons acts perpendicular to the bird's wings, as shown. Modelling the bird as a particle, find, to the nearest degree, the angle  $\theta$  that its wings make with the vertical. [7]



The diagram shows a body which may be modelled as a uniform lamina.


The body is suspended from the point marked  $A$  and rests in equilibrium.

- Calculate, to the nearest degree, the angle which the edge  $AB$  then makes with the vertical. [7]

Frank suggests that the angle between  $AB$  and the vertical would be smaller if the lamina were made from lighter material.

- State, with a brief explanation, whether Frank is correct. [2]

- A uniform rod  $AB$ , of mass  $0.8 \text{ kg}$  and length  $10a$ , is supported at the end  $A$  by a light inextensible vertical string and rests in limiting equilibrium on a rough fixed peg at  $C$ , where  $AC = 7a$ .



- Draw a diagram to show all the forces acting on the rod. [2]
- Find the magnitude of the tension in the string. [3]

Given further that  $AB$  makes an angle of  $20^\circ$  with the horizontal,

- find the magnitude of the normal reaction exerted by the peg on the rod at  $C$ . [4]

- Two particles  $A$  and  $B$ , of mass  $m$  and  $km$  respectively, are moving in the same direction on a smooth horizontal surface.  $A$  has speed  $4u$  and  $B$  has speed  $u$ . The coefficient of restitution between  $A$  and  $B$  is  $e$ .  $A$  collides directly with  $B$ , and in the collision the direction of  $A$ 's motion is reversed. Immediately after the impact,  $B$  has speed  $2u$ .

- Show that the speed of  $A$  immediately after the impact is  $u(3e - 2)$ . [3]
- Deduce the range of possible values of  $e$ . [3]

(iii) Show that  $4 < k \leq 5$ . [6]

7. A ball is projected from ground level with speed  $34 \text{ ms}^{-1}$  at an angle  $\alpha$  above the horizontal, where  $\tan \alpha = \frac{8}{15}$ .

(i) Find the greatest height reached by the ball above ground level. [5]

While it is descending, the ball hits a horizontal ledge 6 metres above ground level.

(ii) Find the horizontal distance travelled by the ball before it hits the ledge. [5]

(iii) Find the speed of the ball at the instant when it hits the ledge. [3]

### MECHANICS 2 (C) TEST PAPER 1 : ANSWERS AND MARK SCHEME

- (i) Deceleration =  $\frac{5}{3} \text{ ms}^{-2}$  Force =  $1200 \times \frac{5}{3} = 2000 \text{ N}$  M1 A1  
 (ii) K.E. lost =  $600(30^2 - 20^2) = 300\,000 \text{ J}$  M1 A1 4
- $P = Fv : 600 = 6F$   $F = 100 \text{ N}$  M1 A1 A1  
 $100 = 96g \sin \alpha + R$   $R = 100 - 8g = 21.6 \text{ N}$  M1 A1 A1 6
- $L \sin \theta = 0.5g = 4.9$   $L \cos \theta = mv^2/r = 0.5 \times 16 \div 2 = 4$  M1 A1 M1 A1  
 $\tan \theta = 4.9 \div 4 = 1.225$   $\theta = 50.8^\circ$  M1 A1 A1 7
- (i)  $600(25, 6) + 600(30, 27) = 1200(\bar{x}, \bar{y})$   $\bar{x} = 27.5$ ,  $\bar{y} = 16.5$  M1 A1 M1 A1 A1  
 $\tan \theta = 16.5 \div 27.5 = 0.6$   $\theta = 31^\circ$  M1 A1  
 (ii) No : centre of mass depends only on area, not on density B1 B1 9
- (i) Diagram showing weight, tension, normal reaction, friction B2  
 (ii) M(C) :  $T(7a \cos \alpha) = 0.8g(2a \cos \alpha)$   $T = 2(0.8g) \div 7 = 2.24 \text{ N}$  M1 A1 A1  
 (iii) Resolve perp. to rod :  $R + T \cos \alpha = 0.8g \cos \alpha$  M1 A1  
 $R = 5.6 \cos 20^\circ = 5.26 \text{ N}$  M1 A1 9
- (i)  $(v_B - v_A)/(u - 4u) = -e$   $2u - v_A = 3eu$   $v_A = u(2 - 3e)$  M1 A1  
 $v_A < 0$ , so speed =  $u(3e - 2)$  A1  
 (ii) Since  $v_A < 0$ ,  $2 - 3e < 0$   $\frac{2}{3} < e \leq 1$  M1 A1 A1  
 (iii)  $4mu + kmv = mv_A + kmv_B$   $v_A + 2ku = 4u + kv$  M1 A1  
 $v_A = u(4 - k)$ , so  $4 - k = 2 - 3e$   $k = 3e + 2$  M1 A1  
 $\frac{2}{3} < e \leq 1$ , so  $4 < k \leq 5$  M1 A1 12
- (i)  $y = (u \sin \alpha)t - \frac{1}{2}gt^2 = 16t - 4.9t^2$  M1 A1  
 When  $y$  is max.,  $16 - 9.8t = 0$   $t = 1.63$   $y = 13.1 \text{ m}$  M1 A1 A1  
 (ii) When  $y = 6$ ,  $4.9t^2 - 16t + 6 = 0$  B1  
 $t = (16 + \sqrt{138.4})/9.8 = 2.83$   $x = (u \cos \alpha)t = 30t = 85.0 \text{ m}$  M1 A1 M1 A1  
 (iii)  $m(34^2) = mg(6) + \frac{1}{2}mv^2$   $v^2 = 1038$   $v = 32.2 \text{ ms}^{-1}$  M1 A1 A1 13