

1. A parcel of mass 2.5 kg is moving in a straight line on a smooth horizontal floor. Initially the parcel is moving with speed 8 m s^{-1} . The parcel is brought to rest in a distance of 20 m by a constant horizontal force of magnitude R newtons. Modelling the parcel as a particle, find

(a) the kinetic energy lost by the parcel in coming to rest,

(2)

(b) the value of R .

(3)



- $$\mathbf{p} = (3t^2 - 6t + 4)\mathbf{i} + (3t^3 - 4t)\mathbf{j}.$$

(a) the velocity of P at time t seconds,

(2)

- (3)

(c) find the velocity of P immediately after the impulse.

(4)

- (a) Show that $\sin \theta = \frac{1}{14}$.

(b) Find the value of y .

4.




Figure 1

A set square S is made by removing a circle of centre O and radius 3 cm from a triangular piece of wood. The piece of wood is modelled as a uniform triangular lamina ABC , with $\angle ABC = 90^\circ$, $AB = 12$ cm and $BC = 21$ cm. The point O is 5 cm from AB and 5 cm from BC , as shown in Figure 1.

- (a) Find the distance of the centre of mass of S from

- (i) AB ,

- (ii) BC .

(9)

The set square is freely suspended from C and hangs in equilibrium.

- (b) Find, to the nearest degree, the angle between CB and the vertical.

(3)





Leave
blank

Question 4 continued



The diagram shows a rod AB of total length $5a$ pivoted at end A. A force of magnitude $4a$ is applied perpendicular to the rod at a point located $4a$ from A. A weight of magnitude a acts vertically downwards at point C, which is a distance a from B. The rod AB makes an angle of 30° with the vertical.

A ladder AB , of mass m and length $4a$, has one end A resting on rough horizontal ground. The other end B rests against a smooth vertical wall. A load of mass $3m$ is fixed on the ladder at the point C , where $AC = a$. The ladder is modelled as a uniform rod in a vertical plane perpendicular to the wall and the load is modelled as a particle. The ladder rests in limiting equilibrium making an angle of 30° with the wall, as shown in Figure 2.

(10)



The diagram shows a particle's path from point A to point B. Point A is at a height of 47.5 m and a horizontal distance of 30 m from point O. The path is a curve starting at A and ending at B. The velocity vector at A is $(2u\mathbf{i} + 5u\mathbf{j}) \text{ m s}^{-1}$.

[In this question, the unit vectors \mathbf{i} and \mathbf{j} are in a vertical plane, \mathbf{i} being horizontal and \mathbf{j} being vertical.]

(a) Show that the time taken for P to move from A to B is 5 s. (6)

(b) Find the value of u . (2)

(c) Find the speed of P at B . (5)



- (b) Find the total kinetic energy lost in the collision. (5)

(c) Calculate the range of values of e for which there will be a second collision between P and Q .

Leave
blank**Question 7 continued**