

- Find the acceleration of the car at the instant when its speed is 15 m s^{-1} .

(5)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Figure 1 shows a ladder AB , of mass 25 kg and length 4 m, resting in equilibrium with one end A on rough horizontal ground and the other end B against a smooth vertical wall. The ladder is in a vertical plane perpendicular to the wall. The coefficient of friction between the ladder and the ground is $\frac{11}{25}$. The ladder makes an angle β with the ground. When Reece, who has mass 75 kg, stands at the point C on the ladder, where $AC = 2.8$ m, the ladder is on the point of slipping. The ladder is modelled as a uniform rod and Reece is modelled as a particle.

- (a) Find the magnitude of the frictional force of the ground on the ladder. (3)
- (b) Find, to the nearest degree, the value of β . (6)
- (c) State how you have used the modelling assumption that Reece is a particle. (1)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- (a) Calculate the work done against friction in moving the block from A to B .

The block passes through A with a speed of 2 m s^{-1} .

- (b) Find the speed of the block at B .


(4)

- $$v = \begin{cases} 10t - 2t^2, & 0 \leq t \leq 6, \\ \frac{-432}{t^2}, & t > 6. \end{cases}$$

$$(a) \ t = 6, \tag{3}$$
$$(b) \ t = 10. \tag{5}$$
This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Leave
blank**Question 4 continued****Q4****(Total 8 marks)**

5.

**Figure 2**

A uniform lamina $ABCD$ is made by joining a uniform triangular lamina ABD to a uniform semi-circular lamina DBC , of the same material, along the edge BD , as shown in Figure 2. Triangle ABD is right-angled at D and $AD = 18$ cm. The semi-circle has diameter BD and $BD = 12$ cm.

- (a) Show that, to 3 significant figures, the distance of the centre of mass of the lamina $ABCD$ from AD is 4.69 cm. (4)

Given that the centre of mass of a uniform semicircular lamina, radius r , is at a distance $\frac{4r}{3\pi}$ from the centre of the bounding diameter,

- (b) find, in cm to 3 significant figures, the distance of the centre of mass of the lamina $ABCD$ from BD . (4)


The lamina is freely suspended from B and hangs in equilibrium.

- (c) Find, to the nearest degree, the angle which BD makes with the vertical. (4)



[illegible]

6.

**Figure 3**

A cricket ball is hit from a point A with velocity of $(p\mathbf{i} + q\mathbf{j}) \text{ m s}^{-1}$, at an angle α above the horizontal. The unit vectors \mathbf{i} and \mathbf{j} are respectively horizontal and vertically upwards. The point A is 0.9 m vertically above the point O , which is on horizontal ground.

The ball takes 3 seconds to travel from A to B , where B is on the ground and $OB = 57.6 \text{ m}$, as shown in Figure 3. By modelling the motion of the cricket ball as that of a particle moving freely under gravity,

- find the value of p , (2)
- show that $q = 14.4$, (3)
- find the initial speed of the cricket ball, (2)
- find the exact value of $\tan \alpha$. (1)
- Find the length of time for which the cricket ball is at least 4 m above the ground. (6)
- State an additional physical factor which may be taken into account in a refinement of the above model to make it more realistic. (1)



- (a) Show that the speed of Q immediately after the collision is $\frac{1}{5}(9e + 4)u$. (5)

(b) Show that $e = \frac{1}{4}$. (4)

(c) Show that P is a distance $\frac{3}{5}d$ from the wall at the instant when Q hits the wall. (4)

(d) find, in terms of d , the distance of the point B from the wall.

