



General Certificate of Education

Mathematics 6360 Statistics 6380

MS/SS1B Statistics 1B

Mark Scheme

2006 examination – January series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Key To Mark Scheme And Abbreviations Used In Marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MS/SS1B

Q	Solution	Marks	Total	Comments
1(a)	Gradient, $b = 0.886$ to 0.887 $b = 0.88$ to 0.89	B2 (B1)		AWFW AWFW
	Intercept, $a = 2.31$ to 2.33 $a = 2.3$	B2 (B1)		AWFW AWRT
	Attempt at Σx Σx^2 Σy Σxy or Attempt at S_{xx} S_{xy} Attempt at a correct formula for b $b = 0.886$ to 0.887 $a = 2.31$ to 2.33	(M1) (m1) (A1) (A1)		72, 624, 87, 720 105.6, 93.6 AWFW AWFW
	Accept a & b interchanged only if $y = ax + b$ stated or subsequently used correctly in either (b) or (c)		4	
	(b) a : average waiting time of 2.32 minutes (139 seconds) when entering empty restaurant	B1		OE; accept minimum waiting time
	b : average increase in waiting time of 0.886 minutes (53 seconds) for each customer in restaurant on entry	B1	2	OE
	(c) Use of $y = a + 5b$ or $y = a + 25b$	M1		
	(i) For $x = 5$ $y = 6.6$ to 6.8			
	(ii) For $x = 25$ $y = 24.3$ to 24.6	A1	2	Both; AFWW
	(d)(i) Reliable as interpolation and small residuals or Reliable as interpolation but large percentage residuals so inconclusive or Large percentage residuals so unreliable	B1 B1 (B1) (B1) (B1)		Within range OE OE
(ii) Unreliable as extrapolation	B1	3	Outside range OE	
	Total		11	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
2(a)	$P(X) = 0.3 \quad P(Y) = 0.4 \quad P(Z) = 0.2$			
(i)	$P(X \cap Y \cap Z) = 0.3 \times 0.4 \times 0.2 = 0.024$	M1	1	
(ii)	$P(X' \cap Y' \cap Z') = 0.7 \times 0.6 \times 0.8$ $= 0.336$	M1 A1	2	At least 2 correct terms CAO
(iii)	$P(X' \cap Y' \cap Z) = 0.7 \times 0.6 \times 0.2$ $= 0.084$	M1 A1		Correct numerical expression CAO
(b)	$P(W Z) = 0.9 \quad P(W Z') = 0.25$			
(i)	$P(Z \cap W) = 0.2 \times 0.9$ $= 0.18$	M1 A1	2	Correct numerical expression CAO
(ii)	$P((Z \cap W') \cup (Z' \cap W))$ or $1 - [P((Z \cap W) \cup (Z' \cap W'))]$			
	$= 0.2 \times (1 - 0.9)$ +	M1		0.2×0.9 or (b)(i)
	$(1 - 0.2) \times 0.25$	M1		$(1 - 0.2) \times (1 - 0.25)$
	$= 0.02 + 0.20$ $= 0.22$	A1	3	Cannot score an M1 in both methods $1 - (0.18 + 0.60)$ CAO
	Total		11	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
3(a)	Mean = $\frac{286.5}{50} = 5.73$	B1		CAO
	Standard deviation = $\sqrt{\frac{45.16}{49 \text{ or } 50}} =$ 0.95 to 0.961	B1	2	AWFW
(b)	99% $\Rightarrow z = 2.57$ to 2.58	B1		AWFW 2.5758
	CI for μ is $\bar{x} \pm z \times \frac{(\sigma \text{ or } s)}{\sqrt{n}}$	M1		Use of Must have $(\div\sqrt{n})$ with $n > 1$
	Thus $5.73 \pm 2.5758 \times \frac{(0.95 \text{ to } 0.961)}{\sqrt{50}}$	A1✓		✓ on z and $s^2 > 0$ but not on \bar{x} Accept only 50 or 49 for n
	$5.73 \pm (0.34 \text{ to } 0.36)$	↑		Dependent
	5.37 to 5.39, 6.07 to 6.09)	A1	4	AWFW
(c)	CI excludes both values of 5 and 6½ so Neither claim appears valid	B1✓ ↑ B1✓		✓ on (b); OE Dependent ✓ on (b); OE
	or CI excludes 5 so claim not valid and CI excludes 6½ so claim not valid	(B1✓) (B1✓)	 2	 ✓ on (b); OE ✓ on (b); OE
	Total		8	


MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
4(a)	$\Sigma fx = 8025$ $\Sigma fx^2 = 739975$			
	Mean (\bar{x}) = 80.2 to 80.3	B2		AWFW 80.25
	Standard Deviation (s_n, s_{n-1}) = 30.9 to 31.2 MPs (x): 25, 35, 50, 70, 90, 110, 135, 165	B2 (B1)		AWFW 30.97882 or 31.13489 At least 4 correct
	Mean (\bar{x}) = $\frac{\Sigma fx}{100}$	(M1)	4	Use of
(b)(i)	Large ($n > 30$) sample or Central Limit Theorem	B1	1	OE
(ii)	Mean (\bar{Y}) = 80.2 to 80.3	B1✓		✓ on (a)
	Standard error (\bar{Y}) = $\frac{30.9 \text{ to } 31.2}{\sqrt{36}}$ = 5.1 to 5.25	M1	2	$\sqrt{s^2} > 0$ in (a) ÷ $\sqrt{36}$ or 6
(iii)	$P(\bar{Y} < 90) = P\left(Z < \frac{90 - (80.2 \text{ to } 80.3)}{(5.1 \text{ to } 5.25)}\right)$	M1 M1		Standardising 90 Using values from (b)(ii) with $\sqrt{s^2/36} > 0$ or $\sqrt{s^2/100} > 0$
	= P($Z < 1.84$ to 1.93) = 0.967 to 0.974	A1	3	AWFW
	Total		10	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
5(a)	Scatter Diagram or or	B2 (B1) (B1)	2	4 labelled points plotted 3 labelled points plotted 4 unlabelled points plotted
(b)(i)	Positive/linear correlation/relationship except for two unusual values/results	B1 B1	2	OE OE
(ii)	0.462	B1	1	CAO; accept 3 rd /final/last value
(c)	C and D C is likely freestyle champion D is likely backstroke champion or C is likely freestyle champion D is likely backstroke champion	B1 B1 (B1) (B1)	2	CAO Style identified
(d)(i)	$r = 0.912$ to 0.913 or $r = 0.91$ to 0.92 or 0.46 to 0.47 or $r = 0.9$	B3 B2 B1		AWFW AWFW AWRT
	Attempt at Σx Σx^2 Σy Σy^2 Σxy or Attempt at S_{xx} S_{yy} S_{xy}			270.4, 9188.46 301.6, 11437.84 10246.53
	Attempt at a correct formula for r $r = 0.912$ to 0.913	(M1) (m1) A1	3	48.94, 67.52, 52.45 AWFW
(ii)	Boys are faster/slower at both strokes or Boys are equally good at both strokes	B1	1	OE; do not accept freestyle times are proportional to backstroke times
	Total		11	

Question 5(a)



(a) Scatter Diagram

4 labelled points plotted	B2
3 labelled points plotted	(B1)
4 unlabelled points plotted	(B1)

Graph = 2

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	B(50, 0.2) $P(R \leq 15) = 0.969$ to 0.97	M1 A1	2	Use of in (a) AWFW 0.9692
	(ii) $P(R = 10) = P(R \leq 10) - P(R \leq 9)$ or $P(R = 10) = \binom{50}{10} (0.2)^{10} (0.8)^{40}$ $= 0.5836 - 0.4437 = 0.139$ to 0.141	M1 A1		2
(iii)	$P(5 < R < 15) =$ $P(R \leq 14 \text{ or } 15) = 0.9393$ or 0.9692	M1		Accept values to 3 dp
	minus $P(R \leq 5 \text{ or } 4) = 0.0480$ or 0.0185 $= 0.89$ to 0.893 or B(50, 0.2) expressions stated for at least 3 of $5 \leq R \leq 15$ Answer	M1 M1 A1 (M1) (A2)		3
(b)	Mean, $\mu = np = 50 \times 0.2 = 10$ or Estimate of p , $\hat{p} = 0.21$ Variance, $\sigma^2 = np(1-p) = 10 \times 0.8 = 8$	B1 B1		Either; CAO CAO
	Mean or Estimate of p is similar to that expected but Variance (standard deviation) is different from that expected Reason to doubt validity of Sly's claim	B1 B1		4
	Total		11	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
7 (a)	Weight, $X \sim N(406, 4.2^2)$			
(i)	$P(X < 400) = P\left(Z < \frac{400 - 406}{4.2}\right)$ $= P(Z < -1.428 \text{ to } -1.43)$ $= 1 - P(Z < 1.428 \text{ to } 1.43)$ $= 0.076 \text{ to } 0.077$	M1 m1 A1	3	Standardising (399.5, 400 or 400.5) with 406 and ($\sqrt{4.2}$, 4.2 or 4.2^2) and/or (406 - x) $\Phi(-z) = 1 - \Phi(z)$ AWRT 0.07636
(ii)	$P(402.5 < X < 407.5) =$ $P(X < 407.5) - P(X < 402.5) =$ $P(Z < 0.36) - P(Z < -0.83)$ $= 0.64058 - (1 - 0.79673) = 0.433 \text{ to } 0.44$	M1 B2,1 A1	4	Difference OE AWRT; ignoring signs AWFW 0.43731
(b)(i)	$0.975 \Rightarrow z = 1.96$ $P(Y < 310) = P\left(Z < \frac{310 - \mu}{\sigma}\right)$ <p>or</p> $x = \mu + / \pm z\sigma$ <p>Thus $\frac{310 - \mu}{\sigma} = 1.96 \Rightarrow$ result</p> <p>or</p> $310 = \mu + 1.96\sigma \Rightarrow$ result <p>NB: Working backwards from given equation \Rightarrow at most M1 M0 mo</p>	M1 M1 m1	3	Accept explanation in words Standardising 310 using μ and σ Accept in words Equating AG Substitution
(ii)	$0.86 \Rightarrow z = 1.08$ $310 - \mu = 1.96\sigma$ $307.5 - \mu = 1.08\sigma$ $2.5 = 0.88\sigma$ $\sigma = 2.84 \text{ to } 2.842$ $\mu = 304.4 \text{ to } 304.5$	B1 M1 A1 A1	4	AWRT 1.0803 Attempt at solving 2 equations each of form $x - \mu = z\sigma$ AWFW 2.841 AWFW 304.43
	Total		14	
	TOTAL		75	