

Version 1.0



**General Certificate of Education (A-level)
January 2012**

Mathematics

MS2B

(Specification 6360)

Statistics 2B

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2012 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MS2B

Question	Solution	Marks	Total	Comments
1(a)	21.05 and 21.15	B1	1	both (allow 21.049 and 21.149)
(b)	$E(X) = 0$ (symmetry)	B1		For $R[-a, a]$: $E(X) = 0$ iff $a = 0.05, 0.1, 0.5$ then:
	$\text{Var}(X) = \frac{1}{12}(0.05 - -0.05)^2 = \frac{1}{12} \times \frac{1}{100}$	M1		$\text{Var}(X) = \frac{1}{12}(a - -a)^2$ or their $a = 0.049$ to 0.05 used for M1
	$\Rightarrow sd(X) = \sqrt{\frac{1}{12} \times \frac{1}{100}} = \frac{1}{20\sqrt{3}}$	A1	3	or $\frac{\sqrt{3}}{60}$ or $\sqrt{\frac{1}{1200}}$ 0.0289 (3sf) A0
(c)	$P(-0.01 \leq X \leq 0.03) = 0.04 \times 10 = 0.4$	B1	1	cao from correct value used $\int_{-0.01}^{0.03} 10dx = [10x]_{-0.01}^{0.03} = 0.4$ oe
	Total		5	

MS2B (cont)

Question	Solution	Marks	Total	Comments
2(a)(i)	$H_0: \mu = 61.4$	B1		(both)
	$H_1: \mu \neq 61.4$			
	$z_{calc} = \frac{65.0 - 61.4}{7.5 / \sqrt{16}}$	M1		Alternative:
	$= 1.92$	A1		$P(\bar{X} > 65.0) = P(Z > 1.92)$
	$z_{crit} = \pm 1.96$			$= 1 - 0.97257$
	or (shown in / implied by diagram)	B1		$= 0.02743$
				$\geq 0.025 \quad \therefore$ Accept H_0
	Accept H_0	Adep1		Use of $t \Rightarrow$ max(B1M1A1)
	Insufficient / No evidence (at 5% level) to suggest / show mean (age has) changed (from 61.4 years.)			dep(B1M1) but not A1B1
	Mean (age) has not changed at 1% level (of significance)	Edep1	6	If incorrect or no hypothesis then B0 \Rightarrow max(M1A1B1) i.e. final Adep1Edep1 not available
(ii)	$61.4 - 3 \times 7.5 = 38.9 > 25$ \Rightarrow none under the age of 25 years. Very unlikely any members < 25 yrs.	B1	1	dep(Adep1) $z = \frac{25 - 61.4}{7.5} = -4.85$ $\Rightarrow P(Z < -4.85) \approx 0$ \Rightarrow none aged under 25 included
(b)(i)	$\bar{y} = \frac{\sum y}{n} = \frac{702}{12} = 58.5$ $s^2 = \frac{\sum (y - \bar{y})^2}{n - 1} = \frac{88.25}{11} = 8.02$	B1		($s = 2.83$) $\left(\sigma^2 = 7.35 \text{ or } \sigma = 2.71 \right)$ iff $\sigma / \sqrt{11}$ used below
	$t_{crit} = \pm 1.796$	B1		Ignore signs for t_{crit} If z used then max(B1B1B0M0A0)
	90% CI for μ : $58.5 \pm 1.796 \times \frac{s}{\sqrt{12}}$ 58.5 ± 1.4685 $= 57.03, 59.97$	M1		(their \bar{y}) $\pm t_{11} \times \frac{(\text{their } s)}{\sqrt{12}}$ OR (their \bar{y}) $\pm t_{11} \times \frac{(\text{their } \sigma)}{\sqrt{11}}$
	$= (57.0, 60.0)$	A1	5	
(ii)	upper limit < 61.4 \Rightarrow recruitment drive lowered the average age of the club membership	B1ft	1	Must refer to 61.4 (on their CI)
	Total		13	

MS2B (cont)

Question	Solution	Marks	Total	Comments									
3(a)(i)	$E_i: \frac{mp}{N}; \frac{mq}{N}; \frac{np}{N}; \frac{nq}{N}$	B2,1	2	B1 any one correct B2 all correct (simplified)									
(ii)	$\left. \begin{aligned} \sum_i E_i &= \frac{mp + mq + np + nq}{N} \\ &= \frac{m(p+q)}{N} + \frac{n(p+q)}{N} \text{ (oe)} \end{aligned} \right\}$ $= \frac{mN}{N} + \frac{nN}{N}$ $= m + n$ $= N$ (since $p + q = m + n = N$)	M1 Mdep1 Adep1	3	$\sum_i E_i = \frac{mp + mq + np + nq}{N}$ $= \frac{m(p+q) + n(p+q)}{N}$ (or use of unsimplified forms) $= \frac{(p+q)(m+n)}{N} = \frac{N \times N}{N} = N$ (AG)									
(b)	H_0 : No association between Andy's results and wind conditions E_i : <table border="1" style="margin-left: 20px;"> <tr> <td>17.82</td> <td>15.18</td> <td>33</td> </tr> <tr> <td>9.18</td> <td>7.82</td> <td>17</td> </tr> <tr> <td>27</td> <td>23</td> <td>50</td> </tr> </table>	17.82	15.18	33	9.18	7.82	17	27	23	50	B1 M1		Attempt E's
17.82	15.18	33											
9.18	7.82	17											
27	23	50											
	$\Rightarrow 0_i - E_i - 0.5 = 2.32$	M1		Yates' correction attempted									
	$X^2 = 0.3020 + 0.3546 + 0.5863 + 0.6883 = 1.93$	M1 A1		Final column attempted awrt									
	$\chi_{10\%}^2(1) = 2.706$	B1		correct value of χ^2 only (allow 2.71)									
	\Rightarrow Accept H_0	Adep1		dep (B1 for H_0)									
	No association (between Andy's results and wind conditions)	Edep1	8	Appropriate conclusion dep(B1 for H_0 ; M1 final column; $\chi_{10\%}^2 = 2.706$)									
Total			13										
(a)(ii)	An example of unsimplified values derived from $a = \frac{mp}{N}$: $\Rightarrow b = m - \frac{mp}{N}; c = p - \frac{mp}{N};$ $d = n - \frac{mp}{n}$ (oe)												

MS2B (cont)

Question	Solution	Marks	Total	Comments
4(a)(i)	Poisson	B1	1	
(ii)	$E(3X - 1) = 3\lambda - 1$ $\text{Var}(3X - 1) = 9\lambda$	B1 B1	2	oe (allow $3^2\lambda$)
(iii)	$P(X = x + 1) = \frac{e^{-\lambda} \times \lambda^{x+1}}{(x+1)!}$ $P(X = x + 1) = \frac{e^{-\lambda} \times \lambda^{x+1}}{(x+1)!}$ $= \frac{e^{-\lambda} \times \lambda^x \times \lambda}{(x+1)x!} \left. \vphantom{\frac{e^{-\lambda} \times \lambda^x \times \lambda}{(x+1)x!}} \right\}$ $= \frac{\lambda}{x+1} \times \frac{e^{-\lambda} \times \lambda^x}{x!} \left. \vphantom{\frac{\lambda}{x+1} \times \frac{e^{-\lambda} \times \lambda^x}{x!}} \right\}$ $= \frac{\lambda}{x+1} P(X = x)$	B1 Mdep1 Adep1	3	dep(B1) AG
(b)(i)	$\lambda_{\text{car}} = 500/\text{hour}$ $\lambda_{\text{coach}} = 10/\text{hour}$ $\Rightarrow \lambda_{\text{vehicle}} = 510/\text{hour} = 8.5/\text{min}$ $P(V \geq 10) = 1 - 0.6530$ $= 0.347$	B1 M1 A1	3	for 8.5 stated / used special case: $\lambda = 10 \Rightarrow$ B1M0A0 B1 $\Rightarrow 1 - 0.458$ or 0.542
(ii)	$\mu_{\text{car}} = 836/\text{hour}$ $\mu_{\text{coach}} = 22/\text{hour}$ $\Rightarrow \mu_{\text{vehicle}} = 858/\text{hour} = 14.3/\text{min}$ $P(V \leq 3) = P(V = 0, 1, 2, 3)$ $= \begin{cases} e^{-14.3} \left[1 + \frac{14.3}{1} + \frac{14.3^2}{2} + \frac{14.3^3}{6} \right] \\ e^{-14.3} \times 604.91283 \\ 0.0003726 \text{ to } 0.000373 \end{cases}$ $= 0.00037$ (2sf)	B1 M1 Adep1	3	for 14.3 stated /used All 4 terms required for any $\lambda > 0$ M0 for use of normal approximation dep M1
Total			12	

MS2B (cont)

Question	Solution	Marks	Total	Comments																		
5(a)	<table border="1"> <thead> <tr> <th>n</th> <th>Outcome</th> <th>$P(N = n)$</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>H</td> <td>0.5 ($\frac{1}{2}$)</td> </tr> <tr> <td>2</td> <td>TH</td> <td>0.25 ($\frac{1}{4}$)</td> </tr> <tr> <td>3</td> <td>TTH</td> <td>0.125 ($\frac{1}{8}$)</td> </tr> <tr> <td>4</td> <td>TTTH</td> <td>0.0625 ($\frac{1}{16}$)</td> </tr> <tr> <td>5</td> <td>TTTTA</td> <td>0.0625 ($\frac{1}{16}$)</td> </tr> </tbody> </table> $E(N) = \left(1 \times \frac{1}{2}\right) + \left(2 \times \frac{1}{4}\right) + \left(3 \times \frac{1}{8}\right) + \left(4 \times \frac{1}{16}\right) + \left(5 \times \frac{1}{16}\right)$ $= \frac{1}{2} + \frac{2}{4} + \frac{3}{8} + \frac{4}{16} + \frac{5}{16} = \frac{31}{16}$ $= 1\frac{15}{16} \quad (1.9375)$	n	Outcome	$P(N = n)$	1	H	0.5 ($\frac{1}{2}$)	2	TH	0.25 ($\frac{1}{4}$)	3	TTH	0.125 ($\frac{1}{8}$)	4	TTTH	0.0625 ($\frac{1}{16}$)	5	TTTTA	0.0625 ($\frac{1}{16}$)	B2,1		B1 for one correct entry for $n = 1, 2, 4$ B2 for all 3 correct Can be implied by correct $E(N)$
n	Outcome	$P(N = n)$																				
1	H	0.5 ($\frac{1}{2}$)																				
2	TH	0.25 ($\frac{1}{4}$)																				
3	TTH	0.125 ($\frac{1}{8}$)																				
4	TTTH	0.0625 ($\frac{1}{16}$)																				
5	TTTTA	0.0625 ($\frac{1}{16}$)																				
		M1		$\sum_{n=1}^{n=5} n \times P(N = n)$ (all 5 terms attempted /seen/ implied)																		
		A1	4	(awfw 1.93 to 1.94)																		
(b)	<table border="1"> <thead> <tr> <th>m</th> <th>Outcome</th> <th>$P(M = m)$</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>H</td> <td>$\frac{1}{4}$</td> </tr> <tr> <td>2</td> <td>TH</td> <td>$\frac{3}{4} \times \frac{1}{4} = \frac{3}{16}$</td> </tr> <tr> <td>3</td> <td>TTH</td> <td>$\left(\frac{3}{4}\right)^2 \times \frac{1}{4} = \frac{9}{64}$</td> </tr> <tr> <td>4</td> <td>TTTH</td> <td>$\left(\frac{3}{4}\right)^3 \times \frac{1}{4} = \frac{27}{256}$</td> </tr> <tr> <td>5</td> <td>TTTTA</td> <td>$\left(\frac{3}{4}\right)^4 \times 1 = \frac{81}{256}$</td> </tr> </tbody> </table>	m	Outcome	$P(M = m)$	1	H	$\frac{1}{4}$	2	TH	$\frac{3}{4} \times \frac{1}{4} = \frac{3}{16}$	3	TTH	$\left(\frac{3}{4}\right)^2 \times \frac{1}{4} = \frac{9}{64}$	4	TTTH	$\left(\frac{3}{4}\right)^3 \times \frac{1}{4} = \frac{27}{256}$	5	TTTTA	$\left(\frac{3}{4}\right)^4 \times 1 = \frac{81}{256}$			(given) (given)
m	Outcome	$P(M = m)$																				
1	H	$\frac{1}{4}$																				
2	TH	$\frac{3}{4} \times \frac{1}{4} = \frac{3}{16}$																				
3	TTH	$\left(\frac{3}{4}\right)^2 \times \frac{1}{4} = \frac{9}{64}$																				
4	TTTH	$\left(\frac{3}{4}\right)^3 \times \frac{1}{4} = \frac{27}{256}$																				
5	TTTTA	$\left(\frac{3}{4}\right)^4 \times 1 = \frac{81}{256}$																				
		B3,2,1	3	(B1 any one correct) (B2 any 2 correct) (B3 all 3 correct)																		
(c)(i)	$P(J, R):$ $P(1,1) = \frac{1}{2} \times \frac{1}{4} = \frac{1}{8} \quad (\text{oe})$ $P(2,2) = \frac{1}{4} \times \frac{3}{16} = \frac{3}{64} \quad (\text{oe})$ $P(3,3) = \frac{1}{8} \times \frac{9}{64} = \frac{9}{512} \quad (\text{oe})$ $P(4,4) = \frac{1}{16} \times \frac{27}{256} = \frac{27}{4096} \quad (\text{oe})$ $P(5,5) = \frac{1}{16} \times \frac{81}{256} = \frac{81}{4096} \quad (\text{oe})$	M1		e.g 0.125 attempt at any $P(n,n)$																		
		A1		any 1 correct to 3sf																		
		A1		all 5 correct to 3sf																		
	$p = \sum_{n=1}^{n=5} P(n,n)$	m1		$\sum_{n=1}^{n=5} P(n,n)$ with all 5 values attempted																		
	$\Rightarrow p = \frac{221}{1024} \quad (0.2158)$	A1	5	(awfw 0.215 to 0.217)																		
(ii)	$= 3 \times \left(\frac{221}{1024}\right)^2 \times \left(\frac{803}{1024}\right)$	M1		(either term with their p used) ($0 < p < 1$)																		
	$+ \left(\frac{221}{1024}\right)^3$	M1		(second term with their p used) ($0 < p < 1$)																		
	$P(X \geq 2) = P(X = 2) + P(X = 3)$ $= 0.120 \quad (3\text{dp})$	Mdep1		dep (M1M1)																		
		A1	4	(allow 0.119; 0.12; 0.121)																		
	Total		16																			

MS2B (cont)

Question	Solution	Marks	Total	Comments
6(a)		B2,1	2	B2 for st. line from (1,0.2) to (5,0.3) B1 st. line ($m > 0$) from $x = 1$ to $x = 5$.
(b)	$E(X) = \frac{1}{40} \int_1^5 x(x+7) dx$ $= \frac{1}{40} \left(\frac{x^3}{3} + \frac{7x^2}{2} \right)_1^5$ $= \frac{1}{40} \left(\frac{125}{3} + \frac{175}{2} - \frac{1}{3} - \frac{7}{2} \right)$ $= 3\frac{2}{15}$	M1 A1 A1	3	Ignore limits Ignore limits cao (accept 3.133̇ or $\frac{47}{15}$ oe <i>exact</i>)
(c)	$F(x) = \int_1^x \frac{1}{40}(x+7) dx$ $= \frac{1}{40} \left[\frac{x^2}{2} + 7x \right]_1^x$ $= \frac{1}{80}(x^2 + 14x - 1 - 14)$ $= \frac{1}{80}(x^2 + 14x - 15)$ $= \frac{1}{80}(x+15)(x-1)$	M1 A1 Adep1 Adep1	4	$F(x) = \int \left(\frac{x}{40} + \frac{7}{40} \right) dx$ $= \frac{x^2}{80} + \frac{7x}{40} + c \Rightarrow \text{(M1A1)}$ $F(1) = 0 \Rightarrow c = -\frac{1}{80} - \frac{7}{40} = -\frac{15}{80}$ or [use of $F(5) = 1$] $\Rightarrow F(x) = \frac{1}{80}(x^2 + 14x - 15)$ $F(x) = \frac{1}{80}(x+15)(x-1) \text{ (AG)}$
(d)(i)	$P(2.5 \leq X \leq 4.5) = F(4.5) - F(2.5)$ $= \frac{1}{80}(19.5 \times 3.5 - 17.5 \times 1.5)$ $= \frac{42}{80} = \frac{21}{40} \text{ (0.525)}$	M1 A1	2	Trapezium Rule $\frac{1}{2} \left(\frac{23}{80} + \frac{19}{80} \right) \times 2$ $= \frac{42}{80} = \frac{21}{40}$
(ii)	$F(m) = \frac{1}{2}$ $\Rightarrow \frac{1}{80}(m^2 + 14m - 15) = \frac{1}{2}$ $(\times 80) \Rightarrow m^2 + 14m - 15 = 40$ $m^2 + 14m - 55 = 0$	B1 M1 Adep1	3	$\int_1^m \frac{1}{40}(x+7) dx = 0.5 \text{ (B1)}$ Correct equation formed AG
(e)	$m = \frac{-14 \pm \sqrt{196 + 220}}{2} = \frac{-14 \pm 20.396}{2}$ $m = \frac{-14 + 20.396}{2} \text{ (since } m > 1)$ $m = 3.198 \text{ (3dp)}$	M1 A1	2	Correct attempt at solving quadratic (by formula, oe). cao
	Total		16	
	TOTAL		75	