

**UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
International General Certificate of Secondary Education

**MARK SCHEME for the October/November 2011 question paper  
for the guidance of teachers**

**0580 MATHEMATICS**

**0580/22**

Paper 2 (Extended), maximum raw mark 70

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

<b>Page 2</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2011</b>	<b>0580</b>	<b>22</b>

### Abbreviations

cao	correct answer only
cso	correct solution only
dep	dependent
ft	follow through after error
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
www	without wrong working

Qu.	Answers	Mark	Part Marks
1	35	2	<b>M1</b> for $4 \times 8 + 3$ or $4 \times 8\frac{3}{4}$ or $4 \times 8\frac{1}{2} + 1$ or $\frac{525}{15}$ or $\frac{510}{15} + 1$ <b>SC1</b> for answer 34
2	$(a + b)(x + y)$	2	<b>M1</b> $x(a + b) + y(a + b)$ or <b>M1</b> $a(x + y) + b(x + y)$
3	(a) 6.25 cao (b) 0.16 cao	1 1	
4	12375 cao 12825 cao	2	<b>B1, B1</b> If no marks scored give <b>M1</b> for 27.5 and 28.5 seen
5	$2\frac{1}{12}$ cao <b>with correct working</b>	3	<b>M1</b> $(1+)\frac{6}{12} + \frac{4}{12} + \frac{3}{12}$ oe <b>A1</b> $(1)\frac{13}{12}$ or $\frac{25}{12}$ oe
6	37.5	3	<b>M1</b> $F = k/d^2$ <b>A1</b> $k = 600$
7	$a = -3$ $b = 4$	3	<b>M1</b> $-3a + 4b = 25$ <b>B1</b> one correct
8	11.3	3	<b>M2</b> $22 \times 1.852 \times 1000/3600$ oe or <b>M1</b> $22 \times$ figs 1852 or $22 \times 1000/3600$
9	(a) $\sqrt{(2n-1)}$ (b) $\sqrt{57}$ or 7.55	2 1ft	<b>M1</b> $\sqrt{(2n+k)}$ or $2n-1$ From their (a)
10	$\frac{2x+2}{(x+10)(x+4)}$ oe	3	<b>B1</b> common denominator $(x+10)(x+4)$ oe seen <b>B1</b> $3(x+4) - (x+10)$ seen oe
11	(a) -3 (b) 1.5	2 2	<b>M1</b> $1/2^3$ or $2^{-3}$ <b>M1</b> $2^{6n}$ or $6n = 9$
12	80 www	4	<b>M1</b> attempting area under the graph <b>M1</b> large or small car area found correctly <b>Dep M1</b> correct final area statement
13	(a) 52 (b) 322	2 2	<b>M1</b> $OAB$ or $OBA = 38$ or $OCT = 90$ <b>M1</b> $BCT = 38$ or $BCO = 52$

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2011	0580	22

14	$y \leq 5$ $x \geq 2$ $y \geq x$	4	<b>B1</b> each inequality but accept <b>any</b> of the four inequality symbols Final <b>B1</b> all 3 symbols correct
15	<b>(a)</b> $(3, 3\frac{1}{2})$ <b>(b)</b> $\begin{pmatrix} 4 \\ 3 \end{pmatrix}$ <b>(c)</b> Correct perpendicular bisector with arcs	1 1 2	<b>B1</b> line through $(3, 3\frac{1}{2})$ perp to $AB$ <b>B1</b> two sets of correct arcs
16	<b>(a)</b> Petrol cao <b>(b)</b> 72 <b>(c)</b> $\frac{1}{10}$	1 2 2	<b>M1</b> for $360 \times 12 \div 60$ <b>B1</b> $\frac{6}{60}$ or $\frac{3}{30}$ or $\frac{2}{20}$ or 0.1 or 10%
17	<b>(a) (i)</b> $3a + c$ <b>(ii)</b> $2\frac{1}{2}a + \frac{1}{2}c$ oe <b>(b)</b> $D$ marked $\frac{3}{4}$ way along $CB$	2 2 2	<b>B1</b> $AO + OC + CB$ or $-a + c + 4a$ <b>M1</b> $a + \frac{1}{2}$ their <b>(a)(i)</b> <b>B1</b> $D$ on $CB$
18	<b>(a)</b> $2.5 \times 10^5$ <b>(b)</b> $C = 1/(Lw^2)$	3 3	<b>B2</b> 250000 oe or <b>M1</b> correct part value seen <b>M1</b> each correct move
19	<b>(a)</b> correct bisector (through $3\frac{1}{2}, 3\frac{1}{2}$ ) <b>(b)</b> $y = 1\frac{1}{2}x - 5$ oe <b>(c)</b> 3.61	2 3 2	<b>B1</b> correct line <b>B1</b> correct arcs <b>B2</b> $y = 1\frac{1}{2}x + k$ or $y = kx - 5$ $k$ any number or <b>B1</b> $1\frac{1}{2}x + k$ or $kx - 5$ If O scored allow one each for $m = 1\frac{1}{2}$ or $c = -5$ clearly identified in working <b>M1</b> $\frac{1}{2} \times L \times L = 6.5$ or <b>M1</b> $\sqrt{(3^2 + 2^2)}$