

A Level Biology A
H420/02 Biological Diversity

Question Set 15

- 1 DNA can be obtained from a variety of plant and animal cells.
- (a) A group of students tried to purify some DNA from leek cells using the following method. They decided that exact volumes were not necessary.
1. Grind a leek leaf to a fine pulp using a pestle and mortar.
 2. Add salt and cold water and mix again for at least 10 s.
 3. Add protease enzyme and mix again for at least 10 s.
 4. Filter the liquid into a test tube and stand for at least 10 min.
 5. Tilt the test tube and gently pour in ice-cold ethanol.
 6. A white layer of DNA forms between the sample and the ethanol.
 7. Extract the white layer carefully using a glass rod.
- (i) State the purpose of step 1. [1]
- (ii) Suggest why a protease enzyme added in step 3 is needed to purify DNA. [1]
- (iii) The students considered using pineapple juice as a source of protease enzyme. Suggest why this would **not** be an appropriate source of protease when attempting to produce a pure sample of leek DNA. [1]
- (iv) State one important step that the students had left out of their method. [1]
- (v) Name the process described in step 6. [1]
- (b) Genes isolated from DNA can be used in gene therapy.
- Cystic fibrosis (CF) is a disease that could be treated using gene therapy.
- Healthy individuals have a gene that codes for a channel protein, called CFTR, found in the plasma membrane of a variety of cells, including those lining the airways of the lungs.
- People suffering from CF have two copies of a recessive allele and so their cells do not synthesise the correct channel protein.
- The allele that codes for the functioning CFTR protein can be inserted into the DNA of CF sufferers. The cells can then synthesise the correct CFTR protein and function as normal.

- (i) The treatment of cystic fibrosis is described as **somatic** gene therapy. Another type of gene therapy is known as **germ-line** gene therapy.

Complete the table below to show **three** differences between somatic gene therapy and germ-line gene therapy.

Somatic	Germ-line

[3]

- (ii) Some attempts at gene therapy have resulted in changes to the functioning of other genes.

Explain how inserting a new gene into a chromosome could affect the functioning of other genes in that chromosome.

[2]

- (iii) CF occurs when individuals have two copies of a recessive allele.

Huntington's disease is a lethal disease caused by a dominant allele that codes for the protein huntingtin.

Suggest why gene therapy is unlikely to work as a treatment for Huntington's disease.

[1]

Total Marks for Question Set 15: 11

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge