

AS Level Biology A
H020/02 Depth in biology

Question Set 5

1. A study was carried out on butterflies in two different habitats in the north of England. The two habitats were farmland and mixed deciduous woodland.

- Surveys were completed in 1992 and then at four year intervals.
- Data were collected from butterfly transect sites in both habitats. Using this data, the total butterfly population in each habitat was estimated.
- In 2012, the general populations of butterflies in these two habitats reached historical lows as a result of the prolonged cold and wet weather.
- Between 1992 and 2012 the woodland had become overgrown due to lack of active management. In particular the number of open spaces in the woodland had decreased.

(a) The data in Fig. 1 shows the total butterfly populations per square kilometre in both habitats between the years 1992 to 2012.




Fig. 1

(i) Calculate the total percentage decrease in the number of butterflies on farmland between 1992 and 2012.

Show your working. Give your answer to the nearest whole number.

Answer.....

[2]

(ii) Using the data given in Fig. 1, compare the changes in the number of butterflies on farmland and on woodland between 1992 and 2012.

[2]

(iii) Both habitats experienced the same weather conditions.

Suggest a reason for the **difference** in the rates of decline in butterfly numbers in woodland and farmland. [1]

(iv) A student made the following statement:

‘These data show that the change in butterfly numbers was caused by changes in weather conditions in England.’

Comment on the validity of this statement. [2]

(v) State **one** variable that scientists should control when carrying out surveys such as this. [1]

(b) Butterfly species in severe decline on farmland include gatekeeper, large skipper, small copper, wall brown, small tortoiseshell and white-letter hairstreak.

Butterfly species in severe decline in woodland include brown argus, common blue, gatekeeper, holly blue, marbled white, meadow brown, peacock, small copper, small heath, small tortoiseshell and wall brown.

State which habitat you would expect to have greater species richness and give **two** reasons for your answer. [2]

(c) Female silver-washed fritillary butterflies, *Argynnis paphia*, are usually an orange-brown colour. However, a deep olive-green colour can be seen in some females, largely in the south of England.

(i) What is the term given to this type of biodiversity? [1]

(ii) Give one possible benefit to the species of this type of biodiversity. [1]

Total Marks for Question Set 5: 12

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge