

- Additional Assessment Materials
- Summer 2021
- Pearson Edexcel GCE in Chemistry 8CH0
- Resource Set 1 Topic Group 1
- Topics included: Topic 1: Atomic Structure and the Periodic Table Topic 2: Bonding and Structure
- (Public release version)

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Additional Assessment Materials, Summer 2021 All the material in this publication is copyright

© Pearson Education Ltd 2021

General guidance to Additional Assessment Materials for use in 2021

Context

- Additional Assessment Materials are being produced for GCSE, AS and A levels (with the exception of Art and Design).
- The Additional Assessment Materials presented in this booklet are an **optional** part of the range of evidence teachers may use when deciding on a candidate's grade.
- 2021 Additional Assessment Materials have been drawn from previous examination materials, namely past papers.
- Additional Assessment Materials have come from past papers both published (those materials available publicly) and unpublished (those currently under padlock to our centres) presented in a different format to allow teachers to adapt them for use with candidate.

Purpose

- The purpose of this resource to provide qualification-specific sets/groups of questions covering the knowledge, skills and understanding relevant to this Pearson qualification.
- This document should be used in conjunction with the mapping guidance which will map content and/or skills covered within each set of questions.
- These materials are only intended to support the summer 2021 series.

2 This question is about the structure of the atom and isotopes.

The following excerpt is taken from the book *Inorganic Chemistry* by Bailey and Snellgrove, fourth impression 1938.

"Some of the electrons are also contained in the nucleus, whilst the remainder are arranged in rings revolving round the nucleus The two isotopes [of chlorine] have therefore 18 and 20 electrons respectively in the nucleus and 17 [electrons] external to it."

(a) Identify and correct two errors in the excerpt.

(2)

(b) What is the structure of a 1+ ion of the carbon-13 isotope?	(1)
A six protons, six neutrons and five electrons	
B six protons, seven neutrons and six electrons	
C six protons, seven neutrons and five electrons	
D seven protons, six neutrons and six electrons	
(c) (i) State what is meant by the term relative atomic mass .	(2)

(ii) A 5.000 g sample of lithium, containing the two isotopes lithium-6 and lithium-7, was found to contain 0.460 g of the isotope lithium-6.

Calculate the relative atomic mass of lithium for this sample. Give your answer to an appropriate number of significant figures.

lsotope	Relative isotopic mass
Lithium-6	6.015
Lithium-7	7.016

(3)

(d) A mass spectrometer was used to analyse a sample of bromine, Br₂, with only the ⁷⁹Br and ⁸¹Br isotopes present.

Explain why a very small peak occurs at m/z = 80.

(2)


(Total for Question 2 = 10 marks)

- 2 This question is about ionisation energies.
 - (a) (i) Which equation represents the second ionisation of bromine?
 - \square **A** Br(g) + e⁻ \rightarrow Br⁻(g)
 - \square **B** Br⁻(g) + e⁻ \rightarrow Br²⁻(g)
 - \square **C** Br(g) 2e⁻ \rightarrow Br²⁺(g)
 - \square **D** Br⁺(g) e⁻ \rightarrow Br²⁺(g)
 - (ii) Which set of successive ionisation energies is most likely to be associated with the element boron?
 - A 738,1451,7733,10541,13629

(1)

(1)


- **B** 801, 2427, 3660, 25026, 32828
- C 1086,2353,4621, 6223,37832
- D 1402,2856,4578, 7475, 9445



(b) (i) Complete the graph to show how the first ionisation energies of the Period 3 elements change across the period. Precise figures are not required.

(3)





1 This question is about covalent bonds.	
--	--

(a)	State what is meant by the term covalent bond.	

(b) Draw a diagram of the ammonia molecule, clearly showing its shape. Include any lone pairs of electrons and the value of the bond angle.	

(2)

(2)

(c) The dot-and-cross diagram of BF_3 is

00	oo F ox	00
00 F 00 00	В	00 6 F 8 00

What is the bond angle in BF_3 ?

- 🖾 🗛 90°
- 🖾 **B** 107°
- ☑ C 109.5°
- ☑ D 120°

(1)


(d) (i)	Ammonia and boron trifluoride react to form a compound NH ₃ BF ₃ which contains a dative covalent bond. Each of the molecules, NH ₃ and BF ₃ , has a different feature of its electronic structure that allows this to happen. Use these two different features to explain how a dative covalent bond is formed.	
	(2)	
(ii)) During this reaction, the bond angles about the nitrogen atom and the boron atom change.	

State the new H—N—H and F—B—F bond angles.

(2)

(Total for Question 1 = 9 marks)

6 (a) The diagram shows bond angles in ammonia and water.



8 The table shows some information about a selection of elements and compounds.

	Graphene	Graphite	Diamond	Magnesium oxide	Potassium bromide	Iron
Melting temperature /K	>4000	3950	3820	3125	1007	1808
Density /gcm ⁻³	not measured	2.2 to 2.8	3.51	3.58	2.75	7.86
Compressive strength /GPa	not measured	2.3 and 15.3	443	152	15	170

(a) Explain the difference in the melting temperatures of magnesium oxide and potassium bromide.

(3)

(b) Explain why the electrical conductivity of solid potassium bromide is poor but an aqueous solution of potassium bromide is a good electrical conductor.

(2)

		(2)
		(4)
<i>·</i> · ·		
(e)	The compressive strength is a measure of the energy required to break	
	The compressive strength is a measure of the energy required to break some of the bonds within a substance.	
	some of the bonds within a substance.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compres	sive
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	ssive
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compres	ssive
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	
	some of the bonds within a substance. Deduce possible reasons why there are two widely different values for the compressive strength of graphite.	

Total for Test – 40 marks