

A Level Mathematics A

H240/03 Pure Mathematics and Mechanics

Question Set 4

- A cyclist starting from rest accelerates uniformly at 1.5 m s⁻² for 4s and then travels at constant speed.
 - (a) Sketch a velocity-time graph to represent the first 10 seconds of the cyclist's motion.
 - (b) Calculate the distance travelled by the cyclist in the first 10 seconds. [2]
- 2 A particle P projected from a point O on horizontal ground hits the ground after 2.4 seconds.

The horizontal component of the initial velocity of P is $\frac{5}{3}d$ m s⁻¹.

- (a) Find, in terms of d, the horizontal distance of P from O when it hits the ground. [1]
- (b) Find the vertical component of the initial velocity of P. [2]


P just clears a vertical wall which is situated at a horizontal distance dm from O.

(c) Find the height of the wall. [3]

The speed of P as it passes over the wall is $16 \,\mathrm{m\,s^{-1}}$.

3

(d) Find the value of d correct to 3 significant figures. [4]



The diagram shows a small block B, of mass $0.2 \,\mathrm{kg}$, and a particle P, of mass $0.5 \,\mathrm{kg}$, which are attached to the ends of a light inextensible string. The string is taut and passes over a small smooth pulley fixed at the intersection of a horizontal surface and an inclined plane.

The block can move on the horizontal surface, which is rough. The particle can move on the inclined plane, which is smooth and which makes an angle of θ with the horizontal where $\tan \theta = \frac{3}{4}$.

The system is released from rest. In the first 0.4 seconds of the motion P moves 0.3 m down the plane and B does not reach the pulley.

- (a) Find the tension in the string during the first 0.4 seconds of the motion. [4]
- (b) Calculate the coefficient of friction between B and the horizontal surface. [5]

4 In this question the unit vectors i and j are in the directions east and north respectively.


A particle R of mass 2kg is moving on a smooth horizontal surface under the action of a single horizontal force FN. At time t seconds, the velocity $vm s^{-1}$ of R, relative to a fixed origin O, is given by $v = (pt^2 - 3t)i + (8t + q)j$, where p and q are constants and p < 0.

(a) Given that when
$$t = 0.5$$
 the magnitude of **F** is 20, find the value of p. [6]

When t = 0, R is at the point with position vector (2i - 3j) m.

When t = 1, R is at a point on the line L, where L passes through O and the point with position vector $2\mathbf{i} - 8\mathbf{j}$.

5



The diagram shows a ladder AB, of length 2a and mass m, resting in equilibrium on a vertical wall of height h. The ladder is inclined at an angle of 30° to the horizontal. The end A is in contact with horizontal ground. An object of mass 2m is placed on the ladder at a point C where AC = d.

The ladder is modelled as uniform, the ground is modelled as being rough, and the vertical wall is modelled as being smooth.

(a) Show that the normal contact force between the ladder and the wall is
$$\frac{mg(a+2d)\sqrt{3}}{4h}$$
. [4]

It is given that the equilibrium is limiting and the coefficient of friction between the ladder and the ground is $\frac{1}{8}\sqrt{3}$.

(b) Show that
$$h = k(a+2d)$$
, where k is a constant to be determined. [7]

Total Marks for Question Set 4: 50 Marks



OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge