

A student has two radiators in her home. They are filled with different liquids and have different power ratings.

Fig. 1.1 shows information about the two heaters.



Oil radiator	Water radiator
 <p>Heater contains 10 kg of oil</p>	 <p>Heater contains 10 kg of water</p>
1000 W heater	1500 W heater

Fig. 1.1

Table 1.1 shows information about oil and water.

Material	Specific heat capacity (J/kg °C)	Freezing point (°C)	Boiling point (°C)
Oil	1 700	-24	250
Water	4 200	0	100

Table 1.1

- (a) The student's conservatory can be very cold. Sometimes the temperature can get as low as -6°C .

She thinks that it may be better to use the oil radiator in the conservatory than the water radiator.

Suggest why.

Use the information in Table 1.1 to help you answer.

At -6°C , water is a solid yet oil is a liquid, liquids dissipate heat better than solids. [1]

- (b) Both radiators have a 'cut-out' which prevents them getting hotter than 60°C .

Suggest why.

IF the temperature were to increase beyond 60°C , there is a risk to the students. Nearly boiling oil or water can cause burn if accidentally contacted by the students. [1]

or

there is a risk of water or oil volume expanding as 60°C is close to their boiling points.

(c) The student knows that the oil heater produces 800 J of energy each second.

Calculate the energy produced by the oil heater in 10 minutes.

$$10 \text{ mins} = 10 \times 60 \text{ seconds} = 600 \text{ seconds}$$

$$\begin{array}{l} 800 \text{ J} : 1 \text{ second} \\ \times 600 \quad \downarrow \\ 480,000 \text{ J} : 600 \text{ seconds} \end{array}$$

$$\text{Answer} = \dots\dots\dots 480,000 \dots\dots\dots \text{ J}$$

[2]

(d) The student wants the oil heater to heat up by 40°C

(i) How much energy is needed?

Use the information in **Fig. 22.1** and **Table 22.1** to help you answer.

Show your working.

$$\begin{array}{l} E = mc \Delta\theta \\ \Delta\theta = 40^\circ\text{C} \\ m = 10 \text{ kg} \\ c = 1700 \end{array}$$

$$E = 10 \times 1700 \times 40 = 680000 \text{ J}$$

$$\text{Answer} = \dots\dots\dots 680,000 \dots\dots\dots \text{ J}$$

[2]

(ii) She supplies enough energy to heat up the oil radiator by 40 °C but it only heats up to 32 °C.

Suggest **two** reasons why.

[2]

Because a portion of the energy will be lost to the environment, reducing the temperature increase. (less energy available)

Another portion of the energy will be used for heating the radiator itself, thus less energy is available for the oil to be 40°C.

Total Marks for Question Set 22: 8

Equations in physics

$$(\text{final velocity})^2 - (\text{initial velocity})^2 = 2 \times \text{acceleration} \times \text{distance}$$

$$\text{change in thermal energy} = \text{mass} \times \text{specific heat capacity} \times \text{change in temperature}$$

$$\text{thermal energy for a change in state} = \text{mass} \times \text{specific latent heat}$$

$$\text{energy transferred in stretching} = 0.5 \times \text{spring constant} \times (\text{extension})^2$$

$$\text{potential difference across primary coil} \times \text{current in primary coil} = \text{potential difference across secondary coil} \times \text{current in secondary coil}$$

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge