

AQA Biology A-level


Topic 5: Energy transfers in and between organisms

Notes



Photosynthesis

Photosynthesis is a reaction in which **light energy** is used to produce **glucose** in plants. The process requires **water** and **carbon dioxide**, with the products being **glucose** and **oxygen**. There are two stages of photosynthesis, these are the **light dependent stage** and the **light independent stage**. The rate of photosynthesis is determined by **carbon dioxide concentration, light intensity and well as temperature**.



Chloroplast diagram

Chloroplasts are the site of photosynthesis and are adapted to photosynthesis in the following ways:

- Contains **stacks of thylakoid membranes called grana** which provides a **large surface area** for the attachment of chlorophyll, electrons and enzymes.
- A network of proteins in the grana **hold the chlorophyll in a very specific manner** to absorb the maximum amount of light.
- The granal membrane has **ATP synthase channels** embedded allowing ATP to be synthesised as well as being selectively permeable allowing the establishment of a proton gradient.
- Chloroplasts contain **DNA and ribosomes** allowing them to synthesise proteins needed in the light dependent reaction.

Light Dependent Reaction:

1. **Photons** of light hit chlorophyll molecules in PSII causing the electrons to become excited. This is called **photoionisation**. The charge separation from this drives the process of photolysis.
2. **Photolysis** is the splitting of water with light. One molecule of water requires 4 photons of light to split. When water is split it produces **1 molecule of oxygen, 4 protons and 4 electrons**. The oxygen either naturally diffuses out through the stomata or is used in respiration. The 4 electrons replace those lost from the chlorophyll, whilst the protons move into the **stroma**, later creating a **proton gradient**.
3. The excited electron then moves down a series of protein complexes. At one of the complexes the energy from the electron is used to pump **4 protons from the stroma to the thylakoid space**.
4. The electron then moves down the chain further to PSI. Here more photons of light are absorbed causing the electron to move back up to a high energy level.
5. The electron then moves along the chain to another complex where the electron combines with a proton to form a **hydrogen atom**. This is then used to **reduce NADP**, forming reduced **NADP**.



6. The pumping of protons across the membrane means that there is now a greater concentration of protons in the thylakoid space than the stroma. As a result a proton gradient forms with a **high concentration in the thylakoid space and a low concentration in the stroma**. The protons move across the membrane by diffusion through a protein known as a **stalked particle**. The movement of these protons drives the process of **photophosphorylation**. The enzyme **ATP synthase** phosphorylates ATP from ADP and Pi.


Light Independent Reaction:

1. **CARBON DIOXIDE FIXATION** - carbon dioxide that has diffused in through the stomata is fixed with **ribulose biphosphate (RuBP)** in a process known as **carboxylation**. The enzyme **Rubisco** is needed in order to do this. A 6 carbon sugar is formed first, however this is very unstable and therefore forms 2 molecules of **glycerate-3-phosphate**.
2. **REDUCTION PHASE** - The 2 molecules of glycerate-3-phosphate contain a **-COOH group** and is therefore an **acid**. The reducing power of reduced NADP therefore reduces the glycerate-3-phosphate, with energy being provided by **ATP**. This therefore forms 2 molecule of **triose phosphate**. All of the NADP from the light dependent reaction has now been used with only some of the ATP being used.
3. **REGENERATION OF RuBP** - 5 molecules of triose phosphate are used in order to **regenerate 3 molecules of ribulose biphosphate**. The remaining amount of ATP from the light dependent stage is now used.
4. **ORGANIC MOLECULE PRODUCTION** - 2 molecules of triose phosphate can combine to form the intermediate hexose sugar **fructose 1,6 biphosphate** where after it forms molecules of **glucose**.


6 turns of the Calvin Cycle are required in order to produce 1 molecule of glucose per molecule of CO₂

Respiration

Aerobic respiration is the splitting of a **respiratory substrate**, to release carbon dioxide as a waste product. Hydrogen is reunited with atmospheric oxygen with the release of a large amount of energy. **Anaerobic respiration** occurs in the **absence of oxygen**. Respiration is a multi-step process with each step controlled and catalysed by a specific intracellular enzyme. The steps in respiration are detailed below:




1. **Glycolysis** - this is the first process of both aerobic and anaerobic respiration. It takes place in the **cytoplasm of the cell**. Glucose is **phosphorylated** to produce 2 molecules of **pyruvate**, **2 molecules of ATP** and **2 molecules of NADH**. In anaerobic respiration the pyruvate is further converted into lactate with the help of NADH. **Lactate** is then converted back to pyruvate in the liver. From **one**



molecule of glucose 2 molecules of ATP (net), 2 molecules of reduced NAD (NADH) and 2 molecules of pyruvate are formed.

- Link Reaction** - in the link reaction the 2 molecules of pyruvate are **actively transported into the mitochondria**. The enzyme **decarboxylase** then removes a molecule of CO₂ with a hydrogen also being lost, going on to reduce NAD. The **acetate** formed then combines with coenzyme A to form a molecule of **acetyl coenzyme A**. **Per glucose molecule 2 molecules of acetyl coenzyme A are formed and 0 ATP**
- Krebs Cycle** - The Krebs cycle occurs in the **matrix of the mitochondria**. To begin with the acetyl coenzyme A gives the 2 carbon acetate to a 4 carbon molecule already present. The **6 carbon molecule** that is formed then undergoes a series of reactions. Eventually the starting 4 carbon molecule is regenerated to accept another acetate molecule. **The Krebs cycle turns 2 times per molecule of glucose and therefore per molecule of glucose 2 ATP molecules, 6 NADH molecules, 2 FADH molecules and 4 CO₂ molecules are produced.**
- Oxidation Phosphorylation** - the process of oxidative phosphorylation occurs as follows:

- The reduced NAD (NADH) from the Krebs cycle binds to protein Complex I, releasing its **hydrogen atoms as protons and electrons**. The NAD hydrogen carrier then goes back to the Krebs cycle to be used again. Reduced FAD (FADH) binds to Complex II. It also releases its hydrogen atoms as protons and electrons. The **protons move into the mitochondrial matrix** whilst the electrons released go into the **electron chain**.
- The electrons are then passed down a chain of protein complexes (Complex I to IV) each having a **higher affinity than the previous**. In **Complexes I, III and IV** the energy from the electron is used to pump protons. For each hydrogen released by NADH **4 protons are pumped across**. The protons are pumped into the **intermembrane space**.
- After the electrons have pumped across the protons they are **accepted by the final acceptor oxygen**. The electrons combine with a proton to form a hydrogen atom, which then combines with oxygen to form **water**.
- The pumping of the protons from the mitochondrial matrix to the intermembrane space creates a **proton gradient**. The protons therefore move across the membrane through protein channel called a **staked particle**. The proton motive force provides energy for **ATP synthase** to produce **ATP**.



©1999 Addison Wesley Longman, Inc.



Energy and ecosystems

An **ecosystem** includes all the organisms living in a particular area known as the **community** as well as all the non-living elements of that particular environment. The **distribution** and **abundance** of organisms in a **habitat** is controlled by both **biotic factor** (living) e.g. predators, disease and **abiotic factors** (non-living) such as light levels and temperature. Each species has a particular role in its habitat called its **niche** which consists of its biotic and abiotic interactions with the environment.

The **Sun is the source of all energy** in ecosystems with photosynthetic organisms using this to produce their own food. These can be termed **autotrophs** and are producers. Those organisms that cannot synthesise their own food are called **heterotrophs** with all animals being these. Only around **10% of chemical food energy** is passed on between organisms in the food chain. The other **90% is lost** to the surroundings as:

- uneaten parts e.g. the bones.
- decay of dead material e.g. bacteria may decay some material.
- excretion e.g. energy is lost in faeces
- exothermic reactions e.g. heat lost in respiration.

The efficiency of energy transfer between the trophic levels is worked out using the formula:

$$\text{Percentage efficiency} = \frac{\text{energy available after the transfer}}{\text{energy available before the transfer}} \times 100$$

The **biomass** can be measured in terms of **mass of carbon or dry mass of tissue per given area per given time**. The dry mass is used as the wet mass can vary too much. The chemical energy stored in dry biomass can be estimated using calorimetry. This is carried out in a **bomb calorimeter** in which a sample of known mass is burnt in **pure oxygen**. The bomb calorimeter is submerged in water and therefore the change in water temperature can be used to calculate the energy in the sample.

- **Net primary productivity (NPP)** - the rate at which energy is transferred into the organic molecules that make up new plant biomass, that is the chemical energy store in plant biomass after respiratory losses to the environment have been taken into account
- **Gross primary productivity (GPP)** - the rate at which energy is incorporated into organic molecule in the plants in photosynthesis, that is the chemical energy store in plant biomass, in a given area or volume, in a given time
- Therefore, **$NPP = GPP - R$**
- The net primary production is available for plant growth and reproduction as well as to other trophic levels in the ecosystem such as decomposers and herbivores.
- The net production of consumers (**N**) such as animals can be calculated by: **$N = I - (F + R)$** where **I** represents the chemical energy store in ingested food, **F** represents the chemical energy lost to the environment in faeces and urine and **R** represents the respiratory losses to the environment.



Nutrient cycles

Nutrients are recycled within natural ecosystems, as shown by the nitrogen cycle and the phosphorous cycle.

The Nitrogen Cycle:

Nitrogen is an element used in many biological molecules of which there is a **finite** amount on earth. Due to this it must be recycled from dead organisms and waste products. Most of this is carried out by bacteria in the soil. There are four stages of the nitrogen cycle, these are detailed below:

- **Ammonification** where microbes known as **saprobionts** break down organic matter to ammonia in a two stage process. Firstly, proteins are broken down into amino acids with the use of extracellular **protease** enzymes. These are then subsequently broken down further to remove amino groups with the use of **deaminase** enzymes. Saprobionts use the products of decomposition for respiration.
- **Nitrification** where nitrifying bacteria convert **ammonia to nitrite ions**, NO_2^- , in an oxidation reaction, with a **nitrate ion**, NO_3^- , intermediate. Most plants can take in nitrate ions through their roots.
- **Denitrification** where nitrate ions, NO_3^- , are converted to nitrogen gas, N_2 , by the denitrifying bacteria. This process is wasteful and can be prevented from occurring by soil being well **drained** and **aerated**.
- **Nitrogen fixation** where nitrogen gas is fixed into other compounds by bacteria with nitrogen fixing ability. They do so by reducing nitrogen gas to ammonia which subsequently dissolves to form ammonium ions. Nitrogen fixing bacteria live in **root nodules** of leguminous plants. The relationship between nitrogen fixing bacteria and the plant is known as **mutualistic**, as it is beneficial to both organisms.

The Phosphorus Cycle:

Phosphorus like nitrogen is another element found in many biological molecules that needs to be recycled. Plants can take in phosphate ions, PO_4^{3-} , from soil. Phosphate is released from **sedimentary rocks as a result of weathering, as well as through the decay of bones, shells and the excreta of some birds.**

Mycorrhizae are important in facilitating the uptake of **water and inorganic ions** by plants. These are associations between certain types of fungi and the roots of the vast majority of plants. They **increase the surface area** and act as a **sponge holding water and minerals**. As a result a plant can better **resist drought** and take up **inorganic ions more easily**.

Natural and artificial fertilisers are used to replace the nitrates and phosphates lost by **harvesting plants and removing livestock**.

Nitrogen fertilisers greatly increase crop yields and therefore can help to deal with the demands of a growing human population. However they have negative effects on the environment which include **reducing biodiversity, leaching and eutrophication**.



Leaching is the process by which mineral ions, such as nitrate, dissolve in rainwater and are carried from the soil to end up in rivers and lakes. As a result of this **eutrophication** occurs. This provides **algae** in waterways with enough nitrate ions to grow more **rapidly** than it otherwise would do. As a result this can block out light from other plants, causing decay and the use of oxygen in the water way. This eventually leads to the **death of the ecosystem**.

