

Edexcel B Biology A-Level Core Practical 5

Investigate the effect of temperature on membrane permeability







Cell-surface membranes are made up of a **phospholipid bilayer** which makes them **selectively permeable**. This permeability can be changed by different variables, such as **temperature** and **concentration of solvents** like ethanol.

The permeability of a membrane can be measured by using beetroot cells, which contain a purple **pigment** called **betalain**. When the cell-surface membrane has a higher permeability, more pigment leaks out of cells. The permeability can therefore be measured by the **amount of pigment leaked** from beetroot cells into an aqueous solution using a **colorimeter**.

Equipment

- Water baths
- Thermometer
- Distilled water
- Syringe
- Beetroot
- Cork borer
- White tile
- Knife
- Syringe
- Pipette
- Test tubes
- Colorimeter
- Cuvettes
- Forceps

Method

- 1. Cut beetroot into 8 identical cylinders using a cork borer and wipe/rinse to clean off any pigment released as a result.
- 1. Place each of the cylinders of beetroot in 10 ml of distilled water. Place each test tube in a water bath at a range of temperatures between 0 and 70°C.
- Leave the samples for 15 minutes- pigment will leak out of the beetroot. Record the exact temperature of the water bath using the thermometer.
- 3. Remove the test tubes from the water baths and remove the cylinders of beetroot from them. Decant the liquid into clean test tubes.
- 4. Set the colorimeter to a **blue filter** and **zero** using a cuvette with **distilled water**. Filter each sample into a cuvette using **filter paper**.





5. Measure the absorbance for each solution. A higher absorbance indicates higher pigment concentration, and hence a more permeable membrane.

Risk Assessment

Hazard	Risk	Safety Precaution	In emergency	Risk Level
Scalpel	Cuts from sharp object	Cut away from fingers;use forceps to hold sample whilst cutting; keep away from edge of desk	Elevate cuts; apply pressure; seek medical assistance	Low
Broken glass	Cuts from sharp object	Take care when handling glassware; keep away from edge of desk	Elevate cuts; apply pressure; do not remove glass from wound; seek medical assistance	Low
Hot liquids	Scalding	Handle with care; use tongs to remove boiling tubes from water bath; wear eye protection; keep away from edge of desk	Run burn under cold water; seek medical assistance	Low
Ethanol	Irritant/ flammable	Wear eye protection; keep away from naked flames.	Wash eyes and skin with cold water	Low

Graph

• Plot a graph of absorbance against ethanol concentration/temperature.

•

Conclusion

- As the temperature increases, the permeability of the cell-surface membrane also increases. This is because the proteins in the membrane denature as the heat damages the bonds in their tertiary structure. This creates gaps in the membrane so it is easier for molecules to pass through it.
- At low temperatures, phospholipids have little energy and are packed closely together to make the membrane rigid. This causes a decrease in permeability and restricts molecules from crossing the membrane.

NB: At very low temperatures, ice crystals can form which pierce the cell membrane and increase the permeability.





