



Cambridge International AS & A Level

BIOLOGY**9700/22**

Paper 2 AS Level Structured Questions

May/June 2022

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2022 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **21** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

PUBLISHED**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- 3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- 4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
- 5 'List rule' guidance
 For questions that require *n* responses (e.g. State **two** reasons ...):
 - The response should be read as continuous prose, even when numbered answer spaces are provided.
 - Any response marked *ignore* in the mark scheme should not count towards *n*.
 - Incorrect responses should not be awarded credit but will still count towards *n*.
 - Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
 - Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Mark scheme abbreviations:

;	separates marking points
/	alternative answers for the same marking point
R	reject
A	accept
I	ignore
AVP	any valid point
AW	alternative wording (where responses vary more than usual)
ecf	error carried forward
<u>underline</u>	actual word underlined must be used by candidate (grammatical variants accepted)
max	indicates the maximum number of marks that can be given
ora	or reverse argument

Examples of how to apply the list ruleState **three** reasons.... [3]

A	1	Correct	✓	2
	2	Correct	✓	
	3	Wrong	✗	

F	1	Correct	✓	2
(4 responses)	2	Correct	✓	
	3	Correct CON (of 3)	✗ (discount 3)	

B	1	Correct, Correct	✓, ✓	3
(4 responses)	2	Correct	✓	
	3	Wrong	ignore	

G	1	Correct	✓	3
(5 responses)	2	Correct	✓	
	3	Correct Correct CON (of 4)	✓ ignore ignore	

C	1	Correct	✓	2
(4 responses)	2	Correct, Wrong	✓, ✗	
	3	Correct	ignore	

H	1	Correct	✓	2
(4 responses)	2	Correct	✗	
	3	CON (of 2) Correct	(discount 2) ✓	

D	1	Correct	✓	2
(4 responses)	2	Correct, CON (of 2)	✗, (discount 2)	
	3	Correct	✓	

I	1	Correct	✓	2
(4 responses)	2	Correct	✗	
	3	Correct CON (of 2)	✓ (discount 2)	

E	1	Correct	✓	3
(4 responses)	2	Correct	✓	
	3	Correct, Wrong	✓	

I

PUBLISHED

Question	Answer	Marks
1(a)	<p>I ref. to resolution any one from: Fig. 1.1A is scanning (EM), qualified ; e.g. shows, surface / contour / 3-D, views shows (only the), the outer / external, view (of microvilli) idea of good focus with different depths does not have stated feature of TEM (see mp2)</p> <p>Fig. 1.1B is transmission (EM), qualified ; e.g. showing, 2D / flat, views (of, microvilli / mitochondria) image in one plane showing section through, mitochondria / microvilli / cell I thin shows, internal view of cell / contents of cell / AW does not have stated feature of SEM (see mp1)</p> <p>scanning (EM) stated plus feature <u>and</u> transmission (EM) stated plus feature } ; if Fig. 1.1, A / B not identified</p>	1
1(b)	<p>any two from: max 1 if stating to measure one of the sides of the figure (for each image) measure a number of microvilli and calculate a mean ;</p> <p>same units for actual and measured images / example given ; e.g. if microvillus measurement is in mm convert, microvillus measured length to μm / actual length to mm R if extra detail given is incorrect e.g. <u>divide</u> by 1000 for mm to μm</p> <p>divide, microvillus / image, length by, actual length / AW / $1(\mu\text{m})$; A idea that value for, microvillus / image, length is the magnification</p>	2
1(c)(i)	<p>any two from: (movement of substances) against concentration gradient / from low(er) to high(er) concentration ; (protein has) binding site(s) ; specific ; A description e.g. shape complementary to shape of substance <i>can be in context of protein or binding site</i> ref. to (binding causes protein), conformational change / change of shape ;</p>	2

PUBLISHED

Question	Answer	Marks
1(c)(ii)	<p>any two from:</p> <p>1 glucose is moving against a (concentration) gradient ; AW e.g. low to high</p> <p>2 movement powered by inward movement of Na⁺ ; AW</p> <p>3 Na⁺ movement (into cell from gut lumen) down an electrochemical gradient or down a (Na⁺), concentration / diffusion, gradient or by facilitated diffusion ; A diffusion if response includes <i>ref. to</i> the cotransport protein / cotransporter</p> <p>4 AVP ; e.g. <i>idea that</i>, cotransport needed so that <u>all</u> glucose will be removed from gut lumen or a facilitated diffusion of glucose will only be to equilibrium / AW cotransporter has (specific) binding site for glucose I cotransporter has a complementary shape <i>idea that</i> there is no specific active uptake protein for glucose</p>	2
1(c)(iii)	<p>increases / larger / large, surface area (over which glucose can enter cell per unit time) ; A increases / higher / high, SA:V</p> <p><i>idea of</i> (so) more, transport / carrier / cotransport, proteins A more cotransporters or more proteins, located in membrane / AW ; R pump proteins</p> <p>A <i>idea of</i> more space / places, where proteins can transport glucose</p>	2

PUBLISHED

Question	Answer	Marks										
1(d)	<p><i>any four from:</i></p> <table border="1" data-bbox="344 272 1933 1278"> <thead> <tr> <th data-bbox="344 272 1137 336">suggestion max 3</th> <th data-bbox="1137 272 1933 336">explanation</th> </tr> </thead> <tbody> <tr> <td data-bbox="344 336 1137 571"> <p>1 to replace, old / dead / damaged / injured, cells or to, repair / replace, damaged tissue ;</p> <p>I repair cells I replace old tissues</p> </td> <td data-bbox="1137 336 1933 1050" rowspan="3"> <p>6 <i>idea of function</i> in context of mp1 or 2 or 3 ; e.g. repair allows tissues to function</p> <p>specialisation / differentiation, allows, cells / tissue, to function</p> <p>genetically identical so cells can function as a tissue</p> <p><i>look for other approaches e.g.</i> <i>'replace damaged cells that no longer function'</i> is mp1 and mp6</p> <p>.....</p> <p>7 detail of function (intestines) ;</p> <p>e.g. absorption (of, glucose / nutrients) production of digestive enzymes</p> <p>pass absorbed, nutrients / AW, to blood</p> <p>I nutrition</p> </td> </tr> <tr> <td data-bbox="344 571 1137 805"> <p>2 cells are produced that can, differentiate / become specialised (into intestinal cells); I multipotent / AW</p> <p>R if, stated / strongly implied that, <u>all</u> cells produced by mitosis become specialised</p> </td> </tr> <tr> <td data-bbox="344 805 1137 1050"> <p>3 produce cells that are genetically identical ; A clones</p> <p>A identical / same, DNA / genetic material / genetic information</p> </td> </tr> <tr> <td data-bbox="344 1050 1137 1114"> <p>4 growth ; R growth of cells</p> </td> <td data-bbox="1137 1050 1933 1114"> <p>8 idea of development during, infancy / childhood ;</p> </td> </tr> <tr> <td data-bbox="344 1114 1137 1278"> <p>5 renew / replenish / restore number of / AW, stem cells ; I increase in stem cells</p> </td> <td data-bbox="1137 1114 1933 1278"> <p>9 for, continued production / constant supply / AW, of intestinal cells or so that number of stem cells doesn't decrease ;</p> </td> </tr> </tbody> </table>	suggestion max 3	explanation	<p>1 to replace, old / dead / damaged / injured, cells or to, repair / replace, damaged tissue ;</p> <p>I repair cells I replace old tissues</p>	<p>6 <i>idea of function</i> in context of mp1 or 2 or 3 ; e.g. repair allows tissues to function</p> <p>specialisation / differentiation, allows, cells / tissue, to function</p> <p>genetically identical so cells can function as a tissue</p> <p><i>look for other approaches e.g.</i> <i>'replace damaged cells that no longer function'</i> is mp1 and mp6</p> <p>.....</p> <p>7 detail of function (intestines) ;</p> <p>e.g. absorption (of, glucose / nutrients) production of digestive enzymes</p> <p>pass absorbed, nutrients / AW, to blood</p> <p>I nutrition</p>	<p>2 cells are produced that can, differentiate / become specialised (into intestinal cells); I multipotent / AW</p> <p>R if, stated / strongly implied that, <u>all</u> cells produced by mitosis become specialised</p>	<p>3 produce cells that are genetically identical ; A clones</p> <p>A identical / same, DNA / genetic material / genetic information</p>	<p>4 growth ; R growth of cells</p>	<p>8 idea of development during, infancy / childhood ;</p>	<p>5 renew / replenish / restore number of / AW, stem cells ; I increase in stem cells</p>	<p>9 for, continued production / constant supply / AW, of intestinal cells or so that number of stem cells doesn't decrease ;</p>	4
suggestion max 3	explanation											
<p>1 to replace, old / dead / damaged / injured, cells or to, repair / replace, damaged tissue ;</p> <p>I repair cells I replace old tissues</p>	<p>6 <i>idea of function</i> in context of mp1 or 2 or 3 ; e.g. repair allows tissues to function</p> <p>specialisation / differentiation, allows, cells / tissue, to function</p> <p>genetically identical so cells can function as a tissue</p> <p><i>look for other approaches e.g.</i> <i>'replace damaged cells that no longer function'</i> is mp1 and mp6</p> <p>.....</p> <p>7 detail of function (intestines) ;</p> <p>e.g. absorption (of, glucose / nutrients) production of digestive enzymes</p> <p>pass absorbed, nutrients / AW, to blood</p> <p>I nutrition</p>											
<p>2 cells are produced that can, differentiate / become specialised (into intestinal cells); I multipotent / AW</p> <p>R if, stated / strongly implied that, <u>all</u> cells produced by mitosis become specialised</p>												
<p>3 produce cells that are genetically identical ; A clones</p> <p>A identical / same, DNA / genetic material / genetic information</p>												
<p>4 growth ; R growth of cells</p>	<p>8 idea of development during, infancy / childhood ;</p>											
<p>5 renew / replenish / restore number of / AW, stem cells ; I increase in stem cells</p>	<p>9 for, continued production / constant supply / AW, of intestinal cells or so that number of stem cells doesn't decrease ;</p>											

PUBLISHED

Question	Answer	Marks
2(a)(i)	<p>any two from: oral rehydration (salts / solution / therapy) ; A ORS / ORT A rehydration therapy if <i>ref. to 'drinking'</i> stated within response</p> <p>solution of / water with, glucose and salts ; A named example for salts e.g. sodium / potassium / citrate / chloride</p> <p>AVP ; e.g. antibiotics / doxycycline / tetracycline / furazolidone / choramphenicol / sulfaguanidine (<i>standard treatment in some countries</i>) zinc treatment (<i>standard treatment for children in some countries</i>) <i>ref. to isolation</i></p>	2
2(a)(ii)	<p>any two from: decreases, recovery time ; A cured more quickly I helps recovery / reduces severity of disease</p> <p>reduces time of / stops, diarrhoea production (containing pathogen) ;</p> <p>less likely to / AW, contaminate water ; reduced transmission of pathogen ; AW e.g. breaks transmission cycle</p>	2
2(b)(i)	<p><i>allow antibody or mAb or Mab for monoclonal antibody</i></p> <p>I <i>ref. to receptor</i> I active site I combines / bonds any two from: specific / different, (<i>Vibrio</i> / pathogen) antigen-binding sites / binding sites for antigen ; R if antibody described as enzyme A (each) mAb binds to a, specific / particular, (<i>Vibrio</i> / pathogen) <u>antigen</u></p> <p>(each mAb has) specific / different, tertiary structure / variable region(s) / primary structure / sequence of amino acids ;</p> <p>binding site and <u>antigen</u> have complementary shapes ;</p>	2

PUBLISHED

Question	Answer	Marks
2(b)(ii)	<p><i>allow antibody or mAb or Mab for monoclonal antibody</i> any three from:</p> <p>1 person A, has cholera O1 / is infected with <i>V. cholerae</i> O1 <u>and</u> person B, does not have cholera ;</p> <p>A <i>ref.</i> to person A testing positive for cholera O1 <u>and</u> person B testing negative to cholera</p> <p>2 tests are, valid / working, because, controls / bands 4, shows bands ;</p> <p><i>person A, band / colour in area 3 because</i></p> <p>3 anti-O1 (mobile) antibody binds (specific antigen on) <i>Vibrio</i> O1 ; A antigen-antibody (complex) O1 formed</p> <p>4 antigen-antibody (complex) O1 binds to immobilised antibody <i>ref.</i> to area 3 ; R if stated as, band / colour, in area 2</p> <p>5 person B (in test area 2 and 3) no bands form / no colour / no change, because no antibody binds with antigen / no antigen-antibody complexes formed ;</p>	3
2(c)(i)	95 / 95.1 (%) ; R 95.0%	1
2(c)(ii)	<p>I <i>ref. to</i>, cheaper / no need for (highly) trained personnel / easier to use / easier to obtain results / accurate R <u>more</u> accurate</p> <p><i>any one from:</i> quicker to, obtain result / to do the test ; A less time consuming can carry out test in a greater variety of places / AW ; e.g. can be done at home less labour intensive ; AW e.g. only needs one person / can do it yourself easier to test more people at one time / AW ; portable ; AVP ; e.g. <i>idea of</i> easier to transport or distribute (to reach more people) <i>ref. to</i>, safer / less risk, from pathogen available to more people (<i>consequence of cheaper</i>) <i>suggestion that</i> dipsticks are able to identify different, forms / strains</p>	1

PUBLISHED

Question	Answer			Marks	
3(a)	<p>any two from: I ref. to two units v one unit or bigger v smaller (neutral points)</p>			2	
sucrose			fructose		
disaccharide R if incorrect monosaccharides stated		v	monosaccharide ;		
non-reducing sugar ora A described in terms of reaction with Benedict's reagent		or	reducing sugar ; ora		
glycosidic bond present R glucosidic		or	no glycosidic bond present ;		
AVP ; e.g. always ring structure(s) hydrolysed / broken down before used as, a respiratory substrate / an energy source		v v	ring structure or straight chain form can be used immediately as, a respiratory substrate / an energy source		

PUBLISHED

Question	Answer	Marks
3(b)(i)	<p>I three fatty acids A fatty acid residue <i>for fatty acid</i></p> <p>any three from:</p> <p>1 glycerol ;</p> <p>2 ester, bond / linkage, qualified ; e.g. labelled on Fig. 3.2 / three ester bonds / ester bond between (O of) glycerol and (C of carboxyl of) fatty acid</p> <p>3 palmitate is <u>saturated</u> or oleate(s) are, <u>unsaturated</u> / <u>monounsaturated</u> or two fatty acid(s) are, <u>unsaturated</u> / <u>monounsaturated</u>, <u>and</u> one is <u>saturated</u> ;</p> <p>4 (each fatty acid has a) hydrocarbon chain ;</p> <p>5 AVP ; extra detail e.g. <i>hydrocarbon chain</i> palmitate 15C <u>and</u> oleate 17C palmitate, $\text{CH}_3(\text{CH}_2)_{14}$ <u>and</u> oleate, $\text{CH}_3(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7$</p> <p>palmitate 16C and oleate 18C</p> <p><i>unsaturated fatty acid tail(s) / oleate</i> <i>ref. to</i>, presence of one, C=C / (carbon-carbon) double bond not the maximum number of, hydrogen atoms / hydrogens</p> <p><i>saturated fatty acid tail(s) / palmitate</i> <i>ref. to</i> no, C=C / (carbon-carbon) double bond maximum number of hydrogen atoms / hydrogens</p>	3

PUBLISHED

Question	Answer	Marks
3(b)(ii)	<p>any one from:</p> <p><i>idea that provides greater energy per unit volume ;</i></p> <p>A higher calorific value / greater energy density</p> <p>R produces more energy</p> <p>less mass needed to provide same energy / low(er) mass to energy ratio ; ora higher energy to mass ratio</p> <p>low(er) mass to volume ratio ;</p> <p>AVP ; e.g. glycogen can only be stored in liver and muscles glycogen is hydrated / uses water for storage, so mass increases suggestion that more, areas / adipocytes / adipose tissue, in body to store triglycerides (proportionately) more <u>C-H</u> bonds (to supply H for respiration)</p> <p><i>suggestion that triglycerides provides insulation during flight (at heights)</i> more compact</p>	1
3(c)	<u>closed, double</u> (circulation / circulatory system) ;	1

PUBLISHED

Question	Answer	Marks
3(d)	<p>I <i>ref. to metabolism</i> <i>any two from:</i> need more, / large quantities of, oxygen / glucose or more / large quantities, of, oxygen / glucose, delivered / reaches tissues / transported / AW or <i>idea of enough, oxygen / glucose supplied to meet needs ;</i></p> <p><i>idea that heart pumps blood (to tissues more) efficiently ;</i> e.g. forcefully / in greater volumes / more / with higher pressure / faster <i>more oxygenated blood is mp2 only</i></p> <p>suggestion of reason for needing more oxygen or glucose ; I migration e.g. more active fly / wing muscles search for food longer constantly moving / move for longer time</p> <p>generate heat to maintain body temperature, qualified ; e.g. (generally) smaller size / larger SA:V</p> <p><i>ref. to need to meet demands of carbon dioxide removal (from muscle action / respiration) ;</i> <i>idea that more carbon dioxide produced (so must be removed)</i></p>	2

PUBLISHED

Question	Answer	Marks									
3(e)	<p><i>note: only one mark for both pulmonary vessels</i></p> <table border="1" data-bbox="584 268 1688 730"> <thead> <tr> <th data-bbox="584 268 1245 336">function of blood vessel</th> <th data-bbox="1245 268 1688 336">name of blood vessel</th> </tr> </thead> <tbody> <tr> <td data-bbox="584 336 1245 400">carries blood from the heart to the lungs</td> <td data-bbox="1245 336 1688 400" rowspan="2">pulmonary, artery / arteries ; pulmonary, vein / veins]</td> </tr> <tr> <td data-bbox="584 400 1245 464">carries blood to the heart from the lungs</td> </tr> <tr> <td data-bbox="584 464 1245 528">carries blood from the heart to the rest of body</td> <td data-bbox="1245 464 1688 528"><u>aorta</u> ; I artery</td> </tr> <tr> <td data-bbox="584 528 1245 730">carries blood to the heart from the rest of body</td> <td data-bbox="1245 528 1688 730">I superior and inferior or anterior and posterior I vein venae cavae / vena cava ;</td> </tr> </tbody> </table>	function of blood vessel	name of blood vessel	carries blood from the heart to the lungs	pulmonary, artery / arteries ; pulmonary, vein / veins]	carries blood to the heart from the lungs	carries blood from the heart to the rest of body	<u>aorta</u> ; I artery	carries blood to the heart from the rest of body	I superior and inferior or anterior and posterior I vein venae cavae / vena cava ;	3
function of blood vessel	name of blood vessel										
carries blood from the heart to the lungs	pulmonary, artery / arteries ; pulmonary, vein / veins]										
carries blood to the heart from the lungs											
carries blood from the heart to the rest of body	<u>aorta</u> ; I artery										
carries blood to the heart from the rest of body	I superior and inferior or anterior and posterior I vein venae cavae / vena cava ;										

Question	Answer	Marks
4(a)	<p>R if choice given and one is incorrect</p> <p>carbonic acid ;</p> <p>hydrogencarbonate (ions) ; A bicarbonate (ions)</p>	2
4(b)	<p><i>any one from:</i></p> <p>similar / same, active site ;</p> <p>have the same catalytic amino acids ;</p> <p>similar / same, tertiary structure ; A same quaternary structure</p>	1

PUBLISHED

Question	Answer		Marks																		
4(c)	<p><i>allow pre-RNA for primary transcript</i></p> <p><i>either exon or intron statement required for mp2 to 7, one mark each row</i></p> <p><i>any three from:</i></p> <table border="1" data-bbox="344 338 1928 1198"> <thead> <tr> <th data-bbox="344 338 1137 405">exons</th> <th data-bbox="1137 338 1928 405">introns</th> </tr> </thead> <tbody> <tr> <td data-bbox="344 405 1137 472">1 both, are (nucleotide sequences that are) transcribed / form part of the primary transcript ;</td> <td data-bbox="1137 405 1928 472"></td> </tr> <tr> <td data-bbox="344 472 1137 603">2 (DNA/RNA) <u>coding</u> (sequences) R coding genes I exons have the genetic code</td> <td data-bbox="1137 472 1928 603">(DNA/RNA) <u>non-coding</u> (sequences) R non-coding genes</td> </tr> <tr> <td data-bbox="344 603 1137 734">3 (RNA exons) needed / involved / AW, in translation or for formation of, polypeptides / proteins / carbonic anhydrase</td> <td data-bbox="1137 603 1928 734">not involved in translation / AW</td> </tr> <tr> <td colspan="2" data-bbox="344 734 1928 801"><i>if mps 2 and 3 not gained 'exons code for proteins' or ora 1 mark</i></td> </tr> <tr> <td data-bbox="344 801 1137 900">4 not removed, from primary transcript / from RNA / during gene splicing / during RNA splicing</td> <td data-bbox="1137 801 1928 900">removed, from primary transcript / from RNA / during gene splicing / during RNA splicing</td> </tr> <tr> <td data-bbox="344 900 1137 999">5 (join to) form mRNA / contain mRNA codons A remain in mRNA</td> <td data-bbox="1137 900 1928 999">not part of mRNA / in primary transcript only</td> </tr> <tr> <td data-bbox="344 999 1137 1098">6 not involved in regulating activity of genes / AW</td> <td data-bbox="1137 999 1928 1098">may have regulatory roles A other suggestions for role</td> </tr> <tr> <td data-bbox="344 1098 1137 1198">7 AVP e.g. (as part of mRNA) leave the nucleus / go to cytoplasm / go to ribosomes</td> <td data-bbox="1137 1098 1928 1198">AVP e.g. remain in nucleus</td> </tr> </tbody> </table> <p><i>if <u>all</u> are differences and <u>all</u> statements are reversed (i.e. introns for exons and vice versa) max 2 marks</i></p>		exons	introns	1 both, are (nucleotide sequences that are) transcribed / form part of the primary transcript ;		2 (DNA/RNA) <u>coding</u> (sequences) R coding genes I exons have the genetic code	(DNA/RNA) <u>non-coding</u> (sequences) R non-coding genes	3 (RNA exons) needed / involved / AW, in translation or for formation of, polypeptides / proteins / carbonic anhydrase	not involved in translation / AW	<i>if mps 2 and 3 not gained 'exons code for proteins' or ora 1 mark</i>		4 not removed, from primary transcript / from RNA / during gene splicing / during RNA splicing	removed, from primary transcript / from RNA / during gene splicing / during RNA splicing	5 (join to) form mRNA / contain mRNA codons A remain in mRNA	not part of mRNA / in primary transcript only	6 not involved in regulating activity of genes / AW	may have regulatory roles A other suggestions for role	7 AVP e.g. (as part of mRNA) leave the nucleus / go to cytoplasm / go to ribosomes	AVP e.g. remain in nucleus	3
exons	introns																				
1 both, are (nucleotide sequences that are) transcribed / form part of the primary transcript ;																					
2 (DNA/RNA) <u>coding</u> (sequences) R coding genes I exons have the genetic code	(DNA/RNA) <u>non-coding</u> (sequences) R non-coding genes																				
3 (RNA exons) needed / involved / AW, in translation or for formation of, polypeptides / proteins / carbonic anhydrase	not involved in translation / AW																				
<i>if mps 2 and 3 not gained 'exons code for proteins' or ora 1 mark</i>																					
4 not removed, from primary transcript / from RNA / during gene splicing / during RNA splicing	removed, from primary transcript / from RNA / during gene splicing / during RNA splicing																				
5 (join to) form mRNA / contain mRNA codons A remain in mRNA	not part of mRNA / in primary transcript only																				
6 not involved in regulating activity of genes / AW	may have regulatory roles A other suggestions for role																				
7 AVP e.g. (as part of mRNA) leave the nucleus / go to cytoplasm / go to ribosomes	AVP e.g. remain in nucleus																				

PUBLISHED

Question	Answer	Marks						
4(d)	<p><i>allow Hb or hb or Hgb for haemoglobin</i> <i>two marks for points in left column or two marks for points in right column</i></p> <table border="1" data-bbox="349 272 1928 740"> <tr> <td data-bbox="349 272 1137 336">cytoplasm / cytosol ;</td> <td data-bbox="1137 272 1928 336">(bound to) cell surface membrane ;</td> </tr> <tr> <td data-bbox="349 336 1137 539">close to haemoglobin (for H⁺ uptake) R in / on / with, hb or <i>idea of being dispersed within cell because haemoglobin is distributed throughout ;</i></td> <td data-bbox="1137 336 1928 539">close to, anion exchanger / transport protein, for (easy) exit of hydrogen carbonate (ions) ;</td> </tr> <tr> <td data-bbox="349 539 1137 740">quick for hydrogen carbonate (ions) to be transported out of cell / AW or H⁺ (from dissociation) binds rapidly to haemoglobin or cytoplasm provides water for reaction to form carbonic acid ;</td> <td data-bbox="1137 539 1928 740">able to, hydrate / react with, carbon dioxide as it enters cell ;</td> </tr> </table> <p><i>if stated cytoplasm <u>and</u> cell surface membrane, one mark</i> <i>reason can be taken from either column if correctly linked, one mark</i></p>	cytoplasm / cytosol ;	(bound to) cell surface membrane ;	close to haemoglobin (for H ⁺ uptake) R in / on / with, hb or <i>idea of being dispersed within cell because haemoglobin is distributed throughout ;</i>	close to, anion exchanger / transport protein, for (easy) exit of hydrogen carbonate (ions) ;	quick for hydrogen carbonate (ions) to be transported out of cell / AW or H ⁺ (from dissociation) binds rapidly to haemoglobin or cytoplasm provides water for reaction to form carbonic acid ;	able to, hydrate / react with, carbon dioxide as it enters cell ;	2
cytoplasm / cytosol ;	(bound to) cell surface membrane ;							
close to haemoglobin (for H ⁺ uptake) R in / on / with, hb or <i>idea of being dispersed within cell because haemoglobin is distributed throughout ;</i>	close to, anion exchanger / transport protein, for (easy) exit of hydrogen carbonate (ions) ;							
quick for hydrogen carbonate (ions) to be transported out of cell / AW or H ⁺ (from dissociation) binds rapidly to haemoglobin or cytoplasm provides water for reaction to form carbonic acid ;	able to, hydrate / react with, carbon dioxide as it enters cell ;							
4(e)	exocytosis ;	1						
4(f)	<p>any three from:</p> <p>1 acetazolamide / inhibitor, binds / attaches / AW, to, allosteric site / site other than active site ; R alternative active site</p> <p>2 (which) changes, shape / tertiary structure / 3-D structure, of active site ; A active site distorted I protein structure</p> <p><i>allow ecf for mp3 and 4 if competitive described (to max 2)</i></p> <p>3 substrate(s) / CO₂ / HCO₃⁻, cannot, enter / bind to / fit in / AW, active site ; A active site no longer complementary to, substrate / AW A enzyme substrate / ES, complexes cannot form A ESCs cannot form I 'substrate / CO₂ / HCO₃⁻ / 'cannot bind to enzyme' <i>without a link to active site</i></p> <p>4 product, not formed / released ; A named product</p> <p>5 <i>ref. to acetazolamide / inhibitor, leaves, allosteric site / enzyme ;</i> I reversibly binds</p>	3						

PUBLISHED

Question	Answer	Marks
5(a)	<p>any one from: I too small presence of cilia ; A ciliated, cells / epithelial cells / epithelium</p> <p>AVP ; e.g. <i>suggestion that</i> mucus on surface makes detail not easily seen</p>	1
5(b)	<p>epithelium has goblet cells and ciliated epithelial cells ; A ciliated cells A two different cell types present</p> <p>I <i>ref. to</i> cell division</p> <p><i>ref. to</i> staining ; e.g. use of stains gives different, shades / colours differential staining one cell type more neutral / acidic different parts of cells stain differently</p> <p><i>idea of</i> cells have different densities ; e.g. <i>idea of</i> more structures / larger nucleus, gives greater density / AW R no cell structures <i>for lighter cell</i> I <i>ref. to</i> number of cilia (sectioning may include) more of nucleus so, affects light penetration / makes cell darker ora</p> <p>AVP ; <i>idea of</i> different components in cells appearing, darker / lighter</p>	2

PUBLISHED

Question	Answer		Marks	
5(c)	<i>any three from:</i>		3	
	A			
	B / cartilage			
	<i>max two for structure</i>			
	smooth muscle / composed of smooth muscle cells or no chondrocytes	or		no smooth muscle cells or has chondrocytes ;
	cells close together	or		cells, (more) isolated / in gaps / in lacunae ;
	cells, spindle shaped / elongated / AW	or		cells (more) rounded / AW ;
	AVP structure e.g. not surrounded by, perichondrium / AW contains blood vessels	or		surrounded by, perichondrium / AW no blood vessels ;
	<i>function</i>			
	contractile / contracts (and relaxes) I recoil A bronchi walls contract R bronchi contract	or		non-contractile ;
changes diameter of airways A controls size of lumen	or	protective / support ; A described e.g. prevents collapse of, bronchus or keeps bronchus open		
<i>if incorrectly quotes A as the cartilage, use right hand column to allow one mark max (ecf)</i>				

PUBLISHED

Question	Answer	Marks
6(a)	<p>any three from:</p> <p>at (phloem) source, mineral ion entry (with water), from xylem / through plasmodesmata from companion cells ;</p> <p>dissolved in water / as solutes in water / in solution / in sap solution ; <i>in context of entry from xylem or within phloem sap</i></p> <p>(transport) as part of mass flow ; <i>must be in context of phloem</i> flow down a pressure gradient / AW ; R differences in pressure gradient R from high pressure gradient to low pressure gradient</p> <p>AVP ; e.g. suggestion that mineral ions are needed for transport of some assimilates (e.g. complexed with) <i>idea of reallocation of resources / arrive at (phloem) source and are moved to other parts, for use / when needed ;</i></p>	3
6(b)	<p>sieve, plates / pores ; A perforated end plates / end plates with pores I plasmodesmata</p> <p>periphery / edge / AW ; companion ; A transfer</p>	3