

Mark Scheme (Results) October 2017

Pearson Edexcel International Advanced Level In Biology (WBI01) Paper 01 Lifestyle, Transport, Genes and Health



Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2017
Publications Code WBI01_01_1710_MS
All the material in this publication is copyright
© Pearson Education Ltd 2017

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question Number	Answer	Additional guidance	Mark
1(a)	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	ACCEPT structural formula for water or 'water' ACCEPT- COOH ACCEPT 2 separate amino acids drawn in box	(2)

Question Number	Answer	Additional guidance	Mark
1(b)	hydrolysis ;	Accept-hydrolisis	(1)

Question	Answer	Additional guidance	Mark
Number			
1(c)	peptide bond ;		(1)

Question	Answer	Additional guidance	Mark
Number			
1(d)	1. active site that { fits /eq } the { substrate /dipeptide };	mp1 accept-connects to, attaches mp1 accept an ESC is formed/eq	
	2. lowers the activation energy (for the reaction);		
	3. detail of how the activation energy is lowered;	mp3 e.g. by weakening or breaking the peptide bond	(2)

Question	Answer	Mark
Number		
2(a)	2(a). The only correct answer is C	
	A is not correct because although membranes contain carbohydrate this plays no role in the diffusion of oxygen	
	B is not correct because although membranes contain cholesterol this plays no role in the diffusion of oxygen	
	D is not correct because oxygen is non-polar	(1)

Question	Answer	Additional guidance	Mark
Number			
2(b)(i)	Alveoli / alveolus / alveolar wall / eq;	Allow description of alveoli e.g. layer of	
		squamous epithelium	(1)

Question	Answer	Mark	l
Number			i
2(b)(ii)	2(b)(ii). The only correct answer is B		1
	A is not correct because surface area is not at its maximum and the thickness is not at its minimum		
	C is not correct because surface area is not at its maximum and the thickness is not at its		
	D is not correct because surface area is not at its maximum and the thickness is not at its	(1)]

Question Number	Answer	Additional guidance	Mark
2(c)(i)	1. 18 (mm) / 1.8 cm	Accept range from 17-19 mm/1.7-1.9 cm Ecf to mp2	
	2. 0.4 (μm) ;	Accept-0.36/0.38/0.4/0.404 Correct answer with no working gains both marks	(2)

Question	Answer	Additional guidance	Mark
Number			
2(c)(ii)	endothelial / (squamous) epithelial cell;	ACCEPT endothelium, (simple / pavement /	
		flattened) epithelium	(1)

Question	Answer	Additional guidance	Mark
Number			
3(a)	1. phospholipids are able to move;	Mp1 accept ref to them being fluid	
	2. proteins are scattered in the membrane / eq;	Mp2 Ignore ref to them forming a mosaic Mp2 Accept-randomly distributed	(2)

Question	Answer	Mark
Number		
3(b(i)	3(b(i).The only correct answer is D	
	A is not correct because this is too low	
	B is not correct because this is too low	
	C in not correct because this is too low	(1)

Question Number	Answer	Additional guidance	Mark
3(b)(ii)	diffusion;	Accept simple diffusion	(1)

Question Number	Answer	Additional guidance	Mark
3(c)(i)	• increase ;		
	• greater/ higher/eq ;		
	• permeable ;	DNA-semi/partially permeable	
	• osmosis :		(4)

Question Number	Answer	Additional guidance	Mark
3(c)(ii)	 (membrane) proteins are denatured by heat / eq; 	Mp1-Accept proteins are damaged/disrupted DNA- destroyed/killed Ignore ref to membrane ONLY	
	 {solutes / cytoplasm / water/eq } leave the {cells/tissues}; 	Mp2-accept cell contents leave only	(2)

Question Number	Answer	Additional guidance	Mark
4(a)	 if energy { intake/input} exceeds energy requirements/eq; idea that excess energy will be stored as 	Mp1 accept energy expenditure/output/used I-ref to calories	
	fat; 3. resulting in an increase in {body mass/BMI};	Mp3 accept weight Ignore obesity	(2)

Question	Answer	Additional guidance	Mark
Number			
4(b)(i)	EITHER	Correct answer with no working gains	
		full marks	
	1. 1.74 - 0.90 = 0.84 ;		
		DNA-48 as a whole number	
	2. (0.84 ÷ 1.74) x 100 = 48.3% ;	DNA-48.2	
		Accept-48.28 and 48.27	
	OR		
	0 4 74 0 00 0 04		
	3. 1.74 - 0.90 = 0.84 ;		
	4 (0.04 0.0) 400 00.000		(0)
	4. $(0.84 \div 0.9) \times 100 = 93.33\%$		(2)

Question Number	Answer	Additional guidance	Mark
4(b)(ii)	reduced levels of selenium (in the diet) are associated with an increase in BMI/obesity (for	ACCEPT converse ACCEPT inverse / negative correlation;	
	both men and women);	- Control of the cont	(1)

Question Number	Answer	Additional guidance	Mark
4(b)(iii)	1. skin fold thickness;		
	2. waist circumference/eq ;	Mp2 accept size/width/diameter	
	3. waist to hip ratio ;	DNA-hip to waist ratio	
	4. body fat mass/body fat %;	DNA-mass/weight only DNA-BMR	
	5. body fat distribution ;		(1)

Question Number	Answer	Mark
4(c)(i)	4(c)(i). The only correct answer is C	
	A is not correct because a correlation is not shown	
	B is not correct because a correlation is not shown	
	D is not correct because a correlation is not shown	(1)

Question Number	Answer	Additional guidance	Mark
4(c)(ii)	idea that smoking causes platelet damage;	Mp1 accept platelets become "sticky"	
	increases the risk of formation of blood clots;		
	 increases the risk of {plaque formation/atheroma/ atherosclerosis}; 		
	4. increases blood pressure;		(3)

Question Number	Answer	Additional guidance	Mark
5(a)	frog and humans have a double circulatory system;	Please not that each MP must be comparative-ie mention both frog and human to gain mark	
	2. frog and human hearts have two atria;	Ignore-3 v 4 chambers	
	 frog and human have pulmonary circuit /carry blood to lungs; 	Tgriore-5 v 4 chambers	
	4. frog and human have a systemic circuit /carry blood around body;	Mp5 accept frog heart has no septum/is not separated while human heart has septum/is	
	5. frog heart has only one ventricle while human heart has two/eq;	separated	
	 in frog oxygenated blood and deoxygenated blood are mixed but not mixed in humans/eq; 		(4)

Question	Answer	Additional guidance	Mark
Number			
5(b)	oxygenated and deoxygenated circuits are kept separate / not mixed/eq;	Accept converse for all mps	
	2. idea that this maintains a steep gas concentration gradient in the lungs/eq;		
	3. (leading to) a fast rate of diffusion of { oxygen / carbon dioxide / gases } in the lungs		(3)

Question Number	Answer	Additional guidance	Mark
*5(c)	atrioventricular valve opens during atrial systole;	QWC emphasis is clarity of expression Paired responses-but mp1/3/5/7 are stand alone marks.	
	2. to allow blood to flow into the ventricle;	Accept named AV valves eg bicuspid and tricuspid	
	atrioventricular valve closes during ventricular systole;	Accept contraction for systole/relaxation for diastole for all mps	
	4. to prevent blood flowing back into the atria;		
	5. semilunar valve opens during ventricular systole;		
	6. to allow blood into the aorta/arteries;		
	7. semilunar closes during diastole;		
	8. to prevent blood from flowing back into the ventricle;		(5)

Question Number	Answer	Additional guidance	Mark
6(a)(i)	 place a Daphnia (in a drop of solution) on a microscope slide/eq; 		
	2. use an appropriate method to hold <i>Daphnia</i> in place;	Mp2 eg cotton wool/ cover slip/vaseline	
	3. time/stated time to acclimatise in the solution (before measurements are taken);		
	4. count number of heart beats in set time period- eg 10 secs-60 secs ;	MpE accept find heats per minute/hpm	
	5. determine heart rate at each time point;	Mp5 accept find beats per minute/bpm	
	6. named control variable; eg temperature, concentration / volume of glucose / lactose, time for acclimatisation	Mp6 eg temperature, concentration / volume of glucose / lactose, time for acclimatisation, same species or source of Daphnia/pH	(4)

Question Number	Answer	Additional guidance	Mark
6(a)(ii)	idea that glucose has very little effect on the heart rate;	ACCEPT glucose increases heart rate slightly	
	idea that overall lactose causes a decrease in heart rate(compared with control);		
	3. lactose has very little effect on heart rate {in the first 16 minutes/up to 16 minutes};		
	 idea that lactose decreases the heart rate {after 16 minutes/from 16 minutes until the end}; 	Eg at end of experiment/90 mins there is a diff of 140 bpm between lactose and	
	credit correct manipulation of figures compared to the control;	control or x 4.5	(3)

Question	Answer	Mark
Number		
6(a)(iii)	6(a)(iii). The only correct answer is B	
	lactose is NOT made from glucose and fructose	
	only lactose contains a glycosidic bond because it is a disaccharide	
	glucose IS a monosaccharide and lactose is NOT a monosaccharide	(1)

Question Number	Answer	Additional guidance	Mark
6(b)(i)	lactose is larger than glucose/converse;	Accept: lactose has a different shape	
	2. lactose fills the opening in the ion channels;	Mp2 accept glucose passes through the ion channel	
		DO NOT ACCEPT lactose blocks the channel	(2)

Question	Answer	Additional guidance	Mark
Number			
6(b)(ii)	1. named disaccharide ;	e.g. sucrose, maltose,	
	2. slows/decreases (heart rate)/same effect as lactose;		
	or		
	3. named monosaccharide;	e.g. fructose, ribose, deoxyribose, galactose,	
	4. has no/little effect (on heart rate)/same effect as glucose;		(2)

Question	Answer	Mark
Number		
7(a)(i)	7(a)(i). The only correct answer is C	
	A is not correct because hexose sugars are not present	
	B is not correct because hexose sugars are not present	
	D is not correct because uracil is not present	(1)

Question	Answer	Mark
Number		
7(a)(ii)	7(a)(ii). The only correct answer is D	
	A is not correct because alleles occur in pairs	
	B is not correct because there will always be SOME effect	
	C is not correct because being heterozygous is irrelevant	(1)

Question Number	Answer	Additional guidance	Mark
7(b)	 parents correctly shown; three genotypes shown in offspring; six genotypes shown as male and female; 	Allow consequential error (incorrect parents)	
		Ignore Punnet squares	(3)

Question	Answer	Additional guidance	Mark
Number			
7(c)(i)	1. there are 20 (different) amino acids;		
	2. three bases gives 64 possibilities;		
	3. two bases gives 16 possibilities;		
	4. idea that three bases is more than enough, two bases would not be enough;		(2)

Question Number	Answer	Additional guidance	Mark
7(c)(ii)	(QWC - Spelling of technical terms must be correct and the answer must be organised in a logical sequence)	QWC emphasis is logical sequence [penalise once only]	
	 double helix/DNA unwinds/unzips/separates/eq; 		
	2. helicase breaks hydrogen bonds;		
	3. (mono) nucleotides line up along the single strands;	Mp4 accept description of complementary base pairing	
	4. complementary base pairing occurs;	eg A-T/G-C	
	5. hydrogen bonds form (between bases);		
	6. DNA polymerase joins the nucleotides together;		
	7. with phosphodiester bonds;		
	8. reference to semiconservative replication / two identical {copies of DNA/daughter strands} are made;		(5)

Question Number	Answer	Additional guidance	Mark
8(a)	idea that prothrombin is the inactive enzyme;	mp1 ignore protein mp1 ignore inactive only	
	prothrombin is converted to { thrombin /an active enzyme};		
	3. by thromboplastin ;		
	4. (thrombin) converts fibrinogen to fibrin;		
	fibrin traps { platelets/red blood cells} to form clot;		(3)

Question	Answer	Additional guidance	Mark
Number			
8(b)	 as prothrombin concentration increases clotting time decreases; 	Mp1 accept negative correlation/inverse relationship	
	2. more prothrombin to convert into thrombin;		
	3. the greater the concentration of thrombin the faster the production of fibrin;		(3)

Question Number	Answer	Additional guidance	Mark
8(c)(i)	presence of prothrombin mutation increases the risk (of heart attack)/converse;	ACCEPT homozygous for no mutation have lowest risk (of heart attack);	
	 homozygous (recessive) for mutation have the { highest risk / higher risk than heterozygotes } (of heart attack); 		
	3. { other blood groups / not group O } increase the risk of heart attack (for all prothrombin genotypes);	3. ACCEPT converse	
	4. credit correct manipulation of figures;		(3)

Question	Answer	Additional guidance	Mark
Number			
8(c)(ii)	 mutation in prothrombin increases the risk of blood clotting; 	Mn2 accort different order/sequence of	
	mutation results in an altered primary structure for prothrombin;	Mp2 accept different order/sequence of amino acids	
	3. active site of thrombin has altered shape/eq;		
	 thrombin produced has increased enzyme activity; 		(3)

DI	1
-	VI.