SurnameCentre
NumberCandidate
NumberOther Names2



GCE AS/A level

1071/01

BIOLOGY/HUMAN BIOLOGY – BY1

A.M. WEDNESDAY, 21 May 2014

1 hour 30 minutes

For Examiner's use only			
Question	Maximum Mark	Mark Awarded	
1.	11		
2.	6		
3.	6		
4.	10		
5.	6		
6.	9		
7.	12		
8.	10		
Total	70		

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use pencil or gel pen. Do not use correction fluid. Write your name, centre number and candidate number in the spaces at the top of this page. Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the continuation pages at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.


You are reminded of the necessity for good English and orderly presentation in your answers.

The quality of written communication will affect the awarding of marks.




BLANK PAGE


PLEASE DO NOT WRITE ON THIS PAGE



1071 010003










1071 010005

Examiner only The nucleus has pores in the envelope that surrounds it, whereas organelle **A** does not. Describe **one** *other* difference between the membranes that surround organelle **A** and (C) those that surround the nucleus. [1] _____ Describe two differences between the ribosomes found in animal cells and those found (d) in prokaryotic cells. [2] 11





Turn over.




(c)	Describe the biochemical test that could be performed to test for a protein. [2]	Examiner only
(d)	Suggest how the concentration of a specific protein could be measured in a sample o urine.	 .f]
		6
		1071 010009
0 9	© WJEC CBAC Ltd. (1071-01) Turn over	_



The diagram below shows a component of DNA .	E
A B C	
(a) Name the parts A , B and C .	[3]
Α	
Β	
С	
(b) Describe how a polymer of DNA would be different from a polymer of RNA.	[2]

(i)	Name the stage in the cell cycle where DNA replication occurs.	[1]	Examin only	er
(ii)	Vincristine is a drug which prevents the spindle fibres from shortening. Name stage in the cell cycle which would be affected.	the [1]		
(iii)	State three differences between daughter cells produced by the process of mite and those produced by meiosis.	osis [3]		
••••••			10	
				1071 010011
	(ii)	 (ii) Vincristine is a drug which prevents the spindle fibres from shortening. Name stage in the cell cycle which would be affected. (iii) State three differences between daughter cells produced by the process of mite 	 (ii) Vincristine is a drug which prevents the spindle fibres from shortening. Name the stage in the cell cycle which would be affected. [1] (iii) State three differences between daughter cells produced by the process of mitosis 	 (i) Name the stage in the cell cycle where DNA replication occurs. [1] (ii) Vincristine is a drug which prevents the spindle fibres from shortening. Name the stage in the cell cycle which would be affected. [1] (iii) State three differences between daughter cells produced by the process of mitosis and those produced by meiosis. [3]





1071 010013

(a)	Using graph I, name the process by which oxygen is absorbed by the roots, giving a reason for your answer. [2]	Examiner only
(b)	Explain why the rate of uptake of nitrate ions increases between points A and B shown on graph II .	
(c)	In the presence of a respiratory inhibitor such as cyanide, the rate of nitrate uptake falls to zero. Name the process by which nitrate ions are taken up. [1]	
(d)	Water enters root hair cells by osmosis. Calculate the solute potential (Ψ_S) of the root hair cell, when there is no net movement of water, the water potential of the soil water is -100 kPa and the pressure potential (Ψ_P) inside the root hair cell is +200 kPa. Use the formula $\Psi = \Psi_S + \Psi_P$. Show your working and units. [2]	1071
	Answer	6

	E
. The	diagram below shows the fluid mosaic model proposed by Singer and Nicolson in 1972.
x	
(a)	The width of the membrane as shown by X has been measured using transmission electron microscopes. Membrane width does not vary greatly between different organisms. State a value for this width. [1]
	Membrane width =
(b)	Glucose is water soluble. Vitamin A is lipid soluble. Describe and explain how each molecule crosses the membrane shown above. [4]
	Vitamin A
	Glucose
1 4	© WJEC CBAC Ltd. (1071-01)



 (c)	Beet	root vacuoles contain a red pigment called <i>betacyanin</i> . When beetroot discs are cut	Examiner only
		a borer and immersed in a solution of 70% ethanol (an organic solvent) at 15°C, the bigment begins to leak out of the cells into the ethanol turning it red.	
	(i)	Using your knowledge of the structure of cell membranes, explain why this leakage of pigment occurs.	
		[2]	
	·····		
	(ii)	When the experiment was repeated at 30°C, the time taken for the ethanol to turn red decreased. Explain why. [2]	
	•••••		
			9



BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE



7.	lame routir The a col		kaminer only
	(a)	Apple pulp added Gel membrane with immobilised pectinase Tap to control flow rate Juice collected Immobilising enzymes can increase the temperature range over which they can be used. Describe two other advantages of immobilising pectinases. [2]	
	(b)	Suggest why reducing the flow rate of material through the column would result in an increased volume of juice being collected. [2]	
	1 7	© WJEC CBAC Ltd. (1071-01) Turn over.	

Examiner only The extraction of juice using pectinase was compared using equal volumes and concentrations of free enzyme, enzymes bound to the surface of a gel membrane and (C) enzymes immobilised inside alginate beads. The results are shown in the graph below. 100 90 80 Volume of juice extracted (cm³) 70 60 50 40 30 20 10 0 30 40 50 20 60 70 **Temperature (°C)** Free enzyme Enzymes bound to gel membrane surface Enzymes immobilised inside beads Using the graph and your own knowledge of enzymes, answer the following questions. Describe and explain the results for the free enzyme at temperatures above 40°C. (i)

[4]

(ii) 	Explain why a higher yield of juice was obtained when using free enzyme between temperatures of 20°C and 40°C than when using immobilised enzyme. [2]	
(iii)	Suggest a reason for the differences seen in the results for the enzymes bound to the gel membrane surface with those immobilised inside the beads, between temperatures of 20°C and 60°C. [2]	



Answer Any diag	one of rams	f the following questions. included in your answers must be fully annotated.
Either,	(a)	Using examples, describe the functions of carbohydrates and lipids in living organisms. [10
Or	(b)	Describe the structure and function of the rough endoplasmic reticulum, Golg body and lysosomes. [10



Ε	Examiner
	only



E	Examiner only
	Only
END OF PAPER	10
	-



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examine only



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examiner only



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examiner only
		1
]



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examiner only



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examine only



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examiner only
		_

