

# **GCE AS MARKING SCHEME**

**SUMMER 2016** 

BIOLOGY - NEW AS UNIT 1 2400U10-1

#### INTRODUCTION

This marking scheme was used by WJEC for the Summer 2016 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

# WJEC NEW AS BIOLOGY UNIT 1 – BASIC BIOCHEMISTRY AND CELL ORGANISATION

#### MARK SCHEME

#### **GENERAL INSTRUCTIONS**

#### Recording of marks

Examiners must mark in red ink.

One tick must equate to one mark (apart from the questions where a level of response mark scheme is applied).

Question totals should be written in the box at the end of the question.

Question totals should be entered onto the grid on the front cover and these should be added to give the script total for each candidate.

# Marking rules

All work should be seen to have been marked.

Marking schemes will indicate when explicit working is deemed to be a necessary part of a correct answer.

Crossed out responses not replaced should be marked.

Credit will be given for correct and relevant alternative responses which are not recorded in the mark scheme.

#### **Extended response question**

A level of response mark scheme is used. Before applying the mark scheme please read through the whole answer from start to finish. Firstly, decide which level descriptor matches best with the candidate's response: remember that you should be considering the overall quality of the response. Then decide which mark to award within the level. Award the higher mark in the level if there is a good match with both the content statements and the communication statement. Award the middle mark in the level if most of the content statements are given and the communication statement is partially met. Award the lower mark if only the content statements are matched.

# Marking abbreviations

The following may be used in marking schemes or in the marking of scripts to indicate reasons for the marks awarded.

cao = correct answer only
ecf = error carried forward
bod = benefit of doubt

	0	-4:	Moulting dataile			Marks av	ailable		
	Que	stion	Marking details	AO1	AO2	AO3	Total	Maths	Prac
1	(a)	(i)	Telophase (1) Accept cytokinesis						
			Metaphase (1)	2			2		
			Any reference to meiosis – penalise 1 mark						
		(ii)	Chromosomes						
			Accept chromatids		1		1		
			Reject chromatin/ spindle fibres						
		(iii)	DNA is found( in other parts of the cell)(1)						
			in mitochondria (1)		2		2		
			Reject chloroplast						
	(b)		(4 days = 96 hours)						
			96/19 = 5 cycles (accept 5.05 cycles)						
			Evidence of 5 cycles e.g. 96/19 or 5 cycles/ 5.05 cycles/ 5.1						
			cycles = 1 mark						
					3		3	3	
			30 000 x 2 <sup>5</sup> (accept 30 000 x 2 <sup>5.05/5.1</sup> )						
			= 960 000/ 996 000/ 1 028 902 = 2 marks						
			= $9.6 \times 10^5 \text{ cm}^{-3}$ (accept $9.96 \times 10^5 \text{ cm}^{-3} / 1.0 \times 10^6$ ) = 3 marks						
			Question 1 total	2	6	0	8	3	0

	0	-4!	B4 a a la	in a dotaile			Marks A	Available		
	Que	stion	IMARK	ing details	AO1	AO2	AO3	Total	Maths	Prac
2	(a)		genetic material in the epithel	ial cell is DNA (1)						
			Any 3 ( x1) from:				4			
			DNA/ epithelial cell	RNA/ virus						
			double stranded/double helix	single stranded (1)						
			deoxyribose	ribose (1)						
			contains thymine	contains uracil (1)	4			4		
			Larger/ longer molecule	Smaller/ shorter molecule (1)						
			(Genetic material contained in	Genetic material not {contained in						
			nucleus)	nucleus/ membrane bound} (1)						
			Answers must be comparative	e (apart from fifth MP)						
			Reference to prokaryotic cells	is neutral						
	(b)	(i)	{Endocytosis/phagocyctosis}	Accept micropinocytosis	1			1		
		(ii)	pairs (1) Reject DNA 3. Adenine with uracil <u>and</u> cy Accept A:U, C:G 4. RNA polymerase catalyse molecule (1)	ach to their) complementary base		4		4		

Overtion	Mouking details			Marks A	vailable		
Question	Marking details	AO1	AO2	AO3	Total	Maths	Prac
(iii)	<ol> <li>Any five (x1) from:         <ol> <li>Translation (1)</li> <li>mRNA attaches to a ribosome (1)</li> <li>tRNA brings a (specific) amino acid (1)</li> </ol> </li> <li>Complementary base pairing between mRNA codon and tRNA anticodon/ codon anticodon interaction (or description of) (1)</li> <li>Ribosome has room for two tRNA molecules/Brings two amino acids into close proximity/ or description of (1)</li> <li>Condensation reaction occurs/peptide bond formed (between amino acids)(1)</li> <li>Ribosome moves to the next codon (or description of)/ reference to stop codon (1)</li> <li>Post translation processing/ or description of e.g. folding of polypeptide chain role of Golgi(1)</li> </ol>	5			5		
(iv)	It does not possess {any ribosomes/tRNA / the organelles for protein synthesis} (1) Reject: organelles unqualified All other organelles = neutral It does not possess mitochondria / cannot produce its own ATP for the process (1)		2		2		
	Question 2 total	10	6	0	16	0	0

	O ti a				Marks A	vailable		
	Question	Marking details	AO1	AO2	AO3	Total	Maths	Prac
3	(a)	A: nucleus (1) Accept nucleolus B: chloroplast (1)	1	1		2		
	(b)	apparent length = $13\text{mm} / 13 \times 1000 = 13000 \mu \text{m}$ magnification = $13000 / 32.3  \mu \text{m}$ or $13/0.0323  (1)$ magnification = $402.48 / 402.5 / 403 / 402$ (correct answer = $2  \text{marks}$ ) OR apparent length = $13.5  \text{mm} / 13.5 \times 1000 = 13500  \mu \text{m}$ magnification = $13500 / 32.3  \mu \text{m}$ or $13.5 / 0.0323  (1)$ magnification = $417.96 / 418$ (correct answer = $2  \text{marks}$ )  Award 1 mark for evidence of figures showing image size/ actual size		2		2	2	
	(c)	<ol> <li>straight chains of β-glucose/ alternative molecules of β-glucose rotate through 180° / chains cross-linked / form microfibrils (1)</li> <li>which provide {strength/rigidity/inelasticity} to the cell wall (1) support = neutral</li> <li>When the {solute concentration/ solute potential} changes (water potential will change) causing water to {move into /out of} the cell (1)</li> <li>cell wall prevents {osmotic lysis/ cell bursting}/ cell wall prevents cell shrinking (1) turgid/ plasmolysed = neutral Reject if direction of water movement is incorrect</li> </ol>	1		1	4		

Overtion	Mandain	ar dataila			Marks A	Available		
Question	Warkir	ng details	AO1	AO2	AO3	Total	Maths	Prac
3 (d)	Spirogyra – eukaryotic cells & A Similarity: Both contain riboson genetic material (1)  Difference: any 2 from: (1 ma  Spirogyra  {Membrane bound organelles/ named organelle} present  DNA enclosed within nuclear membrane  Linear DNA  Larger/80s ribosomes  Accept ribosomes are different sizes  Accept following  DNA associated with histones	nes / cell membranes / DNA/		1 1		3		
	Mesosome absent	Mesosome present						
	Question 3 total		3	6	2	11	2	0

	0		Moulsing dataile			Marks a	vailable		
	Que	stion	Marking details	AO1	AO2	AO3	Total	Maths	Prac
4	(a)	(i)	Diagram shows the addition of water (1) Glucose and fructose drawn with –OH groups on C1 and C2 in down positions (1)	2			2		
		(ii)	Glycosidic	1			1		
		(iii)	They have the same {chemical formula/ molecular formula/ number of atoms of each element} but different {structural formulae/ structures}/ both $C_6H_{12}O_6$ but different {structural formulae/ structures}	1			1		
		(iv)	Add Biuret solution to a sample of both solutions (1) Reject reference to heat The sucrase will cause a colour change from blue to lilac (and the sucrose will remain blue) (1) OR Heat both solutions with acid, then neutralise with alkali, then (heat) with Benedicts (1) The sucrose will cause a colour change from blue to brick red (and the sucrase will remain blue)(1)	1	1		2		2

0	-4! - ·-	Maulina dataila			Marks a	vailable		
Ques	stion	Marking details	AO1	AO2	AO3	Total	Maths	Prac
(b)	(i)	Any two (x 1) from:  concentration of {sucrose / sucrase / DNS} (1)  temperature (1)  filter/ wavelength of light in the colorimeter (1)  method of mixing (1)		2		2		2
	(ii)	Linear axes with values at origin (1)  X-axis labelled pH and Y-axis labelled mean absorbance + au  (1)  Correct plots (1) tolerance ± ½ small square  Plots joined (1) no extrapolation		4		4	4	4
	(iii)	Hypothesis is incorrect (1)  The {fastest rate of reaction/ most product produced} is {pH 4/ acidic/ between pH3-5}/ the optimum pH is pH 4 (1)  Use of data e.g. mean light absorption was 0.87 au at pH 4 and only 0.30 au at pH 7/ rate is approx three times faster at pH4 than at pH 7 (1)		3		3		3

	0	-4!				Marks a	vailable		
	Que	stion	Marking details	AO1	AO2	AO3	Total	Maths	Prac
4		(iv)	3.8 (μg cm <sup>-3</sup> ) (1) Accept any figure between 3.7-3.8 3.8/2 = 1.9 (μg cm <sup>-3</sup> ) (1) OR 3.7 (μg cm <sup>-3</sup> ) (1) 3.7/2 = 1.85(μg cm <sup>-3</sup> ) (1) OR 4.2 (μg cm <sup>-3</sup> ) (1) Accept any figure between 4.2-4.3 4.2/2 = 2.1 (μg cm <sup>-3</sup> ) (1) OR			2	2	2	2
	(c)		<ul> <li>4.3 (μg cm<sup>-3</sup>) (1)</li> <li>4.3/2 = 2.15 (μg cm<sup>-3</sup>) (1)</li> <li>There is {overlap / high variability} in the repeat readings at {pH 3/4/5/ low pH} (1)</li> <li>Repeat experiment with a range of intermediate values between pH 3-5/</li> </ul>			2	2		2
			Repeat data {to obtain a more reliable mean/ to improve reliability} (1) Reject references to accuracy  Question 4 total	5	10	4	19	6	15

	Overtion	Maylsing data:la			Marks a	available		
	Question	Marking details	AO1	AO2	AO3	Total	Maths	Prac
5	(a)	<ol> <li>Oxygen crosses the membrane by (simple) diffusion (1)         NOT facilitated</li> <li>oxygen passes through the phospholipid (bilayer) (1)         Membrane = neutral</li> <li>Sodium ions pass through {channel/carrier} proteins (1)         intrinsic proteins = neutral</li> <li>by {facilitated diffusion/active transport/ cotransport} (1)         Reject active transport in relation to channel proteins         If oxygen and sodium are not treated separately, allow one         mark max for correct linkage of diffusion with phospholipid         bilayer or {facilitated diffusion/ active transport/ cotransport}         with {channel/ carrier} proteins</li> </ol>	4			4		
	(b)	<ol> <li>The percentage ion composition of {fresh water/ habitat} is lower than that in the carp's blood plasma/ fresh water is hypotonic to the blood/ ORA (1)</li> <li>The water potential of fresh water is {higher/less negative} than the water potential of the carp's blood/ ORA (1)</li> <li>The carp takes in water (through the gills) by osmosis (1)</li> <li>(Producing large volumes of dilute urine) prevents {osmotic lysis/ bursting} of cells (1)</li> <li>Maintaining ion composition (1)         Accept reduces the loss of ions</li> </ol>		1	1	4		

	0	4!	Mauking dataile			Marks a	available		
	Que	estion	Marking details	AO1	AO2	AO3	Total	Maths	Prac
5	(c)	(i)	<ol> <li>Percentage ion composition of sea water is greater than that of the blood plasma/ sea water is hypertonic/ water potential in the fish is higher than the sea water/ ORA (1)</li> <li>ions will move into the blood plasma, {by (facilitated) diffusion/ down a concentration gradient/ or description of}(1)</li> <li>therefore ions must be actively transported out (of the blood plasma into the surrounding sea water) (1)</li> </ol>		1	1	3		
	(c)	(ii)	Cell 1 is from the salmon in sea water (1)  Any one from: as it is {highly folded to give a larger surface} / contains {many/more} mitochondria to {provide ATP/ for respiration/ for active transport } (1)			1	2		

	Overation				Marks a	available		
	Question	Marking details	AO1	AO2	AO3	Total	Maths	Prac
5	(d)	<ol> <li>As the percentage ion composition of the blood plasma is higher than the surrounding water (1)</li> <li>lons will move out of the {fish/gills/blood plasma} by {(facilitated) diffusion/down a concentration gradient} (1)</li> <li>Acidification will {denature/ inactivate/ change the shape of} the (carrier/ channel) proteins in the specialised cells/ denature/ inactivate enzymes (involved in ATP production) (1)</li> <li>Therefore {no/fewer} ions are (actively) transported into the {fish/gills/blood (plasma)}/ fewer ions can diffuse out (1) (in context of damaged proteins)</li> </ol>		1	1 1	4		
		Question 5 total	4	5	8	17	0	0

Ougation	Maultine details	Marks available							
Question	Marking details	AO1	AO2	AO3	Total	Maths	Prac		
6	Indicative content <ul> <li>Pyrophosphatase has tertiary structure</li> <li>The active site has a specific shape</li> <li>Pyrophosphate has a complementary shape</li> <li>fits into the active site/ forms Enzyme Substrate Complex</li> <li>lock and key theory/ induced fit</li> </ul> <li>Phenylalanine must be a non-competitive inhibitor;         <ul> <li>As it has a shape different to pyrophosphate;</li> <li>Phenylalanine bonds to the allosteric site/site away from the active site;</li> <li>Causes a change in the shape of the active site;</li> <li>Pyrophosphate's active site is no longer complementary/prevents pyrophosphate binding</li> <li>fewer enzyme-substrate complexes are formed;</li> <li>The maximum rate of reaction cannot be reached at any concentration of pyrophosphate;</li> </ul> </li>								
	<ul> <li>Phosphate must be a competitive inhibitor;</li> <li>As it has a similar shape to {pyrophosphate/ substrate} / phosphate has a complementary shape the active site;</li> <li>Phosphate {binds to/ competes for} the active site;</li> <li>Prevents pyrophosphate binding / fewer enzymesubstrate complexes are formed;</li> <li>The maximum rate of reaction can be reached at higher concentrations of pyrophosphate;</li> </ul>								

7-9 marks Indicative content of this level is: Detailed explanation of lock and key/ induced fit theory. Detailed explanation of the action of inhibition of phenylalanine. Detailed explanation of the action of inhibition of phosphate.  The candidate constructs an articulate, integrated account, correctly linking relevant points, such as those in the indicative content, which shows sequential reasoning. The answer fully addresses the question with no irrelevant inclusions or significant omissions. The candidate uses scientific conventions and vocabulary appropriately and accurately.			
4-6 marks Indicative content of this level: Any two from:  Explanation of lock and key/ induced fit theory/ or description of.  explanation of the action of inhibition of phenylalanine  explanation of the action of inhibition of phosphate.			
The candidate constructs an account correctly linking some relevant points, such as those in the indicative content, showing some reasoning. The answer addresses the question with some omissions. The candidate usually uses scientific conventions and vocabulary appropriately and accurately.			

Indicative content of this level is: Brief explanation of lock and key/ induced fit theory  OR Brief explanation of the action of inhibition of phenylalanine  OR Brief explanation of the action of inhibition of phosphate.  The candidate makes some relevant points, such as those in the indicative content, showing limited reasoning. The answer addresses the question with significant omissions. The candidate has limited use of scientific conventions and vocabulary.					
<b>0 marks</b> The candidate does not make any attempt or give a relevant answer worthy of credit.					
Question 6 total	4	3	2	9	

# Unit 1: BASIC BIOCHEMISTRY AND CELL ORGANISATION

# SUMMARY OF MARKS ALLOCATED TO ASSESSMENT OBJECTIVES

Question	AO1	AO2	AO3	TOTAL MARK	MATHS	PRAC	
1	2	6	0	8 3		0	
2	10	6	0	16	16 0		
3	3	6	2	11 2		0	
4	5	10	4	19	19 6		
5	4	5	8	17	0	0	
6	4	3	2	9 0		0	
Total	31	36	16	80 11		15	

WJEC GCE AS Biology Unit 1 (New) MS Summer 2016