

Candidate Name	Centre Number				Candidate Number			
					0			

**A LEVEL BIOLOGY****UNIT 4****Variation, Inheritance and Options****SPECIMEN PAPER****(2 hours)****90 marks****ADDITIONAL MATERIALS**

In addition to this examination paper, you will require a calculator and a ruler.

**INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen. Do not use gel pen. Do not use correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet.

For Examiner's use only			
	Question	Maximum Mark	Mark Awarded
Section A	1.	12	
	2.	12	
	3.	14	
	4.	15	
	5.	8	
	6.	9	
Section B	Option	20	
<b>Total</b>		<b>90</b>	

**INFORMATION FOR CANDIDATES**

This paper is in 2 sections, **A** and **B**.


Section **A**: 70 marks. Answer **all** questions. You are advised to spend about 1 hour 30 minutes on this section.

Section **B**: 20 marks; Options. Answer **one option only**. You are advised to spend about 20 minutes on this section.

The number of marks is given in brackets at the end of each question or part-question. The assessment of the quality of extended response (QER) will take place in question 6.

1. Spermatogenesis in humans is controlled by a number of hormones. Attempts to develop a male contraceptive pill have focussed on disrupting the sequence of processes involved in the release of the male hormones.

(a) The photomicrograph below shows a section through the seminiferous tubule of a mammal.




Use the letters on the diagram to identify the following.

[2]

- I. a cell that secretes testosterone .....
- II. a gamete .....
- III. a cell that will divide by mitosis .....
- IV. a cell that must differentiate to become functional .....

- (b) The flow-chart below shows how sperm production in humans is controlled by hormones. Follicle stimulating hormone (FSH) stimulates Sertoli cells to initiate spermatogenesis and luteinising hormone (LH) stimulates the production of testosterone, which is involved in controlling the formation of spermatids and spermatozoa.



Metronidazole is an antibacterial drug that has been shown to have negative effects on spermatogenesis. A study in rats on the effect of the drug on hormone levels produced the following results.

mass of drug administered mg / kg	mean plasma concentrations /arbitrary units		
	FSH	LH	testosterone
0	12.07	9.87	6.12
200	7.81	6.93	3.51
400	6.32	5.43	2.62

- (i) What was the mean percentage decrease in FSH plasma concentration when 200 mg/kg of drug was administered? [1]

.....%

- (ii) Calculate the mass of drug that would have to be administered to a rat with a mass of 550g in order to achieve a relative dose of 200 mg/kg. [1]

.....

The scientists carrying out the study concluded that the metronidazole was most likely to have been acting on a part of the brain.

- (iii) Use the information given to explain why the scientists may have reached this conclusion. [4]


.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

- (iv) Based on the evidence given, suggest two *other* sites where the metronidazole may have been acting, giving reasons for your answers. [4]

.....  
.....  
.....  
.....

2. *Arum maculatum* is a spring-flowering plant found in Welsh woodland. The following observations were made on the structure of its flower through dissection.

The flowers are enclosed in a green, leaf-like structure called a spadix, with a ring of female flowers at the bottom and a ring of male flowers above them. A ring of hairs above the male flowers prevents insects that enter the spadix from leaving until fertilisation has occurred. The male flowers then ripen and the hairs wither. The spadix secretes a faecal odour and the temperature within the spadix can be up to 15°C warmer than the temperature of the surroundings. Once pollinated, the spadix dies, exposing the developing fruits, that turn a bright red colour as they ripen.



- (a) Describe the sequence of events that take place during double fertilisation. [4]

.....

.....

.....

.....

- (b) What conclusions can you make regarding the types of pollination used by *Arum maculatum* to ensure fertilisation of its ovules? [5]

.....

.....

.....



.....

.....

.....

.....

- (c) The graphs below show the growing season of *Arum maculatum* and the relative light intensity reaching the ground in the woodlands where this plant grows. The table gives the mean monthly temperature in these woodlands.



Month	J	F	M	A	M	J	J	A	S	O	N	D
Mean temperature (°C)	5	4	8	10	12	13	15	16	10	8	5	4


Using your knowledge of photosynthesis explain how limiting factors affect the growth of *Arum maculatum* in April, July and November. [3]

.....

.....

.....

3. Dietary ethanol can be broken down by cells in the human body into ethanal which can then be converted into a two carbon molecule which is fed into the Krebs Cycle. Some people have mutations in the enzyme that catalyses this reaction and they are unable to metabolise ethanol.
- (a) Ethanol dehydrogenase is found in the cytoplasm of cells, particularly in the liver. The structure of the molecule is shown below.



A mutation in the DNA base sequence of the gene for ethanol dehydrogenase caused a change in the primary structure of the protein at point **X** on the diagram. This reduces the ability of liver cells to breakdown toxic ethanol.

- (i) What is meant by a gene mutation? [1]

.....  
 .....

- (ii) Use your knowledge of protein synthesis to explain how this type of mutation could lead to a change in the tertiary structure of the protein. [4]

.....  
 .....

- (iii) During the conversion of ethanol into ethanal hydrogen atoms are removed. Name a molecule that could accept these hydrogens. [1]

.....

- (iv) Suggest how this conversion could increase ATP production. [1]

.....  
 .....  
 .....

- (b) The All Wales Medical Genetics Service (AWMGS) provides specialist genetic services to individual patients and families with, or concerned about, rare genetic conditions.

DNA was extracted from two patients suffering from a defect in their ability to metabolise ethanol. Enzymes were used to cut samples of their DNA and the fragments were then separated using gel electrophoresis. The table below shows some of the enzymes available to cut DNA in this analytical technique. The arrows indicate where the enzymes cut the DNA.

Enzyme	Recognition Site
AluI	AG↓CT
BamHI	G↓GATCC
EcoRI	G↓AATTC
HaeIII	GG↓CC
HindIII	A↓AGCTT
NotI	GC↓GGCCGC
PstI	CTGCA↓G
TaqI	T↓CGA

- (i) Name the type of enzyme used to cut the DNA. [1]

.....

A part of the DNA base sequence for the ethanol dehydrogenase gene of the two patients is shown below.


<b>Patient A</b>				
TTTGTGAACC	AACACCTGTG	CGGCTCACAC	CTGGTGAAG	CTCTCTACCT 50
AGTGTGCGGG	GAACGAGGCT	TCTTCTACAC	ACCCAAGACC	CGCCGGGAGG 100
AACAATGCTG	GGCATTGTGG	TACCAGCATC	TGCTCCCTCT	ACCAGCTGGA 150
<b>Patient B</b>				
TTTGTGAACC	AACACCTGTG	CGGCTCCCAC	CTGGTGAAG	CTCTCTACCT 50
AGTGTGCGGG	GAACGAGGCT	TCTTCTACAC	ACCCAAGACC	GGTATCGAGG 100
AACAATGCTG	GGCCTTGTGG	TACCAGCATC	TGCTCCCTCT	ACCAGCTGGA 150

- (ii) From the information provided, conclude which enzymes were used to cut the DNA samples. [2]

.....

.....

Gel electrophoresis of the fragments obtained for Patient A resulted in the bands shown below.



(iii) Draw lines on the graph to show the position of the fragments obtained for Patient B. [2]

(iv) Explain why the DNA fragments can be separated in this way. [2]

.....

.....

.....

.....

4. The common primrose has flowers that vary in the position of their anthers and the length of their styles. These characteristics are controlled by single genes as shown below:

Low anther position	A	Long style	T
High anther position	a	Short style	t

Plants, pure breeding for long style and low anther position, were crossed with plants that were homozygous recessive for both characteristics. All the F<sub>1</sub> produced flowers that had low anther positions and long styles. Some of the F<sub>1</sub> offspring were then crossed together.

- (a) Complete the diagram below to show the expected genotypes, phenotypes and phenotype ratio of the F<sub>2</sub> if inheritance of these characteristics follows standard Mendelian dihybrid inheritance. [5]

Parent phenotypes	Long style, low anther	x	Short style, high anther
Parent genotypes	.....	x	.....
Parent gametes	.....	x	.....
F <sub>1</sub> genotype	.....	x	.....
F <sub>1</sub> phenotype	Long style, low anther	x	Long style, low anther
F <sub>1</sub> gametes	.....	x	.....

F<sub>2</sub> phenotypes

.....

F<sub>2</sub> genotypes

.....

.....

F<sub>2</sub> phenotype ratio

.....

- (b) One of the F<sub>1</sub> was back-crossed with the double homozygous recessive parent. The results of this back-cross are shown below.

Low anther, long style	24
Low anther, short style	10
High anther, long style	13
High anther, short style	25

It was suggested that these characteristics are not inherited following Mendel's second law as the observed numbers of this cross did not follow the expected 1:1:1:1 ratio.

Use  $\chi^2$  (chi<sup>2</sup>) to test if the anther position and the style length are inherited according to a standard dihybrid inheritance pattern or not.

The null hypothesis for this test is that there is no statistical difference between the observed and the expected results.

- (i) Complete the table to calculate the  $\chi^2$  value for this set of data.

[3]

Phenotype	Observed Numbers (O)	Expected Numbers (E)			
Low anther, long style	24				
Low anther, short style	10				
High anther, long style	13				
High anther, short style	25				
	$\Sigma$				

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

$$\chi^2 = \dots\dots\dots$$

- (ii) Use your calculations and the probability table below to conclude whether to accept or reject the null hypothesis giving a reason for your answer. [4]

.....

.....

.....

.....

.....

**Probability Table for  $\chi^2$  Test**


Degrees of freedom	0.90 90%	0.80 80%	0.70 70%	0.50 50%	0.30 30%	0.20 20%	0.10 10%	0.05 5%	0.02 2%	0.01 1%
1	0.026	0.06	0.15	0.46	1.07	1.64	2.71	3.84	5.41	6.64
2	0.21	0.45	0.71	1.39	2.41	3.22	4.61	5.99	7.82	9.21
3	0.58	1.01	1.42	2.37	3.67	4.64	6.25	7.82	9.84	11.34
4	1.61	2.34	3.00	4.35	6.06	7.29	9.24	11.07	13.39	15.09

- (c) One explanation for the deviation shown between the observed and expected results was that the alleles for these characteristics are carried on the same chromosome.

With the use of annotated diagrams, explain how the F<sub>1</sub> plant could give rise to four different gametes if the alleles show linkage.

[3]

5. Pregnancy testing involves detecting changes in the concentration of the hormones progesterone or human chorionic gonadotrophin (HCG) in blood or urine. The graph shows how plasma concentration of these hormones changes during pregnancy.



- (a) (i) Identify each of the two hormones shown in the graph and explain which one would provide the best indication of pregnancy in the early stages. [3]

.....

.....

.....

(ii) Describe the role of each of these hormones in the early stages of pregnancy.

I Hormone **A**: [1]

.....

.....

.....

II Hormone **B**. [2]


.....

.....

.....

(b) Pregnancy testing kits were developed in the early 1980s to detect one of these hormones in urine.

The diagram shows the results of pregnancy testing in a sample of 1200 women at 5 and 21 days following ovulation. All women in the sample were subsequently shown to have conceived at the time of testing.



- (i) Explain why the negative results were treated as false negatives. [1]

.....  
.....

- (ii) Calculate the number of women in the study who were given a positive result at 5 days post ovulation.

[1]

Number of women = .....

8


6. The Grand Banks is an area of sea off the coast of Newfoundland in Canada. It was once one of the most productive fishing grounds in the world for Atlantic cod.

The cod was fished heavily for about 50 years. About 60% of the total cod population of reproductive age was harvested annually.

Cod fishing in the Grand Banks was closed in 1992 but by then the population was less than 1% of what it had been.

Cod grow evenly throughout their life.

The cod that remained when fishing was finally closed were much smaller and grew more slowly than the cod that lived in the Grand Bank several decades previously.



With reference to the information provided above and your knowledge of evolution, explain the changes observed in the phenotype of the Atlantic cod.

(The quality of your extended response will be assessed in this question.) [9QER]

.....

.....

.....







**SECTION B: OPTIONAL TOPICS**Option A: **Immunology and Disease**Option B: **Human Musculoskeletal Anatomy**Option C: **Neurobiology and Behaviour**

Answer the question on **one topic only**.


Place a tick (✓) in one of the boxes above, to show which topic you are answering.

**You are advised to spend about 20 minutes on this section.**

### Option A: Immunology and Disease

7. Influenza is endemic in most communities and is characterised by large scale outbreaks that can reach world-wide proportions. There are three distinct varieties, A, B and C that can be subdivided into many serotypes. Classification and identification of strains of the virus are based on immunological analysis of their surface proteins. Attempts to develop a 100% effective vaccine have failed, with most available vaccines affording, at best, about 60% protection.

The diagram shows the structure of an influenza virus.



- (a) Using the information above answer the following questions.
- (i) Define the term *endemic*. [1]
- .....
- (ii) What is the term used to describe '*...large scale outbreaks that ...reach world-wide proportions*'? [1]
- .....
- (b) Research into vaccine development has focussed largely on the neuraminidase and haemagglutinin proteins.
- (i) Describe how you could test for the presence of protein in a sample of the virus. [2]
- .....
- .....

- (ii) Explain why research into influenza vaccines has focussed on these proteins. [2]

.....

.....

.....

.....

.....

- (iii) It has been found that the haemagglutinin protein stimulates a strong immune response but shows a high degree of antigenic variation over time while the neuraminidase protein is less immunologically active but shows far less variation between strains and over time.

Evaluate the choice of neuraminidase as the better option for research into developing a vaccine that would provide effective, long-term immunity against the influenza virus. [3]

.....


.....

.....

.....

.....

- (c) The graph shows how blood antibody concentration against *Rubella* changed in a 12 month old child following injection with *Rubella* antigen at day 0 followed by a booster injection at day 40.



- (i) Following the first injection with *Rubella* antigen the concentration of antibodies against the virus remained low for several days.

State the time taken for a protective level of antibody to be produced following the first injection.

[1]

.....

- (ii) Explain why the time taken to produce the antibody level required for immunity was much shorter following the booster injection.

[2]

.....  
 .....  
 .....  
 .....

(d) The child, who had not been previously exposed to measles, was given a separate immunisation against the measles virus at Day 40 - the same day that she received the booster injection against Rubella.

(i) On the same graph draw a line to show how the blood antibody concentration against measles would change between days 40 and 60. [2]

(ii) Explain why the blood antibody concentration against measles would change in this way. [3]


.....

.....

.....

.....


(e) The photograph shows the result of an ELISA (enzyme-linked immunosorbent assay) used to determine the concentration of antibodies in the plasma. The process is summarised below.



1. Rubella antigen is immobilised on an inert plastic matrix
2. the test sample is diluted and  $0.4\text{cm}^3$  of the diluted plasma is placed in a separate well of the plate
3. anti-human antibodies which have an enzyme attached are then added – when a suitable substrate is added a colour change is observed
4. absorbance of light at  $492\text{nm}$  is then measured
5. the concentration of antibody in the diluted sample is determined from a standard curve of absorbance at  $492\text{nm}$  for known antibody concentrations

A  $0.4\text{cm}^3$  sample of plasma from a person infected with Rubella was diluted by a factor of  $10^{-3}$  from its original concentration. It was found to have a mean absorbance of 0.24 at 492nm.

Using the standard curve of the absorbance of light at 492nm of known antibody concentrations shown below, calculate the mass of antibody present in  $1\text{cm}^3$  of the original sample of plasma. Give your answer in  $\text{mg cm}^{-3}$ . [3]




Antibody concentration: ..... $\text{mg cm}^{-3}$  of plasma.

20

**Option B: Human Musculoskeletal Anatomy**

8. The photomicrograph shows a section of compact bone.



Complete the table below by giving the function of the two types of cell which could be found in the lacuna.

[1]

Name of cell	Function
Osteoblast	
Osteoclast	

- (ii) Name **two** structures *other* than nerve fibres which are found in the Haversian canal.

[1]

.....

.....

- (iii)
- (iii) The photographs below show human vertebrae. Identify which image shows a cervical vertebra and use the photograph to describe how its structure is adapted to its function in the vertebral column. [2]

**A**



**B**



**C**



**Cervical vertebra** .....

Adaptations to function .....

.....

.....

.....

- (b) Health workers in Wales have observed a rise in the number of cases of rickets in some parts of the country. It has been suggested that a reduction in outdoor play and the consumption of fewer dairy products could be contributing towards this.  
Explain how these lifestyle changes could result in the development of rickets in young children. [5]

.....

.....

.....

.....


.....

.....

.....

.....

(c) The electron micrograph shows a section of muscle.



(i) Name the molecules labelled **A** and **B** in the micrograph above. [2]

**A** .....

**B** .....

(ii) Identify the type of section shown in the micrographs above. [1]

.....

(iii) Micrograph 2 above shows a section taken through the outside of the A band of the sarcomeres.

Draw a labelled diagram below to represent a similar section taken through the I band of the sarcomere. [1]

- (d) The length of the A band in human muscle is 1.85  $\mu\text{m}$ .  
0.25  $\mu\text{m}$  at the centre of the myosin filament does not have myosin heads.  
There are 6 myosin heads in 40 nm of myosin filament.

Calculate how many myosin heads there are in each myosin filament.  
Show your working.

[3]

Number of myosin heads = .....

- (e) Doctors examined a patient suffering from weak, uncoordinated muscular contraction. Tests showed that the patient had lower than normal blood calcium levels. Doctors concluded that this was a possible cause of the symptoms observed.

Explain why the low calcium levels could have resulted in the symptoms observed.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

### Option C: Neurobiology and Behaviour

9. Below is a diagram of the left hemisphere of the brain showing four areas that are involved in speech and language.



- (a) (i) Identify areas **A**, **B**, **C** and **D** in the diagram: [3]

**A** .....

**B** .....

**C** .....

**D** .....

- (ii) Describe the role played by Wernicke's area and Broca's area in speech and language. [2]

.....

.....

.....

- (b) Functional Magnetic Resonance Imaging (fMRI) is a technique for examining activity of the brain. During investigations, the subjects were given a book to read while their brain activity was monitored. The images produced were then examined and the results shown on the table below.

Individual	Brightest areas of brain			
	A	B	C	D
1	✓	✓	✓	✓
2			✓	✓

The scientists concluded that individual 2 was reading silently. What conclusions should be reached regarding the activities of individual 1? Explain your answer.

[1]


.....

.....

.....

.....

- (c) The control of ventilation is brought about by the autonomic nervous system and is shown in the diagram below. Within the medulla are the inspiratory and expiratory centres, which send nerve impulses to the intercostal muscles and diaphragm. When the expiratory centre is stimulated it 'switches off' the inspiratory centre.



- (i) Use the diagram and your knowledge of ventilation to describe how inspiration is initiated. [2]

.....

.....

.....

- (ii) Use the diagram and your knowledge to explain the mechanism by which the ventilation rate is increased during exercise. [4]

.....


.....

.....

.....

.....

- (d) A student set up an experiment to investigate the effectiveness of different types of insect repellent. The diagram below shows the apparatus that she used.



The insecticide was sprayed into the branch of the glass tube labelled E and the mosquitoes were introduced to the bottom of the tube. The number of mosquitoes that entered the branch of the tube labelled F in two minutes. The student repeated the experiment and the apparatus was rinsed with water after each trial. The same volume and concentration of insecticide was sprayed in each trial. The data obtained is shown below.

Repellent	Number of mosquitoes entering branch F						
	Trial 1	Trial 2	Trial 3	Trial 4	Trial 5	mean	Standard deviation
Control	10	9	8	12	11	10.0	1.41
Brand A	11	14	12	13	11	12.0	1.17
Brand B	16	17	19	18	17	17.4	1.02
Brand C	19	16	13	10	9	13.4	

- (i) Suggest how the apparatus was set up for the control experiment. [1]

.....

- (ii) What type of behaviour is being exhibited by the mosquitoes in response to the repellent? [1]

.....

- (iii) Calculate the standard deviation for brand **C** by completing the table below and using the formula given. [3]

$x$	$x - \bar{x}$	$(x - \bar{x})^2$
19	5.6	31.36
16	2.6	6.76
13	-0.4	0.16
10	-3.4	11.56
9	-4.4	19.36
		$\Sigma =$

The formula for standard deviation is:

$$\sqrt{\frac{\Sigma (x - \bar{x})^2}{N}}$$

- Where  $x$  = individual trial results for Brand **C**  
 $\bar{x}$  = mean results for Brand **C**  
 $N$  = number of trials for Brand **C**  
 $\Sigma$  = sum of

Standard deviation for Brand **C** = .....

- (iv) What conclusion could be reached regarding the effectiveness of the different types of insect repellent? Comment on the strength of evidence for this conclusion. [3]

.....  
 .....  
 .....