


Q1. Cow's milk contains the sugar lactose. Many cats are unable to digest cow's milk because they are lactose intolerant.

Cow's milk can be made suitable for these cats by treating it with the enzyme lactase to hydrolyse lactose. This makes the cow's milk lactose-free. Beads are coated with lactase and placed in a tube, as shown in the diagram below. Cow's milk flows over the beads and the lactose is hydrolysed.



(a) Attaching lactase to the beads is a more efficient use of lactase than adding the lactase directly to cow's milk.

Suggest **three** reasons why it is more efficient to attach lactase to the beads.

- 1
-
- 2
-
- 3
-

(3)

(b) Monosaccharides and disaccharides taste sweet. The lactose-free milk made after hydrolysis with lactase tastes sweeter than the cow's milk containing lactose. Suggest why.

.....
.....
.....
.....
.....

(2)
(Total 5 marks)

Q2.(a) (i) Give **two** ways in which the structure of starch is **similar** to cellulose.

1

2

(2)

(ii) Give **two** ways in which the structure of starch is **different** from cellulose.


1

2

(2)

(b) In plants, mass transport of sugars takes place through columns of sieve cells in the phloem. Other cells, called companion cells, transport sugars into, and out of, the sieve cells.

The diagram shows the structure of phloem.



Structures **I** and **J** allow the transport of sugars between cells.

- (i) Using the diagram, suggest and explain **one** other way in which sieve cells are adapted for mass transport.

.....

.....

.....

.....

.....

(2)

- (ii) Using the diagram, suggest and explain **one** other way in which companion cells are adapted for the transport of sugars between cells.

.....

.....

.....


.....

.....

(2)
(Total 8 marks)

Q3. Starch and cellulose are two important plant polysaccharides.

The following diagram shows part of a starch molecule and part of a cellulose molecule.



(a) Explain the difference in the structure of the starch molecule and the cellulose molecule shown in the diagram above.

.....

(2)

(b) Starch molecules and cellulose molecules have different functions in plant cells. Each molecule is adapted for its function.

Explain **one** way in which starch molecules are adapted for their function in plant cells.

.....

(2)

(c) Explain how cellulose molecules are adapted for their function in plant cells.

.....
.....
.....
.....
.....
.....

(Extra space)

.....
.....

(3)
(Total 7 marks)

Q4.A student investigated the effect of chewing on the digestion of starch in cooked wheat.

He devised a laboratory model of starch digestion in the human gut. This is the method he used.

1. Volunteers chewed cooked wheat for a set time. The wheat had been cooked in boiling water.
2. This chewed wheat was mixed with water, hydrochloric acid and a protein-digesting enzyme and left at 37 °C for 30 minutes.
3. A buffer was then added to bring the pH to 6.0 and pancreatic amylase was added. This mixture was then left at 37 °C for 120 minutes.
4. Samples of the mixture were removed at 0, 10, 20, 40, 60 and 120 minutes, and the concentration of reducing sugar in each sample was measured.
5. Control experiments were carried out using cooked wheat that had been chopped up in a blender, not chewed.

(a) What reducing sugar, or sugars, would you expect to be produced during chewing? Give a reason for your answer.

.....
.....
.....
.....

(2)

- (b) In this model of digestion in the human gut, what other enzyme is required for the complete digestion of starch?

.....

(1)

- (c) What was the purpose of step 2, in which samples were mixed with water, hydrochloric acid and pepsin?

.....

.....

(1)

- (d) In the control experiments, cooked wheat was chopped up to copy the effect of chewing.

Suggest a more appropriate control experiment. Explain your suggestion.

.....


.....

.....

.....

(2)

- (e) The figure below shows the student's results.



Explain what these results suggest about the effect of chewing on the digestion of starch in wheat.

.....

.....

.....

.....

.....

.....

(3)
(Total 9 marks)

Q5.(a) In humans, the enzyme maltase breaks down maltose to glucose.
This takes place at normal body temperature.

Explain why maltase:

- only breaks down maltose
- allows this reaction to take place at normal body temperature.

.....

.....

.....

.....

.....

.....
.....
.....
.....
.....

(Extra space)

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

(5)

- (b) Scientists have investigated the effects of competitive and non-competitive inhibitors of the enzyme maltase.

Describe competitive and non-competitive inhibition of an enzyme.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

.....

.....

.....

(Extra space)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(5)
(Total 10 marks)

Q6. Many sports drinks contain water, sodium chloride and carbohydrates. The manufacturers of the sports drinks claim that carbohydrates provide an energy boost. The sodium chloride is used to increase absorption of glucose in the small intestine.

Scientists investigated the effect of a sports drink on the performance of runners in 5 km races.


They recruited 100 runners who had previously run a 5 km race in similar times. During this race, Race 1, they had water they could drink.

The scientists divided the runners into two equal groups, **P** and **Q**. Both groups ran a second 5 km race, Race 2. During this race:

- group **P** had water available
- group **Q** had the sports drink available.

The scientists recorded the mean time for each group to complete this race.

Figure 1 shows their results.




The glycaemic index (GI) is a measure of the increase in blood glucose concentration after eating a given mass of a food compared with eating the same mass of pure glucose. The GI of pure glucose has a value of 100.

The GI of a food depends on several factors such as how much starch and sugars it contains. High GI foods include those containing lots of simple sugars or white flour. The carbohydrates in these foods are rapidly digested and absorbed. Low GI foods include wholegrain bread and breakfast cereals that contain a lot of fibre. The carbohydrates in these foods are digested and absorbed more slowly.

Figure 2 shows changes in blood glucose concentration after eating meals of high GI food and meals of low GI food.

Figure 2



Explain how a sports drink could provide an energy boost when running.

.....

.....

.....

.....

.....

.....

(Extra space)

.....

.....

.....

.....

(3)
(Total 3 marks)