- **M1.**(a) 1. Similarity directional response (to a stimulus) / movement towards / away from a stimulus;
  - 2. Difference taxis (whole) organism moves <u>and</u> tropism a growth (response).

Must be clear which one, taxis or tropism, they are referring to

Taxis occurs in animals / motile organisms <u>and</u> tropism occurs in plants

2

(b) 1. Grow in direction of / towards (pull of) gravity;

Accept: tropism for growth Ignore: pulled by gravity

Accept: positively geotropic / gravitropic

2. Grow away from salt;

Accept: negatively chemotropic / halotropic 1 and 2. Ignore: references to bends / moves

3. Salt has more effect (than gravity).

Accept: converse statement for gravity

Note: all three points may appear in one sentence

3

- (c) 1. More carriers in (cell) L / lower in R;

  Accept: left for L and right for R / side nearer salt for L
  - Accept. Test for L and right for R7 side hearer sait for L
  - (So) less IAA in (cell) L / more IAA in (cell) R;
     Accept: more IAA moves out of L / less IAA moves out of R
  - (So) more (elongation) growth in L / less (elongation) growth in R.
     Accept: less inhibition of growth in L / more inhibition of growth in R;

[8]

3

**M2.**(a) 1. (Taxis is) movement towards / away from a stimulus / a directional response / movement (to a stimulus);

2. (Move towards) temperature they were used to / cultured in;

Movement towards temperature they were used to = 2 marks

2 max

- (b) 1. Hungry, so seeking food / in absence of food respond to temperature;

  Ignore references to temperature and enzymes

  Must be stated not inferred from other statements
  - 2. Move towards temperature they were used to / cultured in;
  - 3. Associate (this temperature) with food;

    Accept they think food is here

    Stated not inferred
  - 4. (Then) stay in this temperature;

3 max

- (c) 1. (Dim) worms live in soil / dark / affected by bright light / dim light is like normal environment / what they are used to;
  - (Even) because worms might move towards / away from bright light / to avoid creating light gradient / prevent worms showing phototaxis / all parts of surface exposed to same light;

Accept to avoid kinesis due to light

3. (Dim light) ensures heat from light not a variable / heat from lamp could kill / dry out worms;

Not just to control variables / factors

2 max

1

[7]

**M3.**(a) Push – legume

Pull – grass;

Both needed for mark

(b) 1. Set up tape measures on two sides of the plot / make grid of plot; Allow 'Number each plant'. With this approach mp3 cannot

be awarded.

- 2. Use random number table / calculator / generator; Allow 'Select from a hat' idea.
- 3. To generate coordinates;

3

- (c) 1. To prevent competition between the maize and the grass;
  - 2. For light / nutrients / water;

OR

- 3. Idea of limits movement of pest (between grass and maize);
- 4. Only eating / damaging grass;

2 max

 Nitrogen-fixing bacteria convert nitrogen (in the air) into ammonium compounds (in the soil) which are converted into nitrates / nitrification occurs;

Accept 'ammonia' for 'ammonium compounds'.

- 2. Maize uses nitrates (in soil) for amino acid / protein / ATP / nucleotide production;
  - 2. Must be in the context of maize. Ignore ionic formulae unless only these are given.

2

- (e) 1. Reduced % damage to maize plants / increased maize grain yield;
  - 2. Calculation to justify mp 1;
  - 3. Standard deviation shows no overlap but need stats to show significance of this difference:
  - 4. More profit / net income / greater income than additional cost (with push-pull);
  - 5. \$322 extra / 408% more / \$401 v \$79 profit;

Accept '\$350 extra income compared to \$28 extra spend'. Mp5 gains credit for both mp4 and 5

3 max

[11]

## **M4.**(a) Three changes described;;;

Neutral nucleus shrinks, since it doesn't

Eg

- 1. Formation / growth of vacuole;
- 2. Formation of starch grains / amyloplasts;
  - 2. Accept starch grains get bigger
- 3. Movement of grains / amyloplasts towards bottom of cell;

  Note list rule applies
- 4. Cells get longer / wider / larger;

3 max

(b) 1. Grows sideways before starch grains form;

Q

- 2. Bending starts when / as grains form;
- 3. More bending as grains increase in number;
  - 3. Ignore starch grain growth references
- 4. More elongation (of cells) / growth (of roots) downwards as starch grains increase / move;
- 5. Bending starts before grains move down;
- 6. Could be related to vacuole;
  - 6. Ignore references to nucleus

3 max

- (c) 1. (IAA) at bottom of root / where IAA concentration high inhibits expansion / elongation (of cells);
  - 2 and 3 need reference to expansion / elongation, not just growth
  - 2. (IAA) at top of root / where IAA concentration low leads to expansion / elongation (of cells);
    - 2. Accept less inhibition

2

**M5.** (a) 1. (Seedlings) respond to light / are phototropic;

Reject: roots are positively phototropic / grow towards light

OR

Neutral: 'to control a variable'

2. (Only) measuring the effect of gravity / response to gravity;

Neutral: light affects growth / results

1

(b) 1. (Cells in) root tip detect gravity / respond to gravity;Must refer to root tip and not just the root

OR

2. IAA / auxin is produced in the root tip;

1

- (c) (i) 1. IAA / auxin moves to lower side / more IAA / auxin on lower side;

  Accept: references to 'cell elongation' instead of 'growth'
  - 2. Lower side grows less / slower / upper side grows more / faster / inhibits growth on lower side;

Note: if auxin is placed at upper side, mark point 2 can still be awarded

Need idea of 'less / slower' or 'more / faster' for mark point 2

2

- (ii) 1. Less IAA / auxin (produced);
  - 2. Lower side grows more / faster / less inhibition of growth on lower side;

Must refer to the lower side

2

[6]