

1	Droplets containing the organisms that cause TB are released into the air when a person suffering from TB coughs. Transmission of TB occurs if these droplets are inhaled into the alveoli of the lungs. In the lungs, the organisms are taken up by macrophages and carried to lymph respectively.		ffering from TB coughs. Transmission of TB occurs if these droplets are	
			es.	
	(a) (i)	St	ate one characteristic symptom of TB other than coughing.	(1)
	(ii)		ace a cross ⊠ in the box next to the name of the organism that causes TB.	(1)
	\times	A	Macrobacterium tuberculosis	
	\times	В	Microbacterium tuberculosis	
	X	C	Monobacterium tuberculosis	
	\times	D	Mycobacterium tuberculosis	
	(iii)	De	escribe how the organisms that cause TB are taken up by macrophages.	(3)
	(iv)		gesting food containing these organisms is unlikely to lead to the evelopment of TB. Give an explanation for this.	(2)

*(b) The graphs below show data related to TB and HIV infections in the population of central Africa from 1990 to 2008.

Graph 1 – The percentage of the population infected by TB



Graph 2 – The number of cases of HIV infection per 100 000 population



The increase in HIV infection in central Africa has led to an increase in TB infection.	2
in 15 injection.	(4)
(Total for Question 1 = 11	marks)

Discuss how far the data in the graphs support the following hypothesis.

2	and viruses.	
	(a) Give two differences between the genetic material of bacteria and viruses.	(2)
	(b) Tuberculosis (TB) is caused when droplets, containing the bacterium <i>Mycobacterium tuberculosis</i> , are inhaled into the lungs.	
	In the lungs, large numbers of the bacterium are formed rapidly. These can be ingested by macrophages. Eventually, tubercles (tissue masses), containing dormant bacteria inside macrophages, may form.	
	(i) Describe how macrophages ingest the bacteria.	(2)
	(ii) Suggest why treatment with antibiotics may not be effective against the dormant bacteria in the tubercles.	(2)

	(iii) TB can be prevented by vaccination. Explain how a person can develop artificial active immunity following vaccination.	
		(3)
(c	c) In a person with TB, the dormant bacteria in tubercles may be activated after several years. The bacteria multiply rapidly, resulting in severe lung damage.	
	The bacteria are released from the tubercles. These bacteria can inhibit the activity of T cells and infect other organs.	
	Explain why the activity of these bacteria and the inhibition of T cells means that	
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)
	a person may quickly develop severe symptoms leading to death.	(4)

3	Bacteria and viruses can cause human diseases.				
	(a) Distinguish between the structure of bacteria and viruses	(3)			
	(b) Infection with a bacterium can result in the development of active immunity	/ to			
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.				
	that bacterium. This results in the production of antibodies by plasma cells.				
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.	sma			
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.	sma			
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.	sma			
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.	sma			
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.	sma			
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.	sma			
	that bacterium. This results in the production of antibodies by plasma cells. (i) Describe how infection with a bacterium results in the production of plasma.	sma			

(ii) Explain how antibodies help a person to recover from an infection.	(2)
(iii) A person who has had an organ transplant has to take immunosuppressive drugs. This prevents the immune system from destroying the organ transplant. Some of these drugs work by inhibiting the production of cytokines.	
Suggest what effect these drugs could have on a person infected with a bacterium or a virus.	
	(4)
(Total for Question 3 = 13 m	arks)

(a) The table below sl For each feature, p	nows some feature blace one cross ⊠	es found in bacter in the appropriate	viruses and bacteria. ria and viruses. e box, in each row, to show in both bacteria and viruses. (3)
Feature	Bacteria only	Viruses only	Both bacteria and viruses
Glycogen granules		\boxtimes	
lucleic acids	×	×	×
medications.	severe cases of BR	D will involve the	use of antibiotics and other
(b) Mild cases of BRD The treatment of s medications.	can usually be tresevere cases of BR	ated using antibion	
(b) Mild cases of BRD The treatment of s medications. (i) Suggest why	can usually be tresevere cases of BR	ated using antibion	use of antibiotics and other , are needed to treat the most
(b) Mild cases of BRD The treatment of s medications. (i) Suggest why	can usually be tresevere cases of BR	ated using antibion	use of antibiotics and other , are needed to treat the most
(b) Mild cases of BRD The treatment of s medications. (i) Suggest why	can usually be tresevere cases of BR	ated using antibion	use of antibiotics and other , are needed to treat the most

(ii) The table below shows the effectiveness of various antibiotics on three species of bacteria that can contribute towards severe cases of BRD.

Antibiotic	Effectiveness of various antibiotics on three BRD bacterial pathogens (%)				
Antibiotic	Mannheimia haemolytica	Pasteurella multocida	Histophilus somni		
Danofloxacin	71	88	84		
Enrofloxacin	83	93	95		
Florfenicol	85	90	95		
Oxytetracycline	56	70	55		
Spectinomycin	72	76	67		
Tilmicosin	61	64	93		

A group of cattle has BRD but the bacteria pathogen has not been identified. Suggest which antibiotics would be the most suitable to use to treat these cattle.

(3)

Give reasons for your answer.

treatment of these cattle, once the pathogen has be	
, ,	(3)
/-	
(Total	for Question 4 = 11 marks)