## **Animal Responses**

v T glycogen granules

**1.** The image below is a transmission electron micrograph of a section of skeletal muscle.

Which row, A to D, shows the correct labels?

	Organelle T	Region U	Region V	Region between X and Y
Α	Golgi body	I-line	actin	Z-band
В	mitochondrion	Z-line	myofibril	sarcomere
С	sarcoplasmic reticulum	A-band	collagen	I-band
D	mitochondrion	I-band	myosin	sarcoplasmic reticulum

Your answer

[1]

**2.** The drug metoprolol prevents stimulation of post-synaptic receptors in the sympathetic nervous system.

Which of the following conditions could this drug be used to treat?

- 1 Muscle fatigue
- 2 Tachycardia
- 3 High blood pressure
- **A** 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1



[1]

**3.** The image below is a diagram of the human brain.



Which of the labelled regions would be directly involved in learning to play a musical instrument?

A W and X
B W and Y
C W and Z
D Y and Z

Your answer

[1]

4. Collagen is found in tendons. Tendons attach muscles to bones.

Which of the following lists of properties, **A** to **D**, makes collagen suitable for this role?

- A strong, inflexible, insoluble
- **B** strong, flexible, soluble
- ${\bm C}$  strong, inflexible, soluble
- **D** strong, flexible, insoluble

Your answer

[1]

- 5. Which of the following statements is / are correct?
  - 1 The autonomic nervous system contains sensory and motor neurones.
  - 2 Somatic and parasympathetic motor neurones use different neurotransmitters.
  - 3 Somatic motor neurones stimulate skeletal muscles whereas autonomic motor neurons stimulate only glands.
    - A 1, 2 and 3 are correct
    - B only 1 and 2 are correct
    - C only 2 and 3 are correct
    - D only 1 is correct

Your answer

[1]

6. Multiple sclerosis (MS) is an autoimmune disease that damages the nervous system.

Guillain–Barré syndrome is another autoimmune condition in which neurones are damaged and the rate of nervous impulses is reduced.

affects the central nervous system.

Guillain-Barré syndrome affects the peripheral nervous system.

i. Suggest **two** symptoms of MS that might **not** be present in people with Guillain–Barré syndrome.

Explain your answers.

1\_\_\_\_\_\_ \_\_\_\_\_\_ 2\_\_\_\_\_\_ \_\_\_\_\_\_\_[2]

ii. Multiple sclerosis and Guillain–Barré syndrome both cause muscle weakness and loss of muscle function.

Suggest and describe how the function of neuromuscular junctions will be affected by multiple sclerosis and Guillain–Barré syndrome.

\_\_\_\_\_[2]



7. Fig. 16.1 shows a drawing of a dissected human heart.. Fig. 16.2 is a diagram of a section through the human brain.



Fig. 16.1

i. Identify the letter and name of the structure in **Fig. 16.2** that is connected by nerves to structure **A** in **Fig. 16.1**.

Letter	
Name	

[2]

Normal human resting heart rate is approximately 70 beats per minute (bpm). Cutting the parasympathetic nerve to the heart increases this to approximately 100 bpm. ii.

Suggest **two** conclusions that could be made from this observation about the control of resting heart rate in normal humans.

-	1 	
-	 2 	
-		
	[2]	
	<ul> <li>Injury to the parts of the brain labelled G and H in Fig. 16.2 can lead to a range of symptoms including:</li> <li>fatigue</li> <li>weight gain</li> <li>menstrual irregularities</li> <li>low blood pressure or dizziness</li> <li>increased sensitivity to cold.</li> </ul> Outline how injury to G and H is able to cause such a wide range of symptoms.	
 iv.	Suggest why it can be difficult for a doctor to conclude that the symptoms described in part (iii) are definitely caused by damage to parts of the brain.	<u>2]</u>
		[1]

**8. Fig. 20.1** is a flow diagram that shows the sequence of events in the body once a threat is perceived. The response is often described as the 'fight or flight' response as it prepares the body to respond physically to the threat in the short-term.



i. Identify two signalling molecules named in Fig. 20.1.

1	1	_	_	 	 	_	_	-	_	_	_	-	 	 	 _	_	_	 	_	_	-	 _	 	 _	_	-	 _	_	_	 	_	_	 	 	_	_	_	_	 	 	_	_	_	 	 _	 	 -
2	2	_	_	 	 	_	_	_	_	_	_	_	 	 	 _	_	_	 		_		 _	 	 _	_		 _	_	_	 			 	 	_		_		 	 	_	_	_	 	 	 	 _

[1]

ii. Adrenaline acts on a variety of cell types with a variety of responses.

Complete the table by stating the effects of stimulating each target cell. The first one has been completed for you.

Target cell	Response	Role in the 'fight or flight' response
Smooth muscle in bronchioles	Muscle relaxes	Bronchioles dilate and allow more oxygen to reach blood
Sino-atrial node		
Liver cell		
Erector muscle in skin		

iii. Describe the sequence of actions that occur once adenylyl cyclase is activated in the target liver cells.

	[2]
iv.	The response in <b>Fig. 20.1</b> also occurs when a person is subjected to stress. However, the body does not need to respond physically to the stimulus and so, for example, the bronchioles do not dilate.
	From the information given and your own knowledge, suggest the long term adverse effects of continued exposure to stress on body function.
	[2].

9. Reflex actions are rapid responses that protect the body from harm.

The Moro reflex is found in babies up to five months of age, and occurs when the baby feels its head is suddenly no longer supported. The Moro reflex is made up of the following responses:

- The baby spreads out its arms then brings them together rapidly.
- The baby cries.

i. Suggest how the Moro reflex helps to prevent harm to a newborn baby.

\_\_\_\_\_

ii. The Moro reflex gradually disappears and usually stops completely after babies reach nine months. Other reflexes develop as children grow older.

Describe a reflex response a 3-year-old child would make to an object moving towards their eyes **and** explain the advantage of this response.

[3]

**10(a).** Squid are an order of aquatic animals. Many species of squid show a reflex response to changes in their body position.

- •A sensory receptor called a statocyst detects changes in the body position of the squid in their aquatic environment.
- •The reflex response corrects the squid's body position.

i. Describe the components and events involved in a reflex arc.

\_\_\_\_\_

ii. Suggest one benefit to the squid of the reflex response described above.

\_\_\_\_\_[1]

iii. A statocyst sensory receptor is shown in Fig. 2.1.



What can you conclude from Fig. 2.1 about the mechanism by which a statocyst acts as a transducer?

[3]

(b). Lobsters and sea cucumbers also have statocysts. From this information, the student concluded:

'I think this means squids, lobsters and sea cucumbers evolved from a recent common ancestor and are probably classified in the same phylum.'

Evaluate the extent to which the information about lobsters and sea cucumbers supports the student's conclusion.

\_\_\_\_\_

**11.** Complete the following passage using the most appropriate terms.

The knee-jerk reflex is an example of a spinal reflex as it only goes through the ...... whereas the blinking reflex is a cranial reflex as the pathway goes through the brain.

Reflexes are very fast as they only involve one or two ....., which are the slowest parts of nerve transmission. This is one reason why reflexes give a survival advantage.

[2]

12. \* Water reabsorption in the kidney is controlled by the endocrine and nervous systems.

Aldosterone and ADH are hormones that act on the kidney.

Aldosterone causes sodium ions to be pumped from the collecting duct cells into tissue fluid.

Describe how the endocrine **and** nervous systems work together to increase water reabsorption from the collecting duct.

[6]

**13.** Acetylcholine (ACh) is a neurotransmitter in mammals. Studies have suggested that it also functions as a hormone in some invertebrate species, such as squid.

When ACh comes into contact with specialised cells in squid skin, it causes them to change colour. These colour changes allow the squid to communicate and to camouflage itself.

ACh is made by cells in the centre of the squid's body.

Explain how it is possible for ACh to have an effect on cells in the skin of the squid.

 [2]

14. Fig. 16.1 shows a drawing of a dissected human heart.



Fig. 16.1

i. Identify the structures labelled A to E on Fig. 16.1.

A	
В	
С	
D	
E	

[5]

ii. State which subdivision of the peripheral nervous system supplies structure **A** on **Fig. 16.1**.

[1]

15. A student investigated the heart rates of smokers and non-smokers.

- Each test subject had their resting heart rate measured using an electronic heart rate monitor.
- They ran 1 km on a running track and their heart rate after running 500 m was recorded.
- Their heart rate was recorded for a third time 3 minutes after the completion of the exercise.

All test subjects were 18 years old. Subjects were tested between 9 am and 4 pm on one day, one at a time. Each test lasted approximately 20 minutes in total. The tests were repeated one week later using the same method. Mean heart rates were calculated for each subject.

The student's plan was to compare the heart rates of smokers and non-smokers using Student's *t*-test.

The student's results are shown in Table 6.

Student	Smoker?	Gender	Resting heart rate (bpm)	Heart rate during exercise	Heart rate after exercise
1	Y	Male	60.5	130.0	66.5
2	N	Female	67.0	145.5	73
3	Y	Male	70.0	120	77.0
4	Y	Male	65.5	100	69
5	Y	Male	66.0	128.5	75.5
6	Y	Female	65.5	115.5	74.5
7	Y	Female	73.5	120.5	81
8	N	Female	63.0	118	66
9	N	Female	71.0	95.5	80.5
10	N	Female	65.5	110	71
11	N	Male	64.0	145.5	68
12	N	Male	52.5	140.0	58.5
13	N	Male	54.0	137.5	63
14	N	Female	73.0	130.5	81
15	N	Female	61.5	124	67
16	N	Female	71.0	130	81.5
17	N	Male	60.0	122.5	63
18	N	Female	64.5	118	69
19	Ν	Female	67.5	130.5	73.5
20	Y	Male	72.0	135	82
21	Y	Female	69.5	110	75.5

Table 6

Suggest and explain improvements that the student could make to his experimental method **and** his presentation of data.

In your answer you should explain the benefits of your suggested improvements.

 <b>[6]</b>

16. The student wrote the following summary about the control of heart rate.

When the heart rate is too low the level of carboxylic acid in the blood becomes higher than normal. The vagus nerve sends action potentials to the AVN to increase the contraction rate of the heart muscle. The baroreceptors in the walls of the blood vessels then detect that the pH of the blood is normal, so heart rate can return to resting.

The endocrine system can also change heart rate. Release of the hormone adrenaline from the adrenal medulla causes the smooth muscle of the heart to contract more frequently.

Identify and correct any biological errors in the student's summary.

\_\_\_\_\_

**17.** Phaeochromocytoma is a rare tumour of adrenal gland tissue. It causes increased hormone release from the adrenal glands.

Fig 21.2 shows three ECG traces showing the heart rhythms of three different patients.



- i. Identify patients **X**, **Y** and **Z** by labelling the traces in Fig. 21.2.
- ii. Sketch a trace for a patient who has entered atrial fibrillation.



iii. Suggest why reduced heart rate is sometimes seen in people who are very aerobically fit.

\_\_\_\_\_[2]

[2]

[2]

**18.** Athletic sprinters require large amounts of energy in short periods of time. Many elite sprinters can run 100 metre races in under 10 seconds.

Under normal conditions, exercise requires an increased rate of breathing. It has been observed that some of the best sprinters only take one breath at the start of the race and do not inhale again until the end of the race.

Suggest how these sprinters can expend so much energy without needing to carry out aerobic respiration.

 [2]

## 19.

i. Skeletal muscle is one of the main tissues where glucose is removed from the blood in response to insulin.

Name the other tissue.

\_\_\_\_\_[1]

ii. Explain why glucose is required for the contraction of skeletal muscle.

 [3]

**20.** Prokaryotic cells have cytoskeletons. The molecules in prokaryotic cytoskeletons are different from the molecules in eukaryotic cytoskeletons.

Table 4.1 lists three molecules present in a prokaryotic cytoskeleton.



Table 4.1

## i. Suggest the function of Protein A.

 [1]

 ii. Suggest the function of Protein C.

 [1]

 iii. An antibiotic called A22 binds irreversibly to Protein B. Despite its antibiotic properties, A22 is not used in humans.

 Suggest why scientists have advised that A22 should not be used in humans.

 [1]